Commentary

Paul R. Cooper, M.D., Professor of Neurosurgery, New York University Medical Center

The Role of Vertebrectomy for Degenerative Disease of the Cervical Spine

In this issue of Neurosurgical Focus™, Naderi, et al., in their paper "Cervical spondylotic myelopathy treated with corpectomy" document their experience with this procedure for the treatment of patients with cervical spondylotic myelopathy (CSM). Although vertebrectomy is an important therapeutic option in the management of compressive cervical myelopathy, this article raises as many questions as it answers. A more critical evaluation of the indications for vertebrectomy and its place in the management of CSM might be useful.

Prior to the widespread use of anterior cervical plating techniques, bone grafts that spanned one or more vertebral body levels had to be fitted securely to the recipient site or patients had to be placed in a halo vest to minimize the incidence of graft complications. Graft movement, extrusion, or collapse resulted in a disturbing incidence of kyphotic deformity and necessitated reoperation, even with the most meticulous operative techniques, combined with secure external fixation. The widespread use of anterior cervical plating has resulted in an increase in the use of vertebrectomy, a reduction in the incidence of graft complications, and, for the most part, obviated the use of the halo vest.

Multilevel vertebrectomy combined with anterior cervical plating represents an advantage in the management of patients with ossification of the posterior longitudinal ligament needing extensive decompression. This technique is also useful in the treatment of patients requiring vertebrectomy for single or multilevel metastatic disease, osteomyelitis, or trauma.

The indications for vertebrectomy and its efficacy in patients with multilevel CSM are, in this reviewer's opinion, more controversial. Neurological deficit in patients with CSM at one or more levels results from spinal cord compression by osteophytes that form at or adjacent to the disc space. Although spinal cord compression and neurological deficit from spondylosis are more likely to occur in patients with congenitally narrow spinal canals, a small spinal canal without other pathology is well tolerated.

Because the spondylotic process occurs contiguous to the disc space, removal of the degenerated disc and adjacent osteophyte will result in decompression of the spinal cord and amelioration of neurological deficit. A clear idea of the two-dimensional anatomy of the osteophytic compression is essential for effective decompression. Contiguous computerized tomography slices with appropriate windows will define this anatomy. However, at the time of operation the surgeon must be able to correlate the extent of the decompression he has achieved with the location of osteophytes on imaging studies. It is likely that many patients have not achieved the expected benefits of anterior cervical decompression because of failure to completely remove the compressive lesion. Nevertheless, in skilled hands, removal of 18 to 20 mm of bone and osteophyte in a transverse direction and 10 to 12 mm in a rostrocaudal direction will

adequately decompress the spinal cord.

When osteophytes are present at a single intervertebral disc space, satisfactory decompression can be achieved without resorting to vertebrectomy. Even with the most aggressive decompression there will be enough vertebral body left to accept a bone graft. Although anterior cervical plating has been used in this situation, the advantages it provides, if any, are unclear.

The choice of the most effective operative strategy becomes more difficult when spondylotic compression occurs at two or more adjacent disc spaces. In this situation proponents of vertebrectomy claim that only one bone graft, rather than two or more, can be used with a lower incidence of pseudoarthrosis; resection of a residual vertebral body whose rostral and caudal ends have already been removed makes bone grafting easier. Anterior cervical plating then becomes desirable to minimize the incidence of complications common with the use of a long bone graft.

Nevertheless anterior cervical plating decreases, but does not eliminate the incidence of graft-related complications and is associated with its own problems: plate fracture, screw breakage or back-out, and misplacement of screws. Recent experience would indicate that the longer the length of the plated segment and the greater the instability, the more likely it is that plate complications will occur. If this is so, one might reasonably ask if vertebrectomy and attendant plating are preferable to more conventional decompression and bone grafting at multiple levels.

In fact decompression at adjacent levels without removal of the residual uninvolved vertebral body will adequately treat the cause of spinal cord compression and leave enough vertebral body to accept a bone graft in most patients. Although there may be a higher incidence of pseudoarthrosis when bone grafts are placed at multiple levels compared to single-level fusion, failure of fusion to take place does not always cause symptoms and the incidence may be no higher than the rate of nonfusion with long grafts placed after the vertebrectomy. Moreover, decompression at multiple levels had the advantage of avoiding the rather significant incidence of complications documented by Naderi, et al., in their article.

It is important that the spinal surgeon be prepared to use the surgical technique that is most appropriate to the morphology of the patient's degenerative process. For the most part, complete decompression may be achieved by decompression and grafting at multiple levels without resorting to vertebrectomy. In those less common instances in which imaging studies show the spondylosis at adjacent levels to involve most of the posterior aspect of the vertebral body, vertebrectomy and plating are appropriate and should be the operative procedure of choice.

[Click here to return to article 5.](#)