Neuroimaging findings in pediatric sports-related concussion

Michael J. Ellis, MD,^{1,2,4,6,7} Jeff Leiter, PhD,^{1,6} Thomas Hall, BSc,⁶ Patrick J. McDonald, MD, MHSc,^{1,2,4,6,7} Scott Sawyer, MD,^{2,5} Norm Silver, MD,^{2,5} Martin Bunge, MD,³ and Marco Essig, MD³

Departments of ¹Surgery, ²Pediatrics and Child Health, and ³Diagnostic Imaging and ⁴Section of Neurosurgery, University of Manitoba, ⁵Health Sciences Centre Children's Hospital, ⁶Pan Am Clinic, and ⁷Children's Hospital Research Institute of Manitoba, Canada North Concussion Network, Winnipeg, Manitoba, Canada

OBJECT The goal in this review was to summarize the results of clinical neuroimaging studies performed in patients with sports-related concussion (SRC) who were referred to a multidisciplinary pediatric concussion program.

METHODS The authors conducted a retrospective review of medical records and neuroimaging findings for all patients referred to a multidisciplinary pediatric concussion program between September 2013 and July 2014. Inclusion criteria were as follows: 1) age $\leq$ 19 years; and 2) physician-diagnosed SRC. All patients underwent evaluation and follow-up by the same neurosurgeon. The 2 outcomes examined in this review were the frequency of neuroimaging studies performed in this population (including CT and MRI) and the findings of those studies. Clinical indications for neuroimaging and the impact of neuroimaging findings on clinical decision making were summarized where available. This investigation was approved by the local institutional ethics review board.

RESULTS A total of 151 patients (mean age 14 years, 59% female) were included in this study. Overall, 36 patients (24%) underwent neuroimaging studies, the results of which were normal in 78% of cases. Sixteen percent of patients underwent CT imaging; results were normal in 79% of cases. Abnormal CT findings included the following: arachnoid cyst (1 patient), skull fracture (2 patients), suspected intracranial hemorrhage (1 patient), and suspected hemorrhage into an arachnoid cyst (1 patient). Eleven percent of patients underwent MRI; results were normal in 75% of cases. Abnormal MRI findings included the following: intraparenchymal hemorrhage and sylvian fissure arachnoid cyst (1 patient); non-hemorrhagic contusion (1 patient); demyelinating disease (1 patient); and posterior fossa arachnoid cyst, cerebellar volume loss, and nonspecific white matter changes (1 patient).

CONCLUSIONS Results of clinical neuroimaging studies are normal in the majority of pediatric patients with SRC. However, in selected cases neuroimaging can provide information that impacts decision making about return to play and retirement from the sport.

<http://thejns.org/doi/abs/10.3171/2015.1.PEDS14510>

KEY WORDS neuroimaging; sports-related concussion; pediatric patients; computed tomography; magnetic resonance imaging; trauma

SPORTS-RELATED concussion (SRC) is a form of traumatic brain injury (TBI) representing an emerging public health concern that affects approximately 1.6–3.8 million Americans²⁶ annually, with a significant proportion occurring in children and adolescents. The clinical presentation of acute SRC among pediatric patients is highly variable, often manifesting as a combination of physical, psychological, sleep, and affective symptoms. The vast majority of these patients will achieve

complete neurological recovery within 1–4 weeks; however, 15%–20% will develop postconcussion syndrome, a condition characterized by persistent symptoms lasting more than 1–3 months.^{13,17,34,40} Some consensus statements suggest that the mainstay of management for children and adolescents with SRC is physical and cognitive rest followed by gradual reintegration into full sport and school activities.^{19,22,31–33}

Evidence from experimental animal studies suggests

ABBREVIATIONS GRE = gradient recalled echo; ICH = intracranial hemorrhage; IPH = intraparenchymal hemorrhage; LOC = loss of consciousness; mTBI = mild traumatic brain injury; SRC = sports-related concussion; SWI = susceptibility-weighted imaging.

SUBMITTED September 26, 2014. **ACCEPTED** January 12, 2015.

INCLUDE WHEN CITING Published online June 2, 2015; DOI: 10.3171/2015.1.PEDS14510.

DISCLOSURE The authors report no conflict of interest concerning the materials or methods used in this study or the findings specified in this paper. The Pan Am Clinic Foundation provided financial support for this study.

that SRC results in temporary alterations in neuronal energy metabolism, excitatory neurotransmitter release, and cerebral blood flow,^{12,18} constituting a functional rather than a structural brain injury. Operating on this evidence, various national and international organizations have published position statements in which they concluded that neuroimaging findings are normal in patients with SRC and that these studies add little to the clinical management of this condition.^{4,19,31–33} At present, experts recommend that neuroimaging be used only in acutely injured patients in whom there is suspicion of serious intracranial pathological conditions such as subdural hematoma, epidural hematoma, or intraparenchymal hemorrhage (IPH), or in patients with prolonged disturbance in level of consciousness, focal neurological deficits, or worsening symptoms.^{4,19,21,22,31–33} Despite significant consistency across published position statements,^{4,19,21,22,31–33} these recommendations are derived from expert opinion only (Level 5 evidence).

In recent years, several advances in structural neuroimaging techniques have led to improved detection of parenchymal abnormalities associated with TBI. Susceptibility-weighted imaging (SWI) and gradient recalled echo (GRE) sequences that are exquisitely sensitive to the magnetic properties of blood have permitted the visualization of traumatic axonal injury and cerebral microhemorrhages in pediatric patients with TBI.^{2,5–7,41,42} Although preliminary research studies have failed to detect similar abnormalities in pediatric patients with SRC²⁹ and mild TBI (mTBI),^{30,43} research studies often include subjects with no clinically defined indication for neuroimaging. Although previous neuroimaging studies of mTBI in children and adolescents have included patients participating in sports,^{23,30,36,41} we are not aware of any published studies that have examined clinical neuroimaging findings among an exclusive cohort of pediatric patients with SRC. Furthermore, there are currently no evidence-based return-to-play or retirement practice guidelines for children and adolescents with SRC who demonstrate traumatic lesions on clinical neuroimaging studies.

Accordingly, the purpose of this retrospective review was to summarize the results of clinical neuroimaging studies performed in patients with SRC who were referred to and treated at a multidisciplinary pediatric concussion program. The impact of neuroimaging findings on clinical decision making is discussed.

Methods

Research Design

We conducted a single-center, retrospective review of medical records and neuroimaging findings for all patients referred to the Pan Am Concussion Program at the Pan Am Clinic in Winnipeg, Manitoba, between September 1, 2013, and July 31, 2014. Patient referrals were obtained from pediatric and adult emergency departments, pediatricians, family practitioners, athletic therapists, and coaches. Inclusion criteria were as follows: 1) age $\leq$ 19 years; and 2) neurosurgeon-diagnosed SRC. Exclusion criteria included 1) diagnosed concussion occurring outside of a sporting activity; 2) incomplete medical records; and 3) neuroimaging studies that were undertaken as a part

of research studies. Concussions sustained during nonspecific gym class and schoolyard activities were excluded. This study was approved by the institutional ethics review board at the University of Manitoba.

Definitions of Terms

The diagnosis of SRC was determined during initial consultation for all eligible patients by a single neurosurgeon with fellowship training in concussion spectrum disorders, according to the definition set forth by the International Consensus on Concussion in Sport.³³ Because patients in the study were referred by numerous outside emergency department, pediatric, and family medicine physicians, it was not possible to confirm the diagnostic criteria used initially to refer patients to the Pan Am Concussion Program. Clinical indications for neuroimaging performed prior to referral were also absent for the vast majority of patients. In the case of those patients who underwent neuroimaging while under the care of the Pan Am Concussion Program, all studies were ordered by the same neurosurgeon, based on the patient's clinical history and physical examination findings, and were summarized accordingly. In general, MRI was considered in patients who demonstrated focal neurological deficits or persistent symptoms lasting more than 1–2 months that were not improving, or patients who were referred with abnormal findings on CT imaging studies.

Assessments of SRC

Neuroimaging studies were performed at multiple hospitals by using multiple CT and MRI scanners. All CT scans performed in this study were obtained without contrast. Magnetic resonance imaging protocols also differed across imaging sites but were summarized accordingly. All neuroimaging studies in this cohort were performed without sedation or general anesthetic.

Statistical Analysis

Outcomes examined in this review were the frequency of neuroimaging studies performed and their findings. All data related to patient demographic information, neuroimaging studies performed, and neuroimaging outcomes were summarized using descriptive statistics. All statistical analyses were performed using SPSS 22.0 (IBM Corp.).

Results

During the study period, 212 pediatric patients were referred to the multidisciplinary concussion program. Of those patients, 151 (71%; 65 male and 86 female) received a diagnosis of SRC on initial consultation by the neurosurgeon and were included in the study. A summary of sports played by pediatric patients with SRC is presented in Table 1. Sixty-one patients were excluded due to concussions sustained outside of a sport setting. The mean age of patients in this cohort was 14.26 ± 2.21 years (range 5–19 years). Twenty-four of 151 patients (16%) underwent CT imaging, all of which occurred in the emergency department, with 23 patients undergoing imaging prior to referral to the concussion program and 1 undergoing urgent

TABLE 1. Summary of sports played by 151 pediatric patients with SRC

Sport	% of Patients*
Hockey	41.72
Soccer	20.53
Basketball	7.95
Football	7.28
Baseball	4.64
Skiing/snowboarding	3.97
Ringette	2.65
Volleyball	2.65
Rugby	1.99
Other	6.62

* Less than 2% of patients participated in other sporting activities such as figure skating/ice skating, wrestling, dance/gymnastics, swimming, cycling, trampoline, and track and field.

CT after a syncopal episode at home that occurred after consultation with the concussion program. The results of cranial CT imaging were normal in 19 of 24 (79%) cases, including a patient with a normal anatomical variant of the septum pellucidum. Abnormal CT imaging results included calvarial skull fracture (1 patient), orbital floor fracture (1 patient), posterior fossa arachnoid cyst (1 patient), focal hyperdensity in the basal ganglia suspicious for hemorrhage (1 patient), and isodense lesion in the left sylvian fissure suspicious for hemorrhage into a small sylvian fissure arachnoid cyst (1 patient). There were no cases of definitive intracranial hemorrhage (ICH) observed on CT imaging in this study.

Sixteen of 151 patients (11%) underwent MRI, including 4 patients who also underwent CT imaging. All MRI studies were performed in the outpatient setting, except 1 performed in a hospitalized patient. All MRI studies were obtained following referral to the concussion program, except 1 obtained prior to referral. Clinical indications for MRI in this study include the following: persistent symptoms (12 patients), focal neurological deficit (1 patient), abnormal CT imaging findings (2 patients), and constitutional symptoms (1 patient). The results of cranial MRI were normal in 12 of 16 (75%) patients. One patient was found to have a cavum septum pellucidum that was classified as a normal finding. Another was found to have bilateral basal ganglia calcifications that were also classified as a normal finding. Abnormal MRI results included the following: IPH and a sylvian fissure arachnoid cyst (1 patient); nonhemorrhagic contusion (1 patient); multifocal demyelinating disease (1 patient); and posterior fossa arachnoid cyst, cerebellar volume loss, and nonspecific white matter changes (1 patient). A summary of patients who underwent MRI is shown in Table 2.

Illustrative Cases

Case 1

An 11-year-old boy sustained a brief loss of consciousness (LOC) after a fall onto the ice during a competitive hockey game. He was previously healthy with no history

of concussions or TBI. His family history was significant for a second-generation relative who played professional American football and in whom chronic traumatic encephalopathy was diagnosed on autopsy. The patient was evaluated at a pediatric emergency department, where he complained of headache, nausea, photophobia, and irritability. A CT scan of the brain obtained without contrast demonstrated an extraaxial isodense lesion within the left sylvian fissure that was suspicious for hemorrhage into a small arachnoid cyst. The patient was referred to the multidisciplinary concussion program, where he reported persistent concussion symptoms, including headache, dizziness, and postural imbalance. An MRI study of the brain obtained without contrast was performed 41 days after the injury and confirmed a small left sylvian fissure arachnoid cyst with no evidence of hemorrhage. The GRE imaging revealed a 2.3×1 -cm curvilinear hypointense lesion within the right occipital lobe white matter consistent with an IPH that was not observed on any other MR sequences (Fig. 1). The patient remained symptomatic at 144 days postinjury. Based on the traumatic abnormalities identified on clinical neuroimaging findings and the child's prolonged symptoms, the neurosurgeon advised the parents that in his opinion the child should not return to contact sports.

Case 2

During a ringette game, a 14-year-old girl sustained a fall onto the ice that was not associated with any LOC or amnesia. She was previously healthy, with 1 prior questionable concussion and no history of moderate or severe TBI. She initially became asymptomatic at rest within a few days postinjury but experienced severe headaches on repeated attempts to return to ringette. Two months later, she underwent an MRI study obtained without contrast that demonstrated a left parietooccipital abnormality that was best visualized on T2-weighted and FLAIR sequences (Fig. 2). No evidence of hemorrhage was observed on GRE imaging. Follow-up contrast-enhanced MRI studies performed 7 and 11 months later demonstrated no interval growth of the lesion, making other differential diagnoses such as low-grade glioma less likely. Single-voxel MR spectroscopy performed over the region of interest demonstrated a slightly decreased creatine level relative to choline and *N*-acetylaspartate levels, which correlated with gliosis. Taken together, the neuroimaging features were most consistent with the presence of a nonhemorrhagic parenchymal contusion or gliosis. Formal neuropsychological testing undertaken at 14 months postinjury demonstrated impairments in visual memory, sensory perception, and visuospatial processing. The patient, who was asymptomatic at rest and during physical activity, was referred to the multidisciplinary concussion program for an opinion regarding return to play. Based on the traumatic abnormalities identified on clinical neuroimaging and concordant neuropsychological findings, the neurosurgeon advised the parents that in his opinion the child should not return to contact sports.

Discussion

This review provides preliminary empirical evidence

TABLE 2. Summary of pediatric patients with SRC who underwent MRI

Age (yrs), Sex	Sport	Symptoms	Clinical Indication	Time Since Injury	MRI Details	MRI Results
11, M*	Hockey	HA	Abnormal CT results	41 days	GRE+, C-, SWI-	Lt sylvian fissure arachnoid cyst, rt occipital IPH
17, F	Soccer	HA, polydipsia, cold intolerance	Constitutional symptoms	40 days	C+, GRE-, SWI+, dynamic pituitary study	Normal
17, F	Hockey	HA, blurred vision, dizziness	Persistent symptoms	238 days	GRE+, C-, SWI-	Normal
14, F	Soccer	HA, photophobia, dizziness	Persistent symptoms	183 days	GRE+, C-, SWI-	Normal, cavum septum pellucidum
14, M	Hockey	Blurred vision, HA, vomiting	Focal neuro deficit	9 days	GRE+, C+, SWI-	Multifocal demyelinating disease, optic neuritis
14, F*	Ringette	HA	Persistent symptoms	60 days	GRE+, C+, SWI-	Lt parietal nonhemorrhagic contusion
14, F	Hockey	HA, fatigue, sleep dist	Persistent symptoms	176 days	GRE-, C-, SWI-	Normal
18, F	Hockey	HA, dizziness, visual dist	Persistent symptoms	170 days	GRE+, TOF MRA	Normal
13, F	Gymnastics	HA, fatigue, sleep dist	Abnormal CT results	14 days	GRE+, C+, MRA	Normal, bilat basal ganglia calcifications
11, M	Hockey	HA, dizziness, photophobia	Persistent symptoms	154 days	GRE-, SWI+	Normal
14, M	Volleyball	HA, decreased concentration, language processing impairment	Persistent symptoms	180 days	GRE+, C-, SWI-	Normal
15, F	Basketball	HA, blurred vision, anxiety	Persistent symptoms	213 days	GRE+, C-, SWI-	Normal
14, M	Hockey	Migraine HA	Persistent symptoms	131 days	GRE+, C-, SWI+	Normal
16, F	Soccer	HA, blurred vision, dizziness	Persistent symptoms	143 days	GRE+, C-, SWI-	Normal
15, F	Baseball	HA	Persistent symptoms	199 days	GRE-, C-, SWI-	PF arachnoid cyst, vol loss in Lt cerebellum, nonspecific WM abnormalities
14, F	Figure skating	HA, dizziness, anxiety	Persistent symptoms	210 days	GRE+, C-, SWI-	Normal

C+, C- = study obtained with (+) or without (-) contrast; dist = disturbance; GRE+, GRE- = GRE imaging included (+) or not included (-); HA = headache; MRA = MR angiography; neuro = neurological; PF = posterior fossa; SWI+, SWI- = susceptibility-weighted imaging included (+) or not included (-); TOF = time of flight; WM = white matter.

* Illustrative cases in text.

to suggest that the results of clinical neuroimaging studies performed in children and adolescents with SRC are normal in the vast majority of cases. Overall, the results in 78% of neuroimaging studies were normal in this retrospective review of patients referred to a multidisciplinary concussion program. The results of CT imaging studies performed exclusively in the emergency department were normal in 79% of cases. Clinically indicated MRI studies performed almost exclusively in the outpatient setting yielded normal findings in 75% of cases. Among the patients who underwent neuroimaging studies, traumatic abnormalities were detected in 11% of cases.

Over the past decade, numerous published position statements from national and international organizations have concluded that neuroimaging studies yield normal findings in the vast majority of patients with SRC, and thus neuroimaging contributes little to the clinical management of these patients.^{4,21,31-33} Experts suggest that neuroimaging be used only to rule out structural intracranial pathology in patients with impaired level of consciousness, focal neurological deficits, or persistent symptoms.^{4,19,21,22,31-33} Although such recommendations seem reasonable, there

are no clinical studies that have examined neuroimaging findings in an exclusive population of children and adolescents with SRC. Consequently, these expert findings and recommendations are based on expert opinion only (Level 5 evidence).

The use of CT imaging in pediatric mTBI has been the focus of several multiinstitutional clinical studies. Despite recent widespread increases in the use of CT in the evaluation of pediatric mTBI in Canadian and American emergency departments,²³ only 4%–7% of patients are found to demonstrate an abnormality on CT, with only 0.5% requiring neurosurgical intervention.^{15,23} Head injury occurring during a sporting activity has consistently failed to be identified as a risk factor for an abnormal CT finding across pediatric studies, which includes other risk factors such as low Glasgow Coma Scale score; suspected open, depressed, or basal skull fractures; persistent headache or irritability; large or boggy scalp hematoma; and dangerous mechanism of injury (i.e., motor vehicle accident, fall from bicycle with no helmet).^{3,16,24,27,35-37} These studies have informed the development of several clinical decision rules that help guide the judicious use of CT imaging

FIG. 1. Illustrative Case 1. Axial GRE image (**left**) demonstrates curvilinear hypointensity consistent with right occipital IPH that was not visualized on axial T2-weighted MRI (**right**).

FIG. 2. Illustrative Case 2. Axial T2-weighted (**left**) and FLAIR (**right**) MRI studies demonstrate evidence of left parietal nonhemorrhagic contusion.

in pediatric patients with mTBI who are evaluated in the emergency department.^{3,16,24,27,35–37}

In the present study, 21% of CT imaging studies performed by emergency department physicians in patients with suspected SRC demonstrated abnormal findings, including calvarial skull fracture (1 patient), orbital floor fracture (1 patient), suspected ICH (2 patients), and posterior fossa arachnoid cyst (1 patient). One patient was found to have a normal anatomical variant of the septum pellucidum, including a cavum septum pellucidum and a cavum vergae, which was classified as a normal finding and is present in up to 20% of the general pediatric population on neuroimaging.⁸ Overall, the results of this study and review of the literature suggest that CT imaging findings are normal in the vast majority of pediatric patients with SRC and mTBI, and that this modality should be reserved for use only in the emergency department to rule out ICH or skull fractures in acutely injured patients with appropriate clinical signs and symptoms. Because of concerns related to the oncological and neurocognitive risks associated with exposure to low doses of ionizing radiation in children,^{9,20} the use of CT outside of the emergency department setting is discouraged.

The use of MRI in pediatric SRC has received significant attention among researchers in recent years. Research series suggest that findings on conventional structural MRI studies are also normal in the vast majority of pediatric patients with mild TBI and SRC;^{29,30,43} however, research cohorts frequently include subjects who are recruited out of convenience and may not have a clinical indication for MRI. In such cases, information is lacking regarding clinically driven MRI findings in pediatric patients with SRC. Abnormal findings were detected in 25% of MRI studies performed in this series, with 2 MR images demonstrating traumatic abnormalities and 3 contributing directly to clinical decision making. In 1 case, MRI undertaken for blurred vision and vomiting allowed the timely diagnosis of a potentially life-threatening demyelinating disease and permitted urgent treatment resulting in complete neurological recovery and return to sporting activities. In 2 cases, MRI demonstrated hemorrhagic and nonhemorrhagic traumatic lesions that impacted decision making about return to play.

These cases highlight the need to consider MRI in pediatric patients with focal neurological deficits, worrisome symptoms, or abnormal or inconclusive CT findings. MRI should also be considered in pediatric patients with persistent symptoms for which the definition is unclear. The vast majority of children and adolescents reach symptomatic recovery within 1–4 weeks, although approximately 15%–20% will remain symptomatic beyond 1 month.^{13,17,34,40} Risk factors for prolonged recovery may include female sex, LOC, and a history of concussions, attention deficit hyperactivity disorder, migraine headaches, and learning disorders.^{25,28,39} Physicians caring for pediatric patients with SRC must take into account all of these factors as well as the trajectory of concussion symptoms and physical examination findings when making the decision to order neuroimaging. These decisions are also impacted by the clinical experience and training of the treating physician as well as his or her familiarity with neurological conditions that can potentially mimic or can also co-occur in the setting of SRC. Such complexities underscore the value of multidisciplinary pediatric concussion programs that have the clinical neurological expertise and diagnostic resources to comprehensively address patients with SRC who have persistent symptoms.

At present, there are limited practice guidelines to direct clinical SRC decision making in children and adolescents who are found to have traumatic lesions on neuroimaging. Although there is some literature to suggest that patients with abnormalities on CT or MRI studies should be advised to retire from contact sports,^{10,11} these recommendations do not specifically address recent advances in SWI and GRE imaging that are capable of demonstrating traumatic abnormalities that would have otherwise escaped detection with conventional MRI studies. Whether the presence of traumatic lesions on neuroimaging following SRC is a reflection of an unrecognized genetic susceptibility to head injury is not known. Based on the available literature and guidance by a second opinion from another neurosurgeon, the decision was made to advise both patients who sustained traumatic lesions detected on MRI in this study to retire from contact sports.

Three patients (2 boys and 1 girl) in this study were found to have arachnoid cysts, 1 located in the sylvian fis-

sure (found on CT and MRI) and 2 in the cerebellopontine angle of the posterior fossa (1 found on MRI and 1 found on CT). In a large, single-institution series of neuroimaging studies performed in children and adolescents, the prevalence rate for arachnoid cysts was 2.5%, with these abnormalities found more commonly in male patients.¹ Because of the potential risk of intracranial cyst hemorrhage as well as subdural hematoma and hygroma that can occur following minor head injury,^{14,38} all patients with incidentally discovered arachnoid cysts should undergo consultation with a neurosurgeon to discuss participation in contact sports. In the case of the 2 patients with incidentally discovered posterior fossa arachnoid cysts, both lesions were small with no significant mass effect, and thus both patients were able to return to full sports participation without restriction.

The findings of this study must be evaluated in light of several important limitations. First, patients referred to a multidisciplinary pediatric concussion program may include those with more severe injuries and may thus constitute a population for whom neuroimaging is more likely to be indicated, and with potentially abnormal results, compared with the general pediatric SRC population. Second, patients with traumatic abnormalities identified on initial CT scans following sports-related head trauma may have been referred to other neurosurgeons and not the concussion program, thus reducing the proportion of children with these findings observed in this clinical population. Third, because the majority of CT imaging was performed prior to patient referral to the concussion program, it was not possible to assess whether patients had evidence-based clinical indications for CT imaging. Fourth, the decision to perform MRI was not based on strict predetermined criteria but rather it was based almost entirely on the clinical judgment of a single neurosurgeon. Because of this selection bias, the findings of this study cannot be used to estimate the risk of abnormal MRI findings among a general population of pediatric patients with SRC. Finally, there was considerable variation in MRI protocols across institutions, with not all studies using recent technological advances in SWI or GRE imaging. Therefore, it remains unclear whether standardized incorporation of these sequences in MR imaging protocols would have led to greater detection of cerebral microhemorrhages in this clinical population. To address these limitations, the frequency of abnormal results on neuroimaging in pediatric patients with SRC who have persistent concussion symptoms, focal neurological findings, and an impaired level of consciousness will be addressed by a future prospective study at our institution incorporating these novel structural neuroimaging techniques.

Conclusions

Preliminary empirical evidence suggests that results of clinical neuroimaging studies are normal in the majority of pediatric patients with SRC. However, in selected cases neuroimaging can provide information that impacts decision making about return to play and retirement from the sport. Evidence-based practice guidelines are urgently needed to help direct clinical decision making when neuroimaging studies demonstrate evidence of traumatic intracranial injury.

Acknowledgment

We thank the Pan Am Clinic Foundation for their financial support of this study.

References

1. Al-Holou WN, Yew AY, Boomsaad ZE, Garton HJ, Muraszko KM, Maher CO: Prevalence and natural history of arachnoid cysts in children. *J Neurosurg Pediatr* 5:578–585, 2010
2. Ashwal S, Babikian T, Gardner-Nichols J, Freier MC, Tong KA, Holshouser BA: Susceptibility-weighted imaging and proton magnetic resonance spectroscopy in assessment of outcome after pediatric traumatic brain injury. *Arch Phys Med Rehabil* 87 (12 Suppl 2):S50–S58, 2006
3. Atabaki SM, Stiell IG, Bazarian JJ, Sadow KE, Vu TT, Camarca MA, et al: A clinical decision rule for cranial computed tomography in minor pediatric head trauma. *Arch Pediatr Adolesc Med* 162:439–445, 2008
4. Aubry M, Cantu R, Dvorak J, Graf-Baumann T, Johnston K, Kelly J, et al: Summary and agreement statement of the First International Conference on Concussion in Sport, Vienna 2001. Recommendations for the improvement of safety and health of athletes who may suffer concussive injuries. *Br J Sports Med* 36:6–10, 2002
5. Babikian T, Freier MC, Tong KA, Nickerson JP, Wall CJ, Holshouser BA, et al: Susceptibility weighted imaging: neuropsychologic outcome and pediatric head injury. *Pediatr Neurol* 33:184–194, 2005
6. Beauchamp MH, Beare R, Ditchfield M, Coleman L, Babl FE, Kean M, et al: Susceptibility weighted imaging and its relationship to outcome after pediatric traumatic brain injury. *Cortex* 49:591–598, 2013
7. Beauchamp MH, Ditchfield M, Babl FE, Kean M, Catroppa C, Yeates KO, et al: Detecting traumatic brain lesions in children: CT versus MRI versus susceptibility weighted imaging (SWI). *J Neurotrauma* 28:915–927, 2011
8. Bodensteiner JB, Schaefer GB, Craft JM: Cavum septi pellucidi and cavum vergae in normal and developmentally delayed populations. *J Child Neurol* 13:120–121, 1998
9. Brenner D, Elliston C, Hall E, Berdon W: Estimated risks of radiation-induced fatal cancer from pediatric CT. *AJR Am J Roentgenol* 176:289–296, 2001
10. Cantu RC: Recurrent athletic head injury: risks and when to retire. *Clin Sports Med* 22:593–603, x, 2003
11. Cantu RC, Register-Mihalik JK: Considerations for return-to-play and retirement decisions after concussion. *PM R* 3 (10 Suppl 2):S440–S444, 2011
12. Choe MC, Babikian T, DiFiori J, Hovda DA, Giza CC: A pediatric perspective on concussion pathophysiology. *Curr Opin Pediatr* 24:689–695, 2012
13. Collins M, Lovell MR, Iverson GL, Ide T, Maroon J: Examining concussion rates and return to play in high school football players wearing newer helmet technology: a three-year prospective cohort study. *Neurosurgery* 58:275–286, 2006
14. Cress M, Kestle JR, Holubkov R, Riva-Cambrin J: Risk factors for pediatric arachnoid cyst rupture/hemorrhage: a case-control study. *Neurosurgery* 72:716–722, 2013
15. Davis RL, Mullen N, Makela M, Taylor JA, Cohen W, Rivara FP: Cranial computed tomography scans in children after minimal head injury with loss of consciousness. *Ann Emerg Med* 24:640–645, 1994
16. Dunning J, Daly JP, Lomas JP, Lecky F, Batchelor J, Mackway-Jones K: Derivation of the children's head injury algorithm for the prediction of important clinical events decision rule for head injury in children. *Arch Dis Child* 91:885–891, 2006
17. Eisenberg MA, Meehan WP III, Mannix R: Duration and course of post-concussive symptoms. *Pediatrics* 133:999–1006, 2014

18. Giza CC, Hovda DA: The neurometabolic cascade of concussion. **J Athl Train** 36:228–235, 2001
19. Giza CC, Kutcher JS, Ashwal S, Barth J, Getchius TS, Gioia GA, et al: Summary of evidence-based guideline update: evaluation and management of concussion in sports: report of the Guideline Development Subcommittee of the American Academy of Neurology. **Neurology** 80:2250–2257, 2013
20. Hall P, Adami HO, Trichopoulos D, Pedersen NL, Lagiou P, Ekblom A, et al: Effect of low doses of ionising radiation in infancy on cognitive function in adulthood: Swedish population based cohort study. **BMJ** 328:19, 2004
21. Halstead ME, Walter KD: Clinical report—sport-related concussion in children and adolescents. **Pediatrics** 126:597–615, 2010
22. Harmon KG, Drezner JA, Gammons M, Guskiewicz KM, Halstead M, Herring SA, et al: American Medical Society for Sports Medicine position statement: concussion in sport. **Br J Sports Med** 47:15–26, 2013
23. Klassen TP, Reed MH, Stiell IG, Nijssen-Jordan C, Tenenbein M, Joubert G, et al: Variation in utilization of computed tomography scanning for the investigation of minor head trauma in children: a Canadian experience. **Acad Emerg Med** 7:739–744, 2000
24. Kuppermann N, Holmes JF, Dayan PS, Hoyle JD Jr, Atabaki SM, Holubkov R, et al: Identification of children at very low risk of clinically-important brain injuries after head trauma: a prospective cohort study. **Lancet** 374:1160–1170, 2009
25. Kutcher JS, Eckner JT: At-risk populations in sports-related concussion. **Curr Sports Med Rep** 9:16–20, 2010
26. Langlois JA, Rutland-Brown W, Wald MM: The epidemiology and impact of traumatic brain injury: a brief overview. **J Head Trauma Rehabil** 21:375–378, 2006
27. Maguire JL, Boutis K, Ulerik EM, Laupacis A, Parkin PC: Should a head-injured child receive a head CT scan? A systematic review of clinical prediction rules. **Pediatrics** 124:e145–e154, 2009
28. Makdissi M, Davis G, Jordan B, Patricios J, Purcell L, Putukian M: Revisiting the modifiers: how should the evaluation and management of acute concussions differ in specific groups? **Br J Sports Med** 47:314–320, 2013
29. Maugans TA, Farley C, Altaye M, Leach J, Cecil KM: Pediatric sports-related concussion produces cerebral blood flow alterations. **Pediatrics** 129:28–37, 2012
30. Mayer AR, Ling JM, Yang Z, Pena A, Yeo RA, Klimaj S: Diffusion abnormalities in pediatric mild traumatic brain injury. **J Neurosci** 32:17961–17969, 2012
31. McCrory P, Johnston K, Meeuwisse W, Aubry M, Cantu R, Dvorak J, et al: Summary and agreement statement of the 2nd International Conference on Concussion in Sport, Prague 2004. **Br J Sports Med** 39:196–204, 2005
32. McCrory P, Meeuwisse W, Johnston K, Dvorak J, Aubry M, Molloy M, et al: Consensus Statement on Concussion in Sport: the 3rd International Conference on Concussion in Sport held in Zurich, November 2008. **Br J Sports Med** 43 (Suppl 1):i76–i90, 2009
33. McCrory P, Meeuwisse WH, Aubry M, Cantu B, Dvorák J, Echemendia RJ, et al: Consensus statement on concussion in sport: the 4th International Conference on Concussion in Sport held in Zurich, November 2012. **Br J Sports Med** 47:250–258, 2013
34. Meehan WP III, Mannix RC, Straccioli A, Elbin RJ, Collins MW: Symptom severity predicts prolonged recovery after sport-related concussion, but age and amnesia do not. **J Pediatr** 163:721–725, 2013
35. Oman JA, Cooper RJ, Holmes JF, Viccellio P, Nyce A, Ross SE, et al: Performance of a decision rule to predict need for computed tomography among children with blunt head trauma. **Pediatrics** 117:e238–e246, 2006
36. Osmond MH, Klassen TP, Wells GA, Correll R, Jarvis A, Joubert G, et al: CATCH: a clinical decision rule for the use of computed tomography in children with minor head injury. **CMAJ** 182:341–348, 2010
37. Palchak MJ, Holmes JF, Vance CW, Gelber RE, Schauer BA, Harrison MJ, et al: A decision rule for identifying children at low risk for brain injuries after blunt head trauma. **Ann Emerg Med** 42:492–506, 2003
38. Parsch CS, Krauss J, Hofmann E, Meixensberger J, Roosen K: Arachnoid cysts associated with subdural hematomas and hygromas: analysis of 16 cases, long-term follow-up, and review of the literature. **Neurosurgery** 40:483–490, 1997
39. Scopaz KA, Hatzenbuehler JR: Risk modifiers for concussion and prolonged recovery. **Sports Health** 5:537–541, 2013
40. Tator CH: Concussions and their consequences: current diagnosis, management and prevention. **CMAJ** 185:975–979, 2013
41. Tong KA, Ashwal S, Holshouser BA, Nickerson JP, Wall CJ, Shutter LA, et al: Diffuse axonal injury in children: clinical correlation with hemorrhagic lesions. **Ann Neurol** 56:36–50, 2004
42. Tong KA, Ashwal S, Holshouser BA, Shutter LA, Herigault G, Haacke EM, et al: Hemorrhagic shearing lesions in children and adolescents with posttraumatic diffuse axonal injury: improved detection and initial results. **Radiology** 227:332–339, 2003
43. Yang Z, Yeo RA, Pena A, Ling JM, Klimaj S, Campbell R, et al: An fMRI study of auditory orienting and inhibition of return in pediatric mild traumatic brain injury. **J Neurotrauma** 29:2124–2136, 2012

Author Contributions

Conception and design: Ellis, Leiter, McDonald, Sawyer, Silver. Acquisition of data: Hall. Analysis and interpretation of data: Hall, Bunge, Essig. Drafting the article: Ellis. Critically revising the article: all authors. Reviewed submitted version of manuscript: all authors. Approved the final version of the manuscript on behalf of all authors: Ellis. Study supervision: Ellis.

Supplemental Information

Previous Presentation

This work was presented as a poster at the 2014 AANS Pediatric Section meeting, which was held on Amelia Island, FL, on December 2–5, 2014.

Correspondence

Michael Ellis, Pan Am Clinic, 75 Poseidon Bay, Winnipeg, MB R3M 3E4, Canada. email: mellis3@panamclinic.com.