Neurosurgical Forum

LETTERS TO THE EDITOR

Endoscope-assisted hemispherotomy

TO THE EDITOR: We read with interest the article published by Wagner et al.⁶ (Wagner K, Vaz-Guimaraes F, Camstra K, et al: Endoscope-assisted hemispherotomy: translation of technique from cadaveric anatomical feasibility study to clinical implementation. *J Neurosurg Pediatr* 23:178–186, February 2019).

It is indeed heartening to see that the technique we initially developed^{1–4} is now being performed at other centers. Following our paper, Sood et al.⁵ described a similar technique. Wagner's group is the third group to describe this technique. Wagner et al. provided themselves a good learning curve by progressing first from a cadaveric model to treating patients with atrophic hemispheric pathologies. Such a learning curve allows the surgeon to develop surgical skills and also provides them a sense of comfort before treading into new territories. We have the following comments for the authors.

- 1) Corpus callosotomy. The callosotomy performed for hemispherotomy is slightly different from that where the corpus callosotomy alone is performed. The callosotomy as part of the hemispherotomy has to be performed in such a manner so that the ipsilateral ventricle is opened. In cases in which a callosotomy alone is performed, strict midline sectioning is performed, which passes between the septae into the cavum. Such a technique may be disastrous for hemispherotomy, as any injury to septal and midline structures may lead to significant cognitive deterioration (as all cortical functions are supported by single hemisphere and midline structures). This must be emphasized, especially to newer surgeons who are learning this technique.
- 2) Frontobasal or anterior disconnection. We refer to frontobasal disconnection as “anterior disconnection.” It is again important to start the frontobasal disconnection at the junction of the genu and the A₂ segment and pass laterally starting first in the extraventricular region. This disconnection passes anterior to the caudate head, and, as it passes laterally, it reaches the anterior part of the putamen and passes inside the ventricle. Starting the frontobasal (or anterior) disconnection within the ventricle would bring it too close to the anterior perforated substance and may again injure the diencephalic structures. The anterior disconnection as described in our paper passes along the coronal plane.³

This point was not made adequately clear by the authors.

- 3) Temporal horn access. We refer to this step as “middle disconnection,” as we feel that it is a better term. This passes along the parasagittal plane lateral to the putamen and starts where the anterior disconnection has been completed, and by virtue of its plane of disconnection, it also performs the insular disconnection. The middle disconnection disconnects (and excises) the ventral amygdala and temporal stem anteriorly and joins the temporal horn only posteriorly. This is the most challenging component of the surgery and has to be worked both anterior to posterior and vice versa. Thus, this step is not just about temporal horn access.
- 4) Posterior disconnection. This point again was not clearly made by the authors. This is a line that joins the splenium to the choroid plexus and is a short but important disconnection. This effectively disconnects the temporal efferents, including the fornix and the tail of the caudate nucleus. These structures curve over the roof of the temporal horn along the atrium to pass along the medial and floor of the body of the lateral ventricle.

For our center, endoscopic hemispherotomy has become the sole procedure of choice for all pathologies, atrophic or nonatrophic (n = 45 to date). Since a nonvascular plane is followed, there is minimal blood loss. It does entail, however, a learning curve and experience with handling the endoscope. This is one of the main reasons why we use robotic guidance to hold and guide the endoscope.³ Our technique has opened a new era of use of the endoscope in epilepsy surgery.⁴

P. Sarat Chandra, MCh
Manjari Tripathi, DM

All India Institute of Medical Sciences (AIIMS), New Delhi, India

References

1. Baumgartner JE, Blount JP, Blauwblomme T, Chandra PS: Technical descriptions of four hemispherectomy approaches: from the Pediatric Epilepsy Surgery Meeting at Gothenburg 2014. *Epilepsia* 58 Suppl 1:46–55, 2017
2. Chandra PS, Kurwale N, Garg A, Dwivedi R, Malviya SV, Tripathi M: Endoscopy-assisted interhemispheric transcallosal hemispherotomy: preliminary description of a novel technique. *Neurosurgery* 76:485–495, 2015
3. Chandra PS, Subianto H, Bajaj J, Girishan S, Doddamani R, Ramanujam B, et al: Endoscope-assisted (with robotic

guidance and using a hybrid technique) interhemispheric transcallosal hemispherotomy: a comparative study with open hemispherotomy to evaluate efficacy, complications, and outcome. **J Neurosurg Pediatr** 23:187–197, 2018

4. Chandra SP, Tripathi M: Endoscopic epilepsy surgery: emergence of a new procedure. **Neurol India** 63:571–582, 2015
5. Sood S, Marupudi NI, Asano E, Haridas A, Ham SD: Endoscopic corpus callosotomy and hemispherotomy. **J Neurosurg Pediatr** 16:681–686, 2015
6. Wagner K, Vaz-Guimaraes F, Camstra K, Lam S: Endoscope-assisted hemispherotomy: translation of technique from cadaveric anatomical feasibility study to clinical implementation. **J Neurosurg Pediatr** 23:178–186, 2019

Disclosures

The authors report no conflict of interest.

Correspondence

P. Sarat Chandra: saratpchandra3@gmail.com.

INCLUDE WHEN CITING

Published online November 29, 2019; DOI: 10.3171/2019.7.PEDS19410.

Response

No response was received from the authors of the original article.

©AANS 2020, except where prohibited by US copyright law

Fetal closure of myelomeningocele

TO THE EDITOR: We read with great interest the 75th anniversary invited review of fetal surgery for spina bifida by Dewan and Wellons² (Dewan MC, Wellons JC III: Fetal surgery for spina bifida. JNSPG 75th Anniversary Invited Review Article. *J Neurosurg Pediatr* 24:105–114, August 2019). Fetal closure of myelomeningocele (MMC) is a landmark advance that has revolutionized care for these children. The authors are uniquely positioned to review fetal surgery, as their center was one of the original participants in the Management of Myelomeningocele Study (MOMS) and remains one of the busiest programs in North America for intrauterine MMC closure (IUMC). The MOMS trial is one of the few randomized trials to address a topic of pediatric neurosurgical importance. We congratulate these authors, their co-investigators, and the researchers at other fetal surgery centers, and we celebrate ongoing refinement to mitigate risk and define optimal inclusion criteria for fetal surgery. We do, however, have a couple of concerns that we would like to raise and invite the authors' commentary.

While the efficacy of IUMC was demonstrated by the randomized trial, this was a highly selected group of patients, treated at very experienced centers.¹ As such, the external validity remains unproven, and the class I evidence can only be considered to apply for patients meeting inclusion criteria for the original trial. This important nuance is nearly uniformly lost in discussions pertaining to IUMC.

We must also be cautious that enthusiasm for a new technique does not lead us to overstate the potential bene-

fits and that conclusions are closely supported by the data. For example, Dewan and Wellons discuss the rate of shunt revision in children who have undergone fetal versus postnatal surgery. These data were published in this journal by Tulipan et al. in the updated results of the entire MOMS cohort.⁶ In both the original publication and the current review, a statistically significant difference in shunt revision rate is reported (15.4% vs 40.2%, $p < 0.001$). However, these rates are obtained by dividing the number of patients who underwent shunt revision by the total in the entire cohort (14 revisions in 91 patients for fetal surgery versus 37 revisions in 92 patients in postnatal surgery). It could be questioned whether it is appropriate to include patients who do not have shunts in an analysis of shunt revision. If one considers only those patients who had shunts placed, the result is quite different (14 revisions in 40 patients for fetal surgery: 35% vs 37 revisions in 77 patients for postnatal surgery: 48%). Chi-square testing produces a p value of 0.177 for this comparison. So, should we really conclude that fetal surgery protects against shunt failure?

In the same issue of *Journal of Neurosurgery: Pediatrics* is a study by Mummareddy et al. regarding quality of life (QOL) in children with MMC.⁴ The authors administered QOL surveys to 74 children who had MMC repair between 1997 and 2003. They received 23 responses (31%): 11 with intrauterine closure and 12 with postnatal closure. Those with intrauterine closure scored significantly higher on overall and psychosocial QOL, but they observed no difference in physical QOL.

There are several limitations to that study, and the authors do an excellent job of detailing them. Possible participants were drawn from only one center, and the response rate was only 31%. The 2 groups are different in the proportion with hydrocephalus and the level of MMC, both of which have been shown to be associated with QOL in previous studies.^{3,5} The intrauterine closure group is highly selected by definition, since only families with financial resources, social support, and excellent healthcare access would have been eligible for this treatment in the first place. The study does not control for these factors, and, as the authors point out, the sample size is small, representing less than one-third of the eligible cohort. With this study design and these limitations, a measured conclusion with recognition that these are preliminary data would seem appropriate. Indeed, the text is measured and careful. However, the title of the paper says something quite different. "Intrauterine closure of myelomeningocele is associated with superior long-term quality of life than postnatal closure: a single-center study" strongly implies an answered question with limited room for doubt. Furthermore, the review article by Dewan and Wellons makes reference to this study without note of its limitations. The title would seem to infer a strength of evidence that the data cannot support. We anticipate that further such references to this paper from groups that lack the nuanced knowledge of the IUMC literature would similarly omit awareness or reference to the substantive limitations of the study design in the article by Mummareddy et al.

Another issue pertains to evolution of technique. As Dewan and Wellons point out, enthusiasm is high, and a large number of new centers for fetal surgery have opened

across the country in the last several years. Some centers prefer open while others prefer fetoscopic technique, and there is variation in technique between individual surgeons within these broader divisions. As such, progressively larger numbers of centers are treating a limited number of patients with widely varying experience levels. With limited exceptions, each center's cohort of patients will be underpowered to provide insight into the value or shortcomings of the techniques employed. Therefore, we would suggest that a registry of all patients undergoing IUMC might be useful in capturing and compiling this critical experience. The National Spina Bifida Patient Registry is a Centers for Disease Control and Prevention–sponsored, prospective registry that has enrolled more than 10,000 patients with spina bifida since its inception in 2004. Perhaps a parallel or extended component of the National Spina Bifida Patient Registry would provide fitting infrastructure to capture these important variables and cases. Centers such as Vanderbilt should be at the core of defining this extension of the registry and developing the critical variables.

Intrauterine closure of MMC is a great advance in neurosurgery. It has been shown with class I evidence to be beneficial for families who meet strict criteria.¹ The power and the promise of this work require that we ensure that studies of this treatment are rigorously conducted and reported.

Brandon G. Rocque, MD, MS

Betsy D. Hopson, MSHA

Jeffrey P. Blount, MD, MS

University of Alabama at Birmingham, Birmingham, AL

References

1. Adzick NS, Thom EA, Spong CY, Brock JW III, Burrows PK, Johnson MP, et al: A randomized trial of prenatal versus postnatal repair of myelomeningocele. *N Engl J Med* **364**:993–1004, 2011
2. Dewan MC, Wellons JC III: Fetal surgery for spina bifida. JNSPG 75th Anniversary invited review article. *J Neurosurg Pediatr* **24**:105–114, 2019
3. Flanagan A, Gorzkowski M, Altiok H, Hassani S, Ahn KW: Activity level, functional health, and quality of life of children with myelomeningocele as perceived by parents. *Clin Orthop Relat Res* **469**:1230–1235, 2011
4. Mummareddy N, Dewan MC, Huang A, Basem J, Bennett KA, Shannon CN, et al: Intrauterine closure of myelomeningocele is associated with superior long-term quality of life than postnatal closure: a single-center study. *J Neurosurg Pediatr* **24**:115–119, 2019
5. Rocque BG, Bishop ER, Scogin MA, Hopson BD, Arynychyna AA, Boddiford CJ, et al: Assessing health-related quality of life in children with spina bifida. *J Neurosurg Pediatr* **15**:144–149, 2015
6. Tulipan N, Wellons JC III, Thom EA, Gupta N, Sutton LN, Burrows PK, et al: Prenatal surgery for myelomeningocele and the need for cerebrospinal fluid shunt placement. *J Neurosurg Pediatr* **16**:613–620, 2015

Disclosures

The authors report no conflict of interest.

Correspondence

Brandon G. Rocque: brandon.rocque@childrensal.org.

INCLUDE WHEN CITING

Published online November 29, 2019; DOI: 10.3171/2019.8.PEDS19498.

Response

We read with interest the letter to the editor by Dr. Rocque, Ms. Hopson, and Dr. Blount regarding the invited article on fetal surgery for spina bifida that was written in celebration of the *Journal of Neurosurgery's* 75th anniversary. These concerns spill over to the original MOMS published in the *New England Journal of Medicine* in 2011, the follow-up study specific to hydrocephalus from that same cohort published in 2015, and the report on the results of a long-term study by our group in the same issue.^{1–3} We address these concerns below.

Issue 1 is that of the external validity of MOMS due to the study inclusion criteria. We are not well versed on a study without some sort of inclusion or exclusion criteria. A quick check of an early surgical trial, the Asymptomatic Carotid Atherosclerosis Study published in 1995, shows that even with its limited exclusion criteria, more than 42,000 patients were screened and only 1662 were enrolled. Unless we were to randomize all patients with said disease process, we have to bear the validity issues raised with inclusion criteria. One can state that a given study seems accurate for the studied population, but how valid, or generalizable, it is depends on those criteria. The MOMS inclusion criteria, as detailed in both the original paper and our review, were defined by the original investigators and based on expertise, data, and consensus in the time leading up to the trial. We, the readers and recipients of the results, are left to decide if we then keep the parameters steady in our implementation of the results or push the envelope closer to the edge of safety. Prior placental abruption, age younger than 18 years, and twin pregnancy all seem appropriate exclusions considering the risk entailed with the actual surgery itself.

The main issue that we should be discussing in regard to exclusion criteria is the BMI cutoff and the socioeconomic and racial exclusion that may or may not result from an interaction between those variables. The relationship of obesity to surgical morbidity is well known. At risk, that outside the safety envelope, is death of the fetus, or worse, that of the mother. What onboarding center is willing to risk that? What experienced center is? Yet, certainly there is indeed room to expand the BMI threshold with focused studies, as we do call for in the final paragraph of our review.

One additional point relevant to this issue of access to care is that of public versus private insurance and reimbursement for the procedure. How the reimbursement source lines up on socioeconomic status and racial lines can have a profound effect on what procedure is available to what population. TennCare, for example, has a mechanism in place to reimburse for fetal spina bifida repair. Alabama Medicaid does not. Reimbursement is a key facet of access to care; hence, advocacy at the level of state government by spina bifida experts, such as our colleagues at University of Alabama at Birmingham (UAB), would be key to improving access to the benefits noted in the initial study population for everyone.

So, on the issue of expanding the indications, we agree

with our colleagues. On devaluing a well-executed study because of the time-tested necessity of study criteria, we do not. We would suggest that all of us agree on the issue of reducing disparity of healthcare access.

Issue 2 is that of overstatement of the reduction in shunt burden to those in the intrauterine surgery group. This appears to be a simple issue of how one elects to interpret the concept of shunt revision. The authors of that article, which included the majority of the MOMS investigators, chose to conceptualize that a reduction in shunt revision in the intrauterine surgery cohort was due in part to a reduction in shunt burden.³ The letter authors are correct in their interpretation as well. Over the years, we have come to feel that the best way to reduce shunt infection, for example, is to reduce the need for shunt revision. The best way to reduce the need for shunt revision is to reduce the number of shunts placed, either by relaxing criteria or finding alternate operations, such as ETV or ETV/CPC. We would imagine that our colleagues at UAB find the reduction in the pool of patients in need of shunt revision to be a benefit, indeed a justification, of both.

Issue 3 is that the authors take issue with the title of the manuscript at hand.² “Association” refers to a relationship between variables. This has less strong implications than the term “causation,” which refers to direct cause and effect. These are well-known epidemiological terms, and one would hope the difference between the two is understood to the readers (and writers) of the medical literature. As written, the title plainly describes the association identified—no more, and no less. We reject the implication of a lack of rigor on our part. We have reported on those who answered the survey and the association noted and put stock in the integrity of the editorial process and scientific community in its evaluation of this body of work in the future.

Issue 4 is that of technique evolution. A degree of evolution is only expected as a procedure is adopted across a field, particularly as instruments and techniques advance. As a response to this occurring during the implementation phase since MOMS, we are on record several times calling for a “phase 4—type study” on both intrauterine and postnatal surgery, suggesting variables and outcomes of interest at several national talks and lectures on this topic over the past 3 years. We are encouraged to see that our colleagues at UAB agree and welcome their ideas of incorporating this effort into the National Spina Bifida Patient Registry.

Lastly, and with a final commentary: As we are joined in volume levels and expertise by other centers, we should indeed be sharing our learning, our outcomes, and our data. There may very well be further advances to be made, including BMI, reimbursement, and others, that result in no excluded patients in the future. This will depend on how we move forward as a field. Do we as the corpus of pediatric neurosurgery, with our deep engagement in the spina bifida community, lead this effort, by defining the next steps, designing the studies, and directly affecting the outcomes in the children that we treat? Or do we cede it to alternative subspecialists who operate and walk away, leaving others to care for the short- and long-term effects? We believe the answer is clear. Our colleagues at UAB

represent a strong voice of advocacy for this population. Their comments overall should be taken as intended, to further along the conversation and stoke action.

John C. Wellons III, MD, MSPH

Vanderbilt University Medical Center, Nashville, TN

Michael C. Dewan, MD, MSCI

The Hospital for Sick Children, University of Toronto, Toronto, ON, Canada

References

1. Adzick NS, Thom EA, Spong CY, Brock JW III, Burrows PK, Johnson MP, et al: A randomized trial of prenatal versus postnatal repair of myelomeningocele. *N Engl J Med* **364**:993–1004, 2011
2. Mummareddy N, Dewan MC, Huang A, Basem J, Bennett KA, Shannon CN, et al: Intrauterine closure of myelomeningocele is associated with superior long-term quality of life than postnatal closure: a single-center study. *J Neurosurg Pediatr* **24**:115–119, 2019
3. Tulipan N, Wellons JC III, Thom EA, Gupta N, Sutton LN, Burrows PK, et al: Prenatal surgery for myelomeningocele and the need for cerebrospinal fluid shunt placement. *J Neurosurg Pediatr* **16**:613–620, 2015

INCLUDE WHEN CITING

Published online November 29, 2019; DOI: 10.3171/2019.10.PEDS19505.

©AANS 2020, except where prohibited by US copyright law

Chiari formation

TO THE EDITOR: I read with interest the article on Chiari I “formation” by McCluggage and Oakes¹⁰ (McCluggage SG III, Oakes WJ: The Chiari I malformation. *JNSPG 75th Anniversary Invited Review Article. J Neurosurg Pediatr* **24**:217–226, September 2019). The title indicates that it is a review article and the authors claim that they discuss the evolution in the understanding of the subject. It is rather troublesome to notice that the authors have preferred to entirely omit our contributions to the subject in which it is proposed that atlantoaxial instability is the nodal point of pathogenesis of Chiari I formation and atlantoaxial stabilization is the treatment.

In 2015, our article in the *Journal of Neurosurgery: Spine* proposed that atlantoaxial instability is the primary point of pathogenesis and Chiari formation and a host of other musculoskeletal and neural alterations are secondary and nature’s protective responses.² Accordingly, we preferred the term Chiari “formation” to Chiari “malformation” and craniovertebral “alterations” to craniovertebral “anomalies.”³ It was proposed that tonsillar herniation forms a protective cushion, or is “nature’s airbag,” which protects neural structures from compression between the bones in the event of potential or manifest atlantoaxial instability.²

We proposed an alternative method of assessment of atlantoaxial instability based on the alignment of the facets on lateral profile imaging and evidence of instability during manual manipulation of bones. We introduced a novel concept of “central” or “axial” atlantoaxial instability,

wherein the atlantoaxial joint is unstable despite the traditionally accepted and validated parameter of alteration of the atlantodental interval on dynamic imaging being within normal range, and compression of the dura or neural structures opposite the odontoid process is not present.⁴ Chiari formation is more often associated with central or axial atlantoaxial instability. This form of instability is usually chronic or longstanding in nature. A host of musculoskeletal alterations that include basilar invagination, assimilation of the atlas, platybasia, Klippel-Feil abnormality, bifid arch of the atlas, and C2–3 fusion, and neural alterations that include Chiari formation, syringomyelia, syringobulbia, external syringomyelia, and external syringobulbia are secondary outcomes of chronic atlantoaxial instability. More importantly, all these secondary musculoskeletal and neural abnormalities are reversible following atlantoaxial fixation.^{8,9,11}

In response to our article in 2015,² a number of pediatric neurosurgeons based in North America argued that our concept is valid more for adult cases with Chiari I formation and in those patients in whom there are complex craniovertebral junction bone malformations.¹ We have subsequently published our results of surgical treatment of Chiari I formation by atlantoaxial fixation in pediatric age-group patients⁵ and in cases with Chiari I formation wherein there is no bone deformity.⁶

The authors have identified the more common symptoms of occipital headache, speech disturbance, and sleep apnea in cases with Chiari formation. We have recently shown that all these symptoms are entirely resolved in the immediate postoperative phase following atlantoaxial stabilization.⁷ We also demonstrated reversal of kyphoscoliosis following atlantoaxial fixation.⁹

I wish to take this opportunity to convey to the readers of this esteemed journal that the concept of atlantoaxial instability as a cause of Chiari I formation is here to stay. Any kind of foramen magnum decompression is not necessary for the treatment of Chiari formation.

Atul Goel, MCh

K.E.M. Hospital and Seth G.S. Medical College, Mumbai, India

References

1. Brockmeyer DL, Oakes WJ, Rozzelle C, Johnston J, Rocque BG, Anderson RC, et al: Chiari malformation Type 1 and atlantoaxial instability: a letter from the Pediatric Craniovertebral Society. **J Neurosurg Spine** 23:820–821, 2015 (Letter)
2. Goel A: Chiari malformation—is atlantoaxial instability the cause? Outcome analysis of 65 patients with Chiari malformation treated by atlantoaxial fixation. **J Neurosurg Spine** 22:116–127, 2015
3. Goel A: Is Chiari a “formation” or a “malformation.” **J Craniovertebr Junction Spine** 8:1–2, 2017
4. Goel A, Dhar A, Shah A, Jadhav D, Bakale N, Vaja T, et al: Central or axial atlantoaxial dislocation as a cause of cervical myelopathy: a report of outcome of 5 cases treated by atlantoaxial stabilization. **World Neurosurg** 121:e908–e916, 2019
5. Goel A, Gore S, Shah A, Dharurkar P, Vutha R, Patil A: Atlantoaxial fixation for Chiari I formation in pediatric age-group patients: report of treatment in 33 patients. **World Neurosurg** 111:e668–e677, 2018
6. Goel A, Kaswa A, Shah A: Atlantoaxial fixation for treat-

ment of Chiari formation and syringomyelia with no craniovertebral bone anomaly: report of an experience with 57 cases. **Acta Neurochir Suppl** 125:101–110, 2019

7. Goel A, Prasad A, Shah A, Gore S, Dharurkar P: Voice quality affection as a symptom of Chiari formation. **World Neurosurg** 121:e296–e301, 2019
8. Goel A, Sathe P, Shah A: Atlantoaxial fixation for basilar invagination without obvious atlantoaxial instability (Group B basilar invagination): outcome analysis of 63 surgically treated cases. **World Neurosurg** 99:164–170, 2017
9. Goel A, Vutha R, Shah A, Dharurkar P, Jadhav N, Jadhav D: Spinal kyphoscoliosis associated with Chiari formation and syringomyelia ‘recovery’ following atlantoaxial fixation: a preliminary report and early results based on experience with 11 surgically treated cases. **World Neurosurg** 125:e937–e946, 2019
10. McCluggage SG III, Oakes WJ: The Chiari I malformation. JNSPG 75th Anniversary Invited Review Article. **J Neurosurg Pediatr** 24:217–226, 2019
11. Shah A, Sathe P, Patil M, Goel A: Treatment of “idiopathic” syrinx by atlantoaxial fixation: report of an experience with nine cases. **J Craniovertebr Junction Spine** 8:15–21, 2017

Disclosures

The author reports no conflict of interest.

Correspondence

Atul Goel: atulgoel62@hotmail.com.

INCLUDE WHEN CITING

Published online November 29, 2019; DOI: 10.3171/2019.9.PEDS19511.

Response

We appreciate Dr. Goel’s letter and response to our invited review. Much of the discussion and reasoning concerning his approach was outlined in the letter to the editor¹ following Dr. Goel’s 2015 article and will not simply be repeated here. While a limited number of patients may benefit from fusion of the atlantoaxial joint, we have restricted this procedure to patients with demonstrable pathological movement of that area. To state that “all these [common] symptoms are entirely resolved in the immediate postoperative phase” with stabilization is a very fortunate outcome.

Time will tell if atlantoaxial fusion is the universal answer to this condition or if a more limited application of this technique to those with demonstrable instability is appropriate.

W. Jerry Oakes, MD

Samuel G. McCluggage III, MD

University of Alabama at Birmingham, AL

References

1. Brockmeyer DL, Oakes WJ, Rozzelle C, Johnston J, Rocque BG, Anderson RC, et al: Chiari malformation Type 1 and atlantoaxial instability: a letter from the Pediatric Craniovertebral Society. **J Neurosurg Spine** 23:820–821, 2015 (Letter)

INCLUDE WHEN CITING

Published online November 29, 2019; DOI: 10.3171/2019.10.PEDS19516.

©AANS 2020, except where prohibited by US copyright law