Neurosurgical Forum
LETTERS TO THE EDITOR**A closer look at synovial cysts**

TO THE EDITOR: We read with interest the article by Bruder et al.² (Bruder M, Cattani A, Gessler F, et al: Synovial cysts of the spine: long-term follow-up after surgical treatment of 141 cases in a single-center series and comprehensive literature review of 2900 degenerative spinal cysts. *J Neurosurg Spine* 27:256–267, September 2017). In this study, the authors presented the clinical data of 141 patients with synovial cysts who were treated surgically between 1997 and 2014. The majority of patients underwent hemilaminectomy, and a small subset of patients were treated with laminectomy or interlaminary fenestration. The facet joint was spared in most cases (63%), while the medial two-thirds and one-third of the facet joint were resected in 30% and 7% of patients, respectively. Based on the study results, 88% of patients with preoperative leg pain were “completely pain free” of leg pain after surgical treatment. Interestingly, the rate of low-back pain declined from 90% preoperatively to 5% postoperatively, and 86% of patients with preoperative low-back pain were back “pain free” after surgery. At 9 years of follow-up, 6% reported back pain.

A word of caution is in order here. This study reports that facet-sparing hemilaminectomy led to complete resolution of low-back pain in 86% of cases, and an absence of both leg and back pain on long-term follow-up. These patients were mostly in their 6th decade of life with facet disease, which led to the synovial cyst herniation. Low-back pain is a complex, multifactorial etiology that is unlikely to be resolved with hemilaminectomy or laminectomy. There is an emerging body of evidence that structural pathology, such as facet joint arthropathy and degenerative disc disease, is the main culprit of low-back pain.¹ Although one would argue that the synovial cyst itself is a manifestation of the facet joint degenerative process, surgically removing the cyst without addressing the diseased facet joint, which is the source of back pain, might decrease the severity of low-back pain, but not completely resolve it.^{4,5} For instance, a larger series by Xu et al. showed that patients treated with laminectomy or hemilaminectomy reported a higher incidence of back pain at 2-year follow-up than patients who underwent fusion ($p = 0.002$), although the same patient population showed improvement in back and leg pain immediately after surgery.⁵ Another series of 46 patients (by Weiner et al.) with a longer follow-up time than the current series showed that 28% of patients had low-back pain

after an average of 9.7 years postoperatively.⁴ Therefore, the current conclusion of persistence of back pain in 6% of patients 9 years after surgery is questionable.

Additional factors, including physical, social, psychological, and nonmodifiable genetic ones, could contribute interchangeably, along with structural pathologies, to low-back pain. The multitude of these factors poses a challenge for providers caring for patients with low-back pain. Typically, the presence of mechanical back pain and radiological signs of structural pathology (i.e., widening of the facet joints, hyperintense T2 signal on MRI, collapsed disc with Modic changes, spondylolisthesis) might favor fusion. With this in mind, it is important to recognize that there is no cure, as suggested in Bruden’s study, for low-back pain, even if we were to eliminate the pathological spinal unit with fusion.^{4,5}

Another marked limitation of this study that adds to its lack of statistical insignificance is the small number of follow-up patients (81 patients). Bruden et al. repeatedly stated that their series has the second longest follow-up time. However, their follow-up rate, 57%, was the lowest of all other studies. The authors tried to avoid confronting this limitation by comparing the total number of follow-up patients instead of the percentages, and by stating that they “achieve[d] a reasonable follow-up rate.” This major limitation should have alerted the reviewers of this article at your esteemed journal.

Moreover, the fact that the follow-up process was based on telephone interviews “with structured, questionnaire-based information” raises the following questions: how did the authors assess the patients’ motor function, sensation, and whether the leg pain was residual or new (different dermatome)? More importantly, how was the information about cyst recurrence, the level of recurrence, and rates of delayed fusion obtained?

In their conclusion, the authors stated, “With facet-sparing techniques, the stability of the segment can be preserved.” Although we agree with this statement, it does not hold true for the findings of this study. First, the facet-sparing technique was performed in 63% of the patients while 37% underwent partial facetectomy. Second, the authors did not state how they studied the “segmental instability” given that postoperative imaging was not performed in any of the patients. Finally, a clear distinction must be drawn between “segmental instability” (> 4 mm of motion of the vertebrae on flexion-extension radiographs) and “spondylolisthesis” (vertebral body slippage).³

The authors investigated the presence of postoperative “spondylolisthesis” in the follow-up period, but reported on “segmental instability” in the conclusions. All the previous methodological issues limit the conclusions of this study, and should call into question the relevance of its findings.

In summary, facet arthropathy is a cause of back pain. A hemilaminectomy may be able to remove the cyst and help the patient’s radiculopathy, but we doubt that most neurosurgeons would promise a 94% chance of “back pain free” outcome over the next decade, after a hemilaminectomy, as suggested by this article.

Seba Ramhmdani, MD

Ali Bydon, MD

Johns Hopkins University, Baltimore, MD

References

1. Adams MA: Biomechanics of back pain. *Acupunct Med* 22:178–188, 2004
2. Bruder M, Cattani A, Gessler F, Droste C, Setzer M, Seifert V, et al: Synovial cysts of the spine: long-term follow-up after surgical treatment of 141 cases in a single-center series and comprehensive literature review of 2900 degenerative spinal cysts. *J Neurosurg Spine* 27:256–267, 2017
3. Gopinath P: Lumbar segmental instability: points to ponder. *J Orthop* 12:165–167, 2015
4. Weiner BK, Torretti J, Stauff M: Microdecompression for lumbar synovial cysts: an independent assessment of long term outcomes. *J Orthop Surg Res* 2:5, 2007
5. Xu R, McGirt MJ, Parker SL, Bydon M, Olivi A, Wolinsky JP, et al: Factors associated with recurrent back pain and cyst recurrence after surgical resection of one hundred ninety-five spinal synovial cysts: analysis of one hundred sixty-seven consecutive cases. *Spine (Phila Pa 1976)* 35:1044–1053, 2010

Disclosures

The authors report no conflict of interest.

Response

We read with interest the Letter to the Editor by Ramhmdani and Bydon commenting on our recently published article.

To be clear, assessing the clinical situation via telephone interview is clearly a limitation of our study, as we mentioned in the *Limitations of the Study* section. We also mentioned the assessment of recurrent cysts and delayed fusion in the *Methods* section.

Without doubt, we were not able to contact 42% of patients for long-term follow-up and clearly stated the numbers and percentages, as well as the reasons for losses to follow-up in the text and tables. Nonetheless, we were able to obtain data in 81 patients with a mean follow-up period of more than 9 years. This remains the largest series with a follow-up period of more than 4 years (Supplemental Table 1), proving these data to be worthy of publication.

However, when questioning the results and conclusion of our report and focusing on the symptom “low-back pain,” Ramhmdani and Bydon were somehow misled. We never stated in our report that 86% of patients were free of leg or back pain at the time of long-term follow-up, and we never promised a 94% back-pain free rate or even “cure”

for low-back pain on long-term bases. In our study, additional low-back pain was reported in 127 patients before the operation, and 86% of those patients were pain free at short-term follow-up after surgery. At the time of long-term follow-up, however, 6% of 81 patients of the follow-up group reported low-back pain, and an additional 22% reported low-back pain under physical stress. As stated in Table 3, the overall rate of back pain in the follow-up group was 28% compared with the 22% back pain rate reported in the 594 cases of the literature in which “back pain” at the time of follow-up was mentioned. Therefore, we absolutely agree with Ramhmdani and Bydon that low-back pain has a “complex, multifactorial etiology,” and its treatment is and will be challenging.

In the end, we did not draw our conclusion from the presence or absence of back pain but from the overall long-term clinical situation, which should be the primary goal in degenerative spine surgery.

Markus Bruder, MD

Florian Gessler, MD

Volker Seifert, MD, PhD

Gerhard Marquardt, MD, PhD

Goethe University, Frankfurt, Germany

INCLUDE WHEN CITING

Published online April 20, 2018; DOI: 10.3171/2017.10.SPINE171093.

©AANS 2018, except where prohibited by US copyright law

Thinking laterally: beyond transoral decompression for irreducible ADD with os odontoidium

TO THE EDITOR: I read the recently published article by Wu et al.¹⁰ (Wu X, Wood KB, Gao Y, et al: Surgical strategies for the treatment of os odontoidium with atlantoaxial dislocation. *J Neurosurg Spine* 28:131–139, February 2018). The authors have described surgical strategies for treating patients with atlantoaxial dislocation (AAD) with os odontoidium. They classified patients based on reducibility of dislocation. Those with reducible AAD were treated with posterior stabilization alone, whereas those with irreducible AAD underwent transoral decompression followed by posterior stabilization. The posterior stabilization was short segment in most, with the exception of 4 patients in whom more than 2 segments were fused.

The authors have been practicing erstwhile surgical methods and have failed to take notice of contemporary techniques of direct posterior reduction. The surgical procedures for AAD have undergone significant change in the last 2 decades.^{1,4,6,7} The focus has shifted to the lateral C1–2 joints rather than decompression of the dens. Recent techniques involve manipulation of the C1–2 lateral joints to achieve reduction, followed by C1–2 fusion. In the current era of direct posterior reduction, to preoperatively differentiate reducible from irreducible AAD is to pre-empt the procedural difficulties.¹

FIG. 1. A: Preoperative midsagittal reconstructed CT image showing atlantoaxial dislocation (anteroposterior and vertical/basilar invagination) with os odontoideum. **B:** Parasagittal section through the lateral C1–2 joint showing severe obliquity as compared to the hard palate (*pink lines*). The C-1 lateral mass slips anterior and inferior to the C-2 lateral mass. **C:** Postoperative CT showing reduction of the dislocation. **D:** Parasagittal section passing through the C1–2 joint showing drilling of the joint to near normal. Stabilization has been achieved with C-1 lateral mass and C-2 pedicle screws. Figure is available in color online only.

The C1–2 joint provides maximum neck movements and is relatively less stable. The dens forms a pivot around which C-1 rotates. The atlantodental joint is stabilized by the transverse ligament, anterior arch of C-1, and integrity of the dens and C-2 body. With os odontoideum, the C-1 and dens move in unison. The dislocation is worsened by deformed lateral C1–2 joints or their lax capsules. The dislocation can be in the anteroposterior or lateral plane.^{5,7} The dislocation turns irreducible once the posterior edge of the C-1 lateral mass slips in front of the C-2 lateral mass. The lateral C1–2 joints in these cases can be oblique, giving rise to vertical as well as anteroposterior dislocation.⁹ The dens along with the C-1 anterior arch lies in front of the C-2 body (Fig. 1A). The dislocation in such cases fails to reduce despite traction.

Current surgical techniques involve opening and manipulating the lateral C1–2 joints. The facets can be drilled comprehensively if necessary, to achieve reduction. The presence of os odontoideum with irreducible AAD suggests complete spondyloptosis, and its reduction requires significant manipulation of the lateral masses.¹ Figure 1 shows radiological studies obtained in a patient with irreducible AAD with os odontoideum in whom the C1–2 joints were drilled and manipulated to achieve good reduction. With the current techniques, transoral decompression is circumvented. The procedure has its own complications and is best avoided.

Another concern is occipitocervical fusion. Delayed complications are seen after fusing the occipital squama with the cervical spine, unless multiple levels are fused.⁸ Such multilevel fusion compromises neck movements sig-

nificantly and adversely affects quality of life. Wu et al. have fused C1–3 for cases with anomalous vertebral artery (VA). Anomalous VA can be delineated and safeguarded without compromising the C1–2 fusion.^{2,3} It is rarely a deterrent to C1–2 short-segment fusion.³

A preoperative evaluation of the C1–2 lateral joints and the plane of dislocation helps in planning the surgery.^{7,9} The os odontoideum merely indicates congenital anomalies. Irreducible and reducible AAD can both be managed via a posterior approach alone. However, irreducible AAD is technically more demanding as it requires drilling of the facets and manipulation of the lateral masses. Regardless of the reducibility, the prime focus of the surgical strategy should be on the lateral C1–2 joints and not on the dens (os) and its removal.

Pravin Salunke, MCh
PGIMER, Chandigarh, India

References

- Deepak AN, Salunke P, Sahoo SK, Prasad PK, Khandelwal NK: Revisiting the differences between irreducible and reducible atlantoaxial dislocation in the era of direct posterior approach and C1–2 joint manipulation. *J Neurosurg Spine* **26**:331–340, 2017
- Salunke P, Futane S, Sahoo SK, Ghuman MS, Khandelwal N: Operative nuances to safeguard anomalous vertebral artery without compromising the surgery for congenital atlantoaxial dislocation: untying a tough knot between vessel and bone. *J Neurosurg Spine* **20**:5–10, 2014
- Salunke P, Sahoo S, Deepak AN: Anomalous vertebral artery is not a deterrent to C1–2 joint dissection and manipulation for congenital atlantoaxial dislocation. *Neurol India* **63**:1009–1012, 2015
- Salunke P, Sahoo S, Khandelwal NK, Ghuman MS: Technique for direct posterior reduction in irreducible atlantoaxial dislocation: multi-planar realignment of C1–2. *Clin Neurol Neurosurg* **131**:47–53, 2015
- Salunke P, Sahoo SK, Deepak AN, Garg R: Congenital C1–2 lateral translational dislocation: case illustration with pre- and postoperative imaging. *Pediatr Neurosurg* **51**:218–220, 2016
- Salunke P, Sahoo SK, Deepak AN, Ghuman MS, Khandelwal NK: Comprehensive drilling of the C1–2 facets to achieve direct posterior reduction in irreducible atlantoaxial dislocation. *J Neurosurg Spine* **23**:294–302, 2015
- Salunke P, Sahoo SK, Deepak AN, Khandelwal NK: Redefining congenital atlantoaxial dislocation: objective assessment in each plane before and after operation. *World Neurosurg* **95**:156–164, 2016
- Salunke P, Sahoo SK, Sood S, Mukherjee KK, Gupta SK: Focusing on the delayed complications of fusing occipital squama to cervical spine for stabilization of congenital atlantoaxial dislocation and basilar invagination. *Clin Neurol Neurosurg* **145**:19–27, 2016
- Salunke P, Sharma M, Sodhi HB, Mukherjee KK, Khandelwal NK: Congenital atlantoaxial dislocation: a dynamic process and role of facets in irreducibility. *J Neurosurg Spine* **15**:678–685, 2011
- Wu X, Wood KB, Gao Y, Li S, Wang J, Ge T, et al: Surgical strategies for the treatment of os odontoideum with atlantoaxial dislocation. *J Neurosurg Spine* **28**:131–139, 2018

Disclosures

The author reports no conflict of interest.

Response

We have read Dr. Salunke's comments concerning our paper on the treatment of os odontoideum with AAD. We appreciate his expert attention and advice.

What leads to irreducibility in os odontoideum is still unclear. As Dr. Salunke mentioned, "the dislocation [of the lateral C1–2 joints] turns irreducible once the posterior edge of the C-1 lateral mass slips in front of the C-2 lateral mass." Of course, the dislocation of the C1–2 joints in such cases fails to reduce despite traction. However, this is just one of the reasons for the irreducibility. A retrospective review was conducted by Dlouhy et al. to examine variable factors leading to os odontoideum with irreducible AAD, and the results indicated that abnormality of the transverse ligament and the presence of granulation tissue were the common factors contributing to irreducibility.²

The strategy for addressing os odontoideum with AAD remains controversial. Zhao et al. reported on a series of 279 cases of os odontoideum, whose treatment resulted in high satisfaction ratings and a low rate of complications.⁷ Transoral atlantoaxial joint release was recommended for cases with irreducible AAD before the posterior surgery. A retrospective study by Wang et al. included 904 patients with AAD, 268 of whom had irreducible AAD, which was successfully converted to reducible AAD via transoral atlantoaxial joint release.⁶ In a patient with irreducible AAD, paradental scar tissue and ossification prevented reduction of the atlantoaxial joint, which can be effectively and safely reduced by transoral anterior atlantoaxial release.⁵ According to evidence-based medicine, the transoral decompression procedure was recommended by the American Association of Neurological Surgeons (AANS) in 2001.

On the basis of our clinical work, direct posterior reduction and manipulation of the C1–2 lateral joints is effective for patients with reducible AAD and for many with irreducible AAD. Ni et al. reported on 24 patients with os odontoideum with irreducible AAD who were treated using a modified posterior screw-rod system, which was a safe and effective procedure for AAD secondary to os odontoideum.³ As the study by Deepak et al. showed, a direct posterior approach and C1–2 joint manipulation is feasible for intraoperative reduction.¹ However, for patients with irreducible AAD and anterior osseous obstruction and remaining spinal cord compression due to paradental scar tissue, combined transoral decompression and posterior fusion and stabilization is recommended. With resection of the remaining os odontoideum and soft tissue release around the odontoid and C-1 (residual hyperplasia of ligaments and scar tissue), the reduction can be further achieved by posterior internal fixation and fusion. Each operative method has its own advantages and indications.

Anomalous VAs are commonly observed, such as in high-riding VA (HRVA) and a narrow pedicle of the C-2 vertebra.⁴ Computed tomography scanning is a reliable method to evaluate anomalous VA. For a narrow pedicle of the C-2 vertebra, a C-2 pedicle screw can be successfully placed by an experienced surgeon. For an HRVA, however, a shorter C-2 pedicle screw could be inserted but with the risk of VA injury. Posterior C-2 pedicle screw fixation has

been widely used as a clinical fusion surgery. In cases with an HRVA, extension of the fixation and fusion segment to C-3 has been suggested because one more spinal segment fusion to C-3 can achieve stronger internal fixation with relatively little influence on the activity of the cervical spine.

Once again, we thank Dr. Salunke for his comments and advice and the further discussion of this topic.

Xinghuo Wu, MD

Cao Yang, MD

Union Hospital, Tongji Medical College,
Huazhong University of Science and Technology,
Wuhan, People's Republic of China

References

1. Deepak AN, Salunke P, Sahoo SK, Prasad PK, Khandelwal NK: Revisiting the differences between irreducible and reducible atlantoaxial dislocation in the era of direct posterior approach and C1–2 joint manipulation. *J Neurosurg Spine* 26:331–340, 2017
2. Dlouhy BJ, Policeni BA, Menezes AH: Reduction of atlantoaxial dislocation prevented by pathological position of the transverse ligament in fixed, irreducible os odontoideum: operative illustrations and radiographic correlates in 41 patients. *J Neurosurg Spine* 27:20–28, 2017
3. Ni B, Wu L, Guo Q, Lu X, Chen F, Guo X: Posterior open reduction and interlaminae compression fusion for os odontoideum with atlantoaxial dislocation. *World Neurosurg* 91:326–331, 2016
4. Wajanavisit W, Lertudomphonwanit T, Fuangfa P, Chanplakorn P, Kraiwattanapong C, Jaovisidha S: Prevalence of high-riding vertebral artery and morphometry of C2 pedicles using a novel computed tomography reconstruction technique. *Asian Spine J* 10:1141–1148, 2016
5. Wang C, Yan M, Zhou HT, Wang SL, Dang GT: Open reduction of irreducible atlantoaxial dislocation by transoral anterior atlantoaxial release and posterior internal fixation. *Spine (Phila Pa 1976)* 31:E306–E313, 2006
6. Wang S, Wang C, Yan M, Zhou H, Dang G: Novel surgical classification and treatment strategy for atlantoaxial dislocations. *Spine (Phila Pa 1976)* 38:E1348–E1356, 2013
7. Zhao D, Wang S, Passias PG, Wang C: Craniocervical instability in the setting of os odontoideum: assessment of cause, presentation, and surgical outcomes in a series of 279 cases. *Neurosurgery* 76:514–521, 2015

INCLUDE WHEN CITING

Published online April 27, 2018; DOI: 10.3171/2017.11.SPINE171268.

©AANS 2018, except where prohibited by US copyright law

Rotational thromboelastometry-guided transfusion protocol

TO THE EDITOR: We read with great interest the outstanding article by Guan and colleagues² (Guan J, Cole CD, Schmidt MH, et al: Utility of intraoperative rotational thromboelastometry in thoracolumbar deformity surgery. *J Neurosurg Spine* 27:528–533, November 2017). This timely study showed intraoperative rotational thromboelastometry (ROTEM)-guided transfusion for posterior thoracolumbar

deformity surgery (≥ 7 levels) significantly reduced total blood product transfusion requirements. However, we were left with unresolved questions.

We request clarification of the platelet transfusion pathway in the ROTEM-guided transfusion protocol.² Low EXTEM (external temogram) amplitude at 10 minutes (A10), amplitude at 20 minutes (A20), and maximum clot firmness (MCF) indicate poor clot strength, which is related to either a primary platelet abnormality or functionally low fibrinogen levels.^{5,8} Simultaneous assessment of FIBTEM (fibrinogen temogram) variables—in which platelet activity has been inhibited by the actin polymerization inhibitor cytochalasin D^{4,8}—can further delineate whether a platelet or fibrinogen abnormality is primarily responsible for low EXTEM results.^{5,6,9} Low FIBTEM clot strength may be related to hypofibrinogenemia, whereas adequate FIBTEM clot strength (in the presence of decreased overall non-inhibited platelet EXTEM clot strength) usually indicates platelet deficiency.³

In the Guan et al. study, the ROTEM group with EXTEM A20/MCF < 50 mm or FIBTEM A20/MCF < 7 mm received platelet transfusion.² However, this platelet-inhibited FIBTEM result suggests reduced clot strength from possible hypofibrinogenemia, in which case cryoprecipitate rather than platelet transfusion is generally recommended. The ROTEM-based blood loss treatment algorithms typically recommend administering platelets to patients with low EXTEM clot strength after FIBTEM clot strength is corrected.^{1,3,7,8}

In our experience with major adult spine surgery, intraoperative ROTEM-guided transfusion allowed early identification of hypofibrinogenemia, and significantly more cryoprecipitate was administered to ROTEM patients.⁵ Also, in patients who did not receive tranexamic acid (the only coagulation-modifying medication in our study), ROTEM patients had significantly less blood loss than their non-ROTEM counterparts, with a trend toward less transfusion of packed red blood cells.⁵

Again, we commend the authors on this excellent study.

Thomas J. Buell, MD
Davis G. Taylor, MD
Ching-Jen Chen, MD
Bhiken I. Naik, MBBCh

University of Virginia, Charlottesville, VA

References

1. Gorlinger K, Dirkmann D, Hanke AA, Kamler M, Kottenberg E, Thielmann M, et al: First-line therapy with coagulation factor concentrates combined with point-of-care coagulation testing is associated with decreased allogeneic blood transfusion in cardiovascular surgery: a retrospective, single-center cohort study. *Anesthesiology* **115**:1179–1191, 2011
2. Guan J, Cole CD, Schmidt MH, Dailey AT: Utility of intraoperative rotational thromboelastometry in thoracolumbar deformity surgery. *J Neurosurg Spine* **27**:528–533, 2017
3. Kozek-Langenecker SA, Afshari A, Albaladejo P, Santulano CA, De Robertis E, Filipesco DC, et al: Management of severe perioperative bleeding: guidelines from the European Society of Anaesthesiology. *Eur J Anaesthesiol* **30**:270–382, 2013
4. Lang T, Bauters A, Braun SL, Potzsch B, von Pape KW, Kolde HJ, et al: Multi-centre investigation on reference ranges for ROTEM thromboelastometry. *Blood Coagul Fibrinolysis* **16**:301–310, 2005
5. Naik BI, Pajewski TN, Bogdonoff DI, Zuo Z, Clark P, Terkawi AS, et al: Rotational thromboelastometry-guided blood product management in major spine surgery. *J Neurosurg Spine* **23**:239–249, 2015
6. Schöchl H, Frietsch T, Pavelka M, Jámor C: Hyperfibrinolysis after major trauma: differential diagnosis of lysis patterns and prognostic value of thromboelastometry. *J Trauma* **67**:125–131, 2009
7. Schöchl H, Voelckel W, Grassetto A, Schlimp CJ: Practical application of point-of-care coagulation testing to guide treatment decisions in trauma. *J Trauma Acute Care Surg* **74**:1587–1598, 2013
8. Solomon C, Ranucci M, Hochleitner G, Schöchl H, Schlimp CJ: Assessing the methodology for calculating platelet contribution to clot strength (platelet component) in thromboelastometry and thromboelastography. *Anesth Analg* **121**:868–878, 2015
9. Weiss G, Lison S, Spannagl M, Heindl B: Expressiveness of global coagulation parameters in dilutional coagulopathy. *Br J Anaesth* **105**:429–436, 2010

Disclosures

The authors report no conflict of interest.

Response

We thank Dr. Buell and his team for their careful reading of our paper and their insightful comments. As is obvious from their earlier publication on ROTEM for major spine surgery,¹ they have an excellent understanding of the application of this technology in this challenging patient population.

We agree that for patients with low FIBTEM MCF, cryoprecipitate is likely beneficial, although some authors do suggest transfusing platelets concurrently unless the EXTEM MCF is also high.² The reported algorithm was based on our early experiences with ROTEM for our spinal deformity operations. As they suggest, cryoprecipitate is now utilized as part of our protocol for patients who have low FIBTEM clot strength. Of note, in our study cohort, only one patient had a FIBTEM MCF value low enough to meet our platelet transfusion criteria, and today that patient would have probably received cryoprecipitate.

As mentioned in their letter, ROTEM is an excellent tool for identifying coagulopathies early in patients undergoing major spine operations. We are hopeful that our work will spur further investigation into its use.

Jian Guan, MD
Andrew T. Dailey, MD
University of Utah, Salt Lake City, UT

References

1. Naik BI, Pajewski TN, Bogdonoff DI, Zuo Z, Clark P, Terkawi AS, et al: Rotational thromboelastometry-guided blood product management in major spine surgery. *J Neurosurg Spine* **23**:239–249, 2015
2. Tanaka KA, Bolliger D, Vadlamudi R, Nimmo A: Rotational thromboelastometry (ROTEM)-based coagulation manage-

ment in cardiac surgery and major trauma. *J Cardiothorac Vasc Anesth* 26:1083–1093, 2012

INCLUDE WHEN CITING

Published online April 27, 2018; DOI: 10.3171/2018.1.SPINE1845.

©AANS 2018, except where prohibited by US copyright law

Radiographic differentiation of idiopathic spinal cord herniation versus dorsal arachnoid webs

TO THE EDITOR: We read with great interest the article published by Schultz et al.⁴ (Schultz R Jr, Steven A, Wessell A, et al: Differentiation of idiopathic spinal cord herniation from dorsal arachnoid webs on MRI and CT myelography. *J Neurosurg Spine* 26:754–759, June 2017) regarding the use of MRI and CT myelograms to reliably distinguish between spinal cord herniation (SCH) and dorsal arachnoid webs (DAWs). In their retrospective series, the authors demonstrate that indirect radiographic findings such as the “scalpel sign” described by Reardon et al.³ or a C-shaped dorsal cord indentation can differentiate between these two spinal pathologies.

We agree with the authors’ findings and would like to share a case of a DAW that was diagnosed based on the recognition of the scalpel sign on MRI. A 62-year-old woman was referred to us for evaluation of back pain, numbness, and tingling involving bilateral lower extremities and subtle gait imbalance. Upon recognition of the enlarged CSF space dorsal to the spinal cord on MRI of the cervical spine, a dedicated thoracic spine MRI sequence was performed that clearly demonstrated ventral displacement of the thoracic cord at the level of T-3 and the characteristic scalpel sign, without evidence of syringomyelia (Fig. 1). A diagnosis of DAW was made and surgical treat-

FIG. 1. A: Sagittal T2-weighted MR image demonstrating dorsal indentation of the thoracic spinal cord with characteristic scalpel sign. **B:** Corresponding axial image above the level of the dorsal indentation (at the level indicated by the arrow in panel A). **C:** Corresponding axial image at the level of the dorsal indentation (at the level indicated by the arrowhead in panel A).

ment was offered. Thoracic laminectomy and resection of the arachnoid web was performed. Thick arachnoid bands were clearly visualized at the T-3 and T-4 levels upon opening the dura, with compression of the spinal cord that improved following resection of the web (Fig. 2). The patient had an uneventful course and was neurologically intact after surgery.

The first reference to scalpel sign in relation to a DAW was made by Reardon et al. in 2013.³ While arachnoid webs and arachnoid cysts share a common pathogenesis, the challenge remains in differentiating SCH from an arachnoid web, which can often appear radiologically similar and could be mistaken for each other.² Idiopathic SCH and DAWs are relatively rare abnormalities that have a different pathogenesis and need to be treated differently during surgery.^{1,3} Treatment of SCH involves visualization of the ventral cord and dural defect, while DAW management involves lysis and removal of the intradural extramedullary arachnoid bands over the surface of the cord. While an arachnoid web is often not directly visualized on MRI, it appears that being aware of this entity and recognition of indirect radiographic findings—such as a scalpel-shaped dorsal indentation and preservation of the ventral CSF plane as described by the authors—can allow one to make the diagnosis of DAW sometimes even before

FIG. 2. A: Intraoperative image demonstrating the DAW. **B:** Lysis and resection of the DAW. **C:** Thoracic cord after DAW lysis and resection. Figure is available in color online only.

the development of severe myelopathy or syrinx, and can help plan surgery without the need for CT myelography in the majority of instances.³ The case we encountered lends further support to the diagnostic utility of the scalpel sign as a diagnostic radiological feature of DAW, which was confirmed during surgery.

Osmond C. Wu, MD

Manish K. Kasliwal, MD, MCh

University Hospitals Cleveland Medical Center, Cleveland, OH

References

1. Brus-Ramer M, Dillon WP: Idiopathic thoracic spinal cord herniation: retrospective analysis supporting a mechanism of diskogenic dural injury and subsequent tamponade. *AJNR Am J Neuroradiol* 33:52–56, 2012
2. Haber MD, Nguyen DD, Li S: Differentiation of idiopathic spinal cord herniation from CSF-isointense intraspinal extramedullary lesions displacing the cord. *Radiographics* 34:313–329, 2014
3. Reardon MA, Raghavan P, Carpenter-Bailey K, Mukherjee S, Smith JS, Matsumoto JA, et al: Dorsal thoracic arachnoid web and the “scalpel sign”: a distinct clinical-radiologic entity. *AJNR Am J Neuroradiol* 34:1104–1110, 2013
4. Schultz R Jr, Steven A, Wessell A, Fischbein N, Sansur CA, Gandhi D, et al: Differentiation of idiopathic spinal cord herniation from dorsal arachnoid webs on MRI and CT myelography. *J Neurosurg Spine* 26:754–759, 2017

Disclosures

The authors report no conflict of interest.

Response

We thank Drs. Wu and Kasliwal for their letter describing their experience in the imaging and management of a patient with a compressive dorsal thoracic arachnoid web. Their patient manifested typical imaging findings of this entity, such as a dorsal, upper thoracic location; a “scalpel”-shaped deformity of the cord; and importantly, preservation of the ventral CSF space at the level of the web. These findings are easily observed on the excellent conventional sagittal and axial T2-weighted images provided. However, we have encountered instances in which it has been necessary to perform additional imaging with a volumetric high-resolution T2-weighted sequence (T2 sampling perfection with application-optimized contrasts by using flip angle evolution) or CT myelography to confirm that the ventral subarachnoid space is indeed preserved, to definitively exclude SCH. These techniques may also aid in the differentiation of webs and cysts. Arachnoid webs are gracile structures and are not often directly seen, even using high-resolution imaging. The indirect signs we have described (pattern of deformity, preservation/interruption of the ventral CSF space) can reliably aid in their diagnosis and help differentiate them from idiopathic SCH. We are gladdened that Drs. Wu and Kasliwal were able to use these to successfully diagnose and manage their patient.

Randall Schultz Jr., MD

Prashant Raghavan, MBBS

University of Maryland School of Medicine, Baltimore, MD

INCLUDE WHEN CITING

Published online April 27, 2018; DOI: 10.3171/2017.11.SPINE171194.

©AANS 2018, except where prohibited by US copyright law

Cortical bone trajectory screw technique

TO THE EDITOR: I read with interest the article by Sakaura et al.⁷ (Sakaura H, Miwa T, Yamashita T, et al: Cortical bone trajectory screw fixation versus traditional pedicle screw fixation for 2-level posterior lumbar interbody fusion: comparison of surgical outcomes for 2-level degenerative lumbar spondylolisthesis. *J Neurosurg Spine* 28:57–62, January 2018). The authors describe a “cortical bone trajectory screw technique.” In the opening sentence of the article they state that such a technique is new and is nontraditional. I wish to indicate that the technique Sakaura et al. describe for the lumbar spine is the one we described earlier for use in the cervical spine.^{1,2,4–7} Considering the similarities with our technique, I feel that the authors could have referenced our work.

I entirely agree that the screw that traverses toward the disc space and uses the strong purchase afforded by juxtaposed cortex provides significantly added biomechanical strength to the screw. In 1997, we first discussed bicortical and tricortical techniques for cervical interbody screw fixation.³ The cortex adjoining the disc space is largely cortical in nature in contrast to the cortex in the middle of the vertebral body, which is relatively soft in consistency. In the cervical spine, the course of the screw away from the spinal cord makes screw insertion significantly safer. Larger screws can be used. We labeled this a bicortical method of screw insertion because the strength of cortices of the anterior vertebral body and that adjoining the disc space is exploited for screw insertion. We found that, in such a course, the entire length of the screw travels through the firmer parts of the vertebral body. As an extension of the technique, we indicated that the screw can traverse the disc space and pass through the cortex of adjoining vertebral bodies, and this we labeled the tricortical method of screw insertion. We have indicated that tricortical screw fixation can be used as a standalone procedure for interbody fixation. We have used 2 and even 4 tricortical screws for interbody fixation. Subsequently, in 2000, we described a number of permutations and combinations for bicortical and tricortical screws, using them in a standalone fixation technique or in combination with interbody plates.¹ Wide discectomy, denuding of the articular cartilage, and introduction of bone graft assist in arthrodesis of the vertebral bodies.

Atul Goel, MCh

King Edward VII Memorial Hospital and Seth G.S. Medical College, Mumbai, India

References

1. Goel A: Alternative tricortical methods of screw implantation for anterior cervical plate fixation: a preliminary report. *J Clin Neurosciences* 7:134–136, 2000

2. Goel A: Anterior transdiskal screws. **World Neurosurg** **112**:301, 2018
3. Goel A: Tricortical cervical inter-body screw fixation. **J Postgrad Med** **43**:4–7, 1997
4. Goel A: Tricortical method. **J Neurosurg** **99** (2 Suppl):245–246, 2003 (Letter)
5. Goel A, Cacciola F: Anterior approaches for multilevel cervical spondylosis, in Quinones-Hinjos A (ed): **Schmidek and Sweet's Operative Neurosurgical Techniques**, ed 6. Philadelphia: Elsevier Saunders, 2012, pp 1789–1800
6. Goel A, Cacciola F (eds): **The Craniovertebral Junction: Diagnosis, Pathology, Surgical Techniques**. Stuttgart: Thieme, 2011
7. Sakaura H, Miwa T, Yamashita T, Kuroda Y, Ohwada T: Cortical bone trajectory screw fixation versus traditional pedicle screw fixation for 2-level posterior lumbar interbody fusion: comparison of surgical outcomes for 2-level degenerative lumbar spondylolisthesis. **J Neurosurg Spine** **28**:57–62, 2018

Disclosures

The author reports no conflict of interest.

Response

No response was received from the authors of the original article.

INCLUDE WHEN CITING

Published online April 27, 2018; DOI: 10.3171/2017.12.SPINE171338.

©AANS 2018, except where prohibited by US copyright law