ERRATUM

“Awake” clipping of cerebral aneurysms: report of initial series

TO THE READERSHIP: An error appeared in the article by Abdulrauf et al. (Abdulrauf SI, Vuong P, Patel R, et al: “Awake” clipping of cerebral aneurysms: report of initial series. *J Neurosurg* [epub ahead of print October 21, 2016; DOI: 10.3171/2015.12.JNS152140]).

In Table 1, the number of deaths for the current study was incorrectly reported as 1.5–2.6; the actual mortality was 0. The corrected table is shown below.

The article has been corrected online as of February 3, 2017.

Saleem I. Abdulrauf, MD
St. Louis University, St. Louis, MO

CORRESPONDING ARTICLE See pp 311–318.

INCLUDE WHEN CITING

Published online February 3, 2017; DOI: 10.3171/2017.1.JNS152140a.

©AANS, 2017

TABLE 1. Contemporary (neurophysiological monitoring era) outcomes from clipping of UIAs

Authors & Year	Data Source/Study Design	No. of Patients	LOS (days)*	Discharged to a Facility (not home)†	No. of CVAs (%)	mRS Score (median)	Deaths†
Barker et al., 2003	NIS database/retrospective	3498	NA	565 (16.1)	NA	NA	73 (2.1)
Barker et al., 2004	NIS database/retrospective	3498	5	565 (16.1)	274 (7.8)	NA	73 (2.1)
Higashida et al., 2007	Health Economics and Outcomes Research Group of Boston Scientific Database/retrospective	1881	7.4	NA	249 (13.2)	NA	2.50
Alshekhlee et al., 2010	NIS database/retrospective	3738	4	NA	340 (9)	NA	60 (1.6)
Hoh et al., 2010	NIS/retrospective	4700	8.9 ± 10.1	NA	NA	NA	
Kotowski et al., 2013	60 published studies, meta-analysis/retrospective	9845	NA	NA	692 (7.0)	NA	157 (1.6)
McDonald et al., 2013	Premier Inc. Perspective database/retrospective	1380	NA	232 (17)	194 (14)	NA	10 (0.7)
Jalbert et al., 2015	Medicare provider analysis & review research identifiable files/retrospective	4357	7.1–9.2	(41.9–45.2)	NA	NA	NA
Present study	Single center/prospective	30	3	1 (3)	1 (3)	1	0

NA = not applicable.

* Values are median, mean ± SD, or range.

† Values are number (%) or percent range.