Neurosurgical Forum

LETTERS TO THE EDITOR

Preoperative DVT screening in neurooncology: is it cost-effective to perform duplex sonography in all patients?

TO THE EDITOR: We studied with keen interest the article by Pandey et al.⁵ regarding the role of preoperative deep vein thrombosis (DVT) screening in neurooncology (Pandey A, Thakur B, Hogg F, et al: The role of preoperative deep vein thrombosis screening in neurooncology. *J Neurosurg* 130:38–43, January 2019). We commend the authors for analyzing this pertinent clinical problem. They concluded that preoperative screening for DVT is a worthwhile endeavor in patients undergoing neurosurgical intervention. However, there are a few important points that we would like to bring to the attention of the readers of the *Journal of Neurosurgery*.

The authors have stated that the Wells score or serum D-dimer levels have a high negative predictive value in oncology patients. However, they cited the study by Carrier et al.² and conveyed that a low Wells score and a low D-dimer level are not a common combination in this patient group. They included all cases of cranial tumors in their study including meningiomas, gangliogliomas, and other low-grade gliomas. Since their study does not include only cancer patients, it would be incorrect to assume that a low Wells score and a low D-dimer level would be uncommon in the patients included in their study. We believe that many preoperative cases of brain tumor have a low Wells score and can rule out DVT in a significant portion of the cases. Additional investigation using Doppler ultrasound would not be cost-effective in all the cases. A young, neurologically intact patient without any known medical comorbidities diagnosed with low-grade glioma would have a low Wells score and does not, in our opinion, warrant additional Doppler ultrasonography. The authors found 7 patients to have DVT on screening via bilateral lower-limb venous Doppler ultrasonography. It would be relevant to know the Wells score level in these patients. A high Wells score would point towards the need for Doppler ultrasonography in such patients.

Various studies^{1,3,4,6} have found a prolonged duration of surgery to be a significant factor in the development of DVT. In their study, Pandey and colleagues started

prophylactic-dose enoxaparin in all patients, both those with and those without preoperative DVT findings, usually within 24 hours of surgery. This included patients who underwent biopsy. A patient with a low preoperative Wells score undergoing biopsy (short duration of surgery) for brain tumor does not mandate a DVT chemoprophylaxis in our opinion.

We believe that Doppler ultrasound screening should be reserved only for patients with a high Wells score or high D-dimer levels, and routine screening in all cases of brain tumors may not be cost-effective.

Hardik Sardana, MCh

Madhur Kaura, MBBS

Shweta Kedia, MCh

All India Institute of Medical Sciences, New Delhi, Delhi, India

References

1. Borde TD, Prasad C, Arimappamagan A, Srinivas D, Soman S: Incidence of deep venous thrombosis in patients undergoing elective neurosurgery – a prospective cohort based study. *Neurol India* 65:787–793, 2017
2. Carrier M, Lee AY, Bates SM, Anderson DR, Wells PS: Accuracy and usefulness of a clinical prediction rule and D-dimer testing in excluding deep vein thrombosis in cancer patients. *Thromb Res* 123:177–183, 2008
3. Ganau M, Prisco L, Cebula H, Todeschi J, Abid H, Ligarotti G, et al: Risk of deep vein thrombosis in neurosurgery: state of the art on prophylaxis protocols and best clinical practices. *J Clin Neurosci* 45:60–66, 2017
4. Li Q, Yu Z, Chen X, Wang J, Jiang G: Risk factors for deep venous thrombosis of lower limbs in postoperative neurosurgical patients. *Pak J Med Sci* 32:1107–1110, 2016
5. Pandey A, Thakur B, Hogg F, Brogna C, Logan J, Arya R, et al: The role of preoperative deep vein thrombosis screening in neurooncology. *J Neurosurg* 130:38–43, 2019
6. Smith SF, Biggs MT, Sekhon LH: Risk factors and prophylaxis for deep venous thrombosis in neurosurgery. *Surg Technol Int* 14:69–76, 2005

Disclosures

The authors report no conflict of interest.

Correspondence

Hardik Sardana: hardiksardana@yahoo.co.in.

INCLUDE WHEN CITING

Published online July 12, 2019; DOI: 10.3171/2019.3.JNS19665.

Response

No response was received from the authors of the original article.

©AANS 2019, except where prohibited by US copyright law

Establishing the role of prophylactic antiepileptic drugs in glioma and meningioma surgery

TO THE EDITOR: Joiner et al.³ report the findings of their meta-analysis on the effectiveness of perioperative antiepileptic drug (AED) prophylaxis for early- and late-onset seizures following oncologic surgery (Joiner EF, Youngerman BE, Hudson TS, et al: Effectiveness of perioperative antiepileptic drug prophylaxis for early and late seizures following oncologic neurosurgery: a meta-analysis. *J Neurosurg* 130:1274–1282, April 2019). The authors conclude that AED prophylaxis with phenytoin reduces seizure risk within the 1st week of surgery based on an analysis of 352 patients with 131 gliomas, 108 metastases, 97 meningiomas, and 17 suprasellar tumors. While the analysis did not show an effect on short- and long-term seizures, it nevertheless provides a timely opportunity to revisit the perennial question of the role of prophylactic AEDs in brain tumor surgery and the need for well-designed randomized controlled trials (RCTs) that address both perioperative and late-onset seizure risk. We propose that patients with glioma and meningioma will derive the greatest benefit from preventing seizures. However, unless the issue of equipoise is addressed, these trials are difficult to deliver, especially given the marked variation in practice and the almost universal use of prophylactic AEDs in some geographic territories.¹

In the United Kingdom, routine AED prophylaxis is less prevalent. A survey of members of the Society of British Neurological Surgeons (76 of 349 neurosurgeons responded across 28 of 32 adult neurosurgery units) demonstrated that for meningioma surgery the majority of surgeons (n = 44; 59%) almost never prescribe prophylactic AEDs but that 80% (n = 61) would be willing to participate in an RCT. For gliomas, AED prophylaxis is never used by 71% of surgeons but 74% would participate in an RCT.

In randomized trials of AED prophylaxis, there are several considerations. First, the incidence of postoperative epilepsy in seizure-naïve patients varies between gliomas and meningiomas, so the trial should be tumor specific—for example, based on recent studies on meningiomas,² to detect a reduction in the 12-month seizure rate from 10% to 5%, a power to 80% would require 435 patients in each arm (870 in total). Second, the pathological mechanism underlying seizure generation may be different such that the incurable nature and continued infiltrative growth of gliomas would lead to seizures affecting up to 50% of patients during their illness. In contrast, the inflammatory changes following meningioma surgery carry a much lower risk and might be amenable to novel antiinflammatory agents,

such as interleukin-1 receptor antagonists.⁴ This may provide additional benefit in the longer term by interrupting the epileptogenic process even with short-lived treatment in the acute phase. Third, we should be agnostic about the direction of effect of prophylactic AEDs and accept that their use might equally cause harm or benefit. Fourth, any study should collect data on imaging characteristics, surgical procedure, and postoperative complications in order to determine whether prophylaxis is beneficial in all cases or whether only a subset of patients will benefit.

The United Kingdom provides a unique setting for RCTs of prophylaxis due to the low levels of use of routine AED prophylaxis for glioma and meningioma surgery and the willingness of neurosurgeons to consider wider equipoise and participate in trials. The SPRING trial (Seizure Prophylaxis In Gliomas; <https://www.journalslibrary.nihr.ac.uk/programmes/hta/1631136/#/>) will determine whether in seizure-naïve newly diagnosed glioma patients prophylactic levetiracetam administered preoperatively and for at least 1 year postoperatively will produce a > 50% reduction in the risk of developing seizures when compared to no AED. The study will randomize 806 patients, in 15 neurosurgery units, over 3 years and is due to open in 2019. A similar trial for meningiomas, the Surgical Trial Of Prophylaxis for Epilepsy in Meningioma (STOP'EM), is in development.

While current AEDs suppress seizures, they do not address the underlying pathophysiological mechanisms that lower the threshold for uncontrolled electrical activity at the neuronal level. In the context of preventing seizures, the ideal therapeutic window may require pretreatment before a neuronal insult, which is problematic in many pathologies. However, in the context of meningioma surgery, there is the tantalizing possibility of inhibiting the perioperative inflammatory response to circumvent the induction of epileptogenesis, leading to long-term postoperative seizure control.

While these trials are in progress, this meta-analysis provides the current best evidence for prophylaxis in the immediate perioperative period, but it is limited by the small sample size and mixed tumor types. No doubt neurosurgeons will continue to use prophylactic AEDs, even in the absence of better quality evidence.

Michael D. Jenkinson, FRCS, PhD

Arousa Ali, MBChB, MRes

Abdurrahman I. Islim, MPhil

The Walton Centre NHS Foundation Trust, Liverpool, United Kingdom
Institute of Translational Medicine, University of Liverpool, Liverpool,
United Kingdom

Adel Helmy, FRCS, PhD

University of Cambridge, Cambridge, United Kingdom

Robin Grant, FRCP, MD

Western General Hospital, Edinburgh, United Kingdom

Acknowledgments

M.D.J. and R.G. are recipients of funding from National Institute for Health Research Health Technology Assessment programme (16/31/136).

References

1. Dewan MC, Thompson RC, Kalkanis SN, Barker FG II, Hadjipanayis CG: Prophylactic antiepileptic drug administration following brain tumor resection: results of a recent AANS/CNS Section on Tumors survey. *J Neurosurg* **126**:1772–1778, 2017
2. Islim AI, Ali A, Bagchi A, Ahmad MU, Mills SJ, Chavredakis E, et al: Postoperative seizures in meningioma patients: improving patient selection for antiepileptic drug therapy. *J Neurooncol* **140**:123–134, 2018
3. Joiner EF, Youngerman BE, Hudson TS, Yang J, Welch MR, McKhann GM II, et al: Effectiveness of perioperative antiepileptic drug prophylaxis for early and late seizures following oncologic neurosurgery: a meta-analysis. *J Neurosurg* **130**:1274–1282, 2019
4. Jyonouchi HG, Geng L: Intractable epilepsy (IE) and responses to anakinra, a human recombinant IL-1 receptor antagonist (IL-1ra): case reports. *J Clin Cell Immunol* **7**:456, 2016

Disclosures

The authors report no conflict of interest.

Correspondence

Michael D. Jenkinson: michael.jenkinson@liv.ac.uk.

INCLUDE WHEN CITING

Published online June 28, 2019; DOI: 10.3171/2019.1.JNS19134.

Response

In their letter, Jenkinson et al. address our recent meta-analysis on the effectiveness of AED prophylaxis following brain tumor resection and discuss planned RCTs in this area. As Jenkinson et al. note, our meta-analysis demonstrated a benefit of AED prophylaxis in the reduction of seizures within the 1st week of surgery (RR = 0.352, 95% CI 0.130–0.949, $p = 0.039$) but showed no effect of AED prophylaxis on a combined outcome of short- and long-term seizure rates.

We noted at the time of publication that these results reinforced the need for RCTs to address the question of AED prophylaxis but suggested that equipoise might provide an ethical barrier given the prevalence of AED prophylaxis in the United States and the evidence, albeit imperfect, that AED prophylaxis provides a reduction in early postoperative seizures. Jenkinson et al. argue that the question of equipoise is sufficiently addressed by the low prevalence of AED prophylaxis in the United Kingdom. Indeed, preliminary data from our group based on the MarketScan database (Truven Health Analytics) shows that the US rate of AED prophylaxis is 62.5% following the resection of intraparenchymal tumors and 61.5% after the resection of meningiomas. Jenkinson et al., citing a survey of British neurosurgeons, note that 71% of British neurosurgeons never use AED prophylaxis following glioma resection, and 59% almost never use AED prophylaxis following meningioma resection. Though such survey data are inherently limited, the results do nonetheless suggest that British use of AED prophylaxis may represent the inverse of US practice. As such, Britain does provide a more favorable environment for RCTs of AED prophylaxis than does the US, and lingering questions of equipoise boil down to an assessment of the quality of existing evidence in favor of AED prophylaxis.

We applaud the SPRING and STOP'EM trials to study seizure prophylaxis for glioma and meningioma resection, respectively. We concur that investigations must be stratified by tumor type given that these tumors convey differential risks of postoperative seizure, at least partially due to different underlying mechanisms. Indeed, the inability—based on limited sample size—to stratify patients by tumor type represented one of the key limitations of our meta-analysis. We further agree that any well-designed clinical trial must also incorporate data on imaging characteristics, surgical procedure, and postoperative complications. Such data are historically limited in studies on the subject, making it difficult to assess confounders and determine which patients would best benefit from prophylaxis. We commend Jenkinson et al. for their commitment to evaluate such variables as part of their studies.

One aspect of the SPRING trial design warrants further discussion. Our meta-analysis demonstrated only short-term benefit of postoperative seizure prophylaxis. However, the primary endpoint in the SPRING trial is seizure-frequency reduction of at least 50% at 1 year compared to no AEDs. Though secondary outcomes include time to first seizure and time to first tonic-clonic seizure, the lack of a dedicated short-term (i.e., 1 week) prophylaxis arm will make it difficult to compare the efficacy and side effect profile (mood, personality, fatigue, memory, and quality of life) of what current evidence suggests may be the optimal strategy.³ In our meta-analysis, we hypothesized that because long-term postoperative seizures are more likely to be associated with radiological evidence of tumor progression^{1,4} and that seizures related to tumor progression are not effectively prevented by AEDs,² it is likely that tumor growth gradually dilutes any difference in seizure incidence between AED prophylaxis arms and control arms over time. Thus, a large difference in seizure frequency between the two arms of the SPRING trial at 1 year may be difficult to realize, even if such a difference were present closer to time of surgery.

We maintain that a reduction of seizure frequency in the short-term perioperative period represents a valuable finding in our meta-analysis of previous trials and one that should be further evaluated in trials. It is unclear if the SPRING trial and STOP'EM trial will be designed to capture the potential benefits of this approach. We nonetheless support the development of these trials as a key step toward an evidence-based approach to perioperative AED prophylaxis for patients with brain tumors.

Evan F. Joiner, MD
Brett E. Youngerman, MD
Jeffrey N. Bruce, MD
Columbia University, New York, NY

References

1. Franceschetti S, Binelli S, Casazza M, Lodrini S, Panzica F, Pluchino F, et al: Influence of surgery and antiepileptic drugs on seizures symptomatic of cerebral tumours. *Acta Neurochir (Wien)* **103**:47–51, 1990
2. Glantz MJ, Cole BF, Forsyth PA, Recht LD, Wen PY, Chamberlain MC, et al: Practice parameter: anticonvulsant prophylaxis in patients with newly diagnosed brain tumors. Report

- of the Quality Standards Subcommittee of the American Academy of Neurology. **Neurology** 54:1886–1893, 2000
3. Grant R WC, Jenkinson M, Vale L, Marson T, Hill R, McLenny T: Seizure Prophylaxis in Glioma (SPRING): A phase III randomised trial comparing prophylactic Levetiracetam versus no prophylactic antiepileptic drug in newly diagnosed presumed supratentorial glioma. **National Cancer Research Institute Cancer Conference** (Abstract) (<http://abstracts.ncrri.org.uk/abstract/seizure-prophylaxis-in-glioma-spring-a-phase-iii-randomised-trial-comparing-prophylactic-levetiracetam-versus-no-prophylactic-antiepileptic-drug-in-newly-diagnosed-presumed-supratentorial/>) [Accessed May 22, 2019]
 4. Wu AS, Trinh VT, Suki D, Graham S, Forman A, Weinberg JS, et al: A prospective randomized trial of perioperative seizure prophylaxis in patients with intraparenchymal brain tumors. **J Neurosurg** 118:873–883, 2013

INCLUDE WHEN CITING

Published online June 28, 2019; DOI: 10.3171/2019.2.JNS19156.

©AANS 2019, except where prohibited by US copyright law

Doing more with less and the barebones neurosurgical setup

TO THE EDITOR: We read with interest the article by Corley and Rosseau¹ on the unmet need of global neurosurgery (Corley JA, Rosseau G: Encore careers: a solution to the unmet need in global neurosurgical care. *J Neurosurg* 130:1399–1401, April 2019). Their article has a deep personal meaning for surgeons in countries like ours, and we are very thankful that you have chosen to throw light on this aspect of neurosurgical practice. We deeply appreciate the efforts of all the organizations and individuals who are making proactive decisions regarding our collective responsibility toward global neurosurgical care. As surgeons living and trained in a large middle-income country with several low-income regions, we would like to describe our experiences in neurosurgical care at the ground level. There are two obvious problems faced by low- and middle-income countries (LMICs). The first is the need for skilled manpower. The second is the need for equipment. Within these issues are several gray zones that need to be addressed for the effective application of programs.

Skilled manpower to handle the gigantic gap in services is mired in difficulties related to licensing and incentives. International discussions on this matter need to include the question, “Who should be declared qualified to handle neurosurgical problems and emergencies in particular?” While neurosurgery remains a superspecialty, problems like head trauma and intracranial hemorrhage are so common that we need to ensure a more uniform distribution of services. A skewed ratio of specialist neurosurgeons in the population and an uneven distribution in rural and urban populations adversely affect management. For example, tier-1 cities like New Delhi and Bengaluru have nearly 250 times more neurosurgeons than a rural area just 60 miles away.⁶ These referral centers have the responsibility to manage gaps in manpower and qualified ancillary staff. Once a drill or an x-ray machine malfunctions, it can take weeks if not months to repair the same, thus significantly

compromising patient care. While some issues like having neurosurgically trained nurses or operative theater technicians may not even be an afterthought at many centers, numerous centers in India struggle for even a single support staff to manage cases. The only thing justifying the poor outcomes at these centers is the argument that “some care is better than no care.” Such a school of thought remains questionable. It counters the principle of an “all or none” phenomenon for neurological diseases (which remains as true to the nature of the disease as to the treatment).^{3,4}

There are resource-stricken neurosurgical centers in India where 15 beds have never held less than 25–30 patients, with multiple patients occupying one bed. There is still a dependency on Dandy’s hemostats, steel burrs, and perforators that need to be refurbished. Neurosurgery is considered a high-investment department by many hospitals. While general surgical services are commonly available, neurosurgery has an aura of complication and cost around it. This is a myth that needs to be explored and debunked by the international bodies.^{9,10,12}

The first solution would be to find ways to lower the cost of equipment.^{13,14} This would involve both the scientific and commercial aspects of production. Even if equipment could be manufactured cheaply, regional rules and taxation might raise the cost despite all efforts. Large companies with significant market shares in equipment constantly attempt to buy out and suppress local manufacturers. Market forces, profitability, and even share prices are blind to social responsibility.^{7,8} We cannot safely rely on conscience.

The second solution is the development of cheaper alternatives and techniques.¹¹ We have even tested and reported on the use of stationery binder clips as a cost-effective replacement for Raney clips for scalp hemostasis.¹⁵ Another example of such an alternative is the promotion of radio-frequency lesioning surgeries for various movement disorders and psychosurgery when a large population cannot afford deep brain stimulation procedures. These custom-tailored options should be respected in resource-stricken settings in which skilled manpower is struggling against a dearth of infrastructure.¹⁰

The third aspect of equipment deficits is a very interesting problem. While it is true that a lot of neurosurgical equipment is prohibitively expensive, we looked into historical precedents in our field to assess how neurosurgery was performed and what results were obtained when equipment was simpler. We asked the fundamental question, “What equipment would a surgeon really need to treat the majority of the neurosurgical burden in a low-resource region while providing acceptable results?” We hope to answer this with a sample neurosurgical setup (Table 1). The majority of the neurosurgical workload is operable with relatively cheap equipment as long as it is well maintained. Through our research, we found that the real prohibitive cost lies in ventilator support and imaging machines like CT and MRI units. While low-cost ventilators have been developed and made available in India,⁵ there has been no replacement for imaging machines yet. It is this “rate-limiting factor” that troubles us and needs as much charity as empowerment.

Trained neurosurgeons in LMICs are at a crossroads now. They are trained at the best centers in the world, have

TABLE 1. The basic cost of a barebones neurosurgical setup

Equipment	Expenditure (INR)	Expenditure (USD)
Neurosurgery operating table (made in India)	8–10 lakh	11,200–14,000
Head frame (made in India)	3.5–5 lakh	5,000–7,000
Basic neurosurgical instruments (made in India)	6–8 lakh	8,400–11,200
Craniotomy set		
Laminectomy set		
Basic microsurgical equipment		
Electric drill	9 ± 2 lakh	9,800–15,000
C-arm (made in India)	10–15 lakh	14,000–21,000
Bipolar & monopolar cautery	3–4 lakh	4,000–5,600
Suction machine (made in India)	1–1.5 lakh	1,400–2,200
Microscope (made in India)	20 ± 5 lakh	21,000–35,000
CT scanning machine	1–2.5 crore	140,000–355,000

INR = Indian rupee; USD = United States dollar.

witnessed masters of neurosurgery, and frequently visit various centers of excellence and international conferences for up-to-date knowledge. They also strive to give the best results to their patients but feel constrained by limited resources. Behind every success story, there are numerous tales of failure, which never get published or highlighted. Only stories of success get a place in the press. There is no substitute for a quality-assured setting, yet we strive to give comparable results with limited resources.^{2,9}

The need of the hour is to think outside of the box as regards capacity building. One such example is the training of bus drivers in Ghana to provide basic trauma life support when there is a road traffic accident.⁶ It led to remarkable improvement in neurotrauma care and delivery of services. Leading organizations may help in the capacity building, but innovative ideas need to come from the persons in the resource-stricken settings as only they are sufficiently aware of the difficulties at hand, local customs and rules, public psyche, realistic and imaginary expectations of society, and state of political willingness.

Nishant S. Yagnick, MCh

Paras Group of Hospitals, Gurgaon, India

Harsh Deora, MCh, DNB

Sanjay Gandhi Post Graduate Institute of Medical Education and Research, Lucknow, India

Manjul Tripathi, MCh

Aman Batish, MCh

Sandeep Mohindra, MCh

Postgraduate Institute of Medical Education and Research, Chandigarh, India

References

1. Corley JA, Rosseau G: Encore careers: a solution to the unmet need in global neurosurgical care. **J Neurosurg** **130**:1399–1401, 2019
2. Deora H, Tripathi M, Yagnick NS, Deora S, Mohindra S,

Batish A: Changing hands: why being ambidextrous is a trait that needs to be acquired and nurtured in neurosurgery. **World Neurosurg** **122**:487–490, 2019

3. Deora H, Yagnick NS, Tripathi M, Singh R, Mohindra S, Suri A, et al: Thrift shop hemostasis: I'm going to pop some hemostats, only got 20 dollars in my pocket! **Ann Clin Lab Res** **7**:282, 2019
4. Devi BI, Shukla DP, Bhat DI, Tripathi M, Warren A, Shanbhag NC, et al: Neurotrauma care delivery in a limited resource setting—lessons learned from referral and patient flow in a tertiary care center. **World Neurosurg** **123**:e588–e596, 2019
5. FE Online: Now, a portable ventilator for just Rs 35,000! AIIMS team develops low cost life-saving device. **Financial Express**. December 10, 2018 (<https://www.financialexpress.com/lifestyle/health/now-a-portable-ventilator-for-just-rs-35000-aiims-team-develops-low-cost-life-saving-device/1409357/>) [Accessed May 29, 2019]
6. Mishra RC: “Against the tide, nothing to hide”—the saga of a community neurosurgeon. **Neurol India** **65**:1213–1218, 2017
7. Mukherjee KK, Kumar N, Tripathi M, Oinam AS, Ahuja CK, Dhandapani SS, et al: Dose fractionated gamma knife radiosurgery for large arteriovenous malformations on daily or alternate day schedule outside the linear quadratic model: proof of concept and early results. A substitute to volume fractionation. **Neurol India** **65**:826–835, 2017
8. Mukherjee KK, Tripathi M: “Nuances of Gamma Knife radiosurgery for upper cervical spine lesions.” **Acta Neurochir (Wien)** **158**:2237, 2016
9. Suri A, Roy TS, Lalwani S, Deo RC, Tripathi M, Dhingra R, et al: Practical guidelines for setting up neurosurgery skills training cadaver laboratory in India. **Neurol India** **62**:249–256, 2014
10. Suri A, Tripathi M: Neurosurgery skills training laboratories and curriculum: a supplement to Halstedian practice. **J Neurosurg** **125**:1612–1613, 2016 (Letter)
11. Tiska MA, Adu-Ampofo M, Boakye G, Tuuli L, Mock CN: A model of prehospital trauma training for lay persons devised in Africa. **Emerg Med J** **21**:237–239, 2004
12. Tripathi M: Dose-fractionated Gamma Knife radiosurgery for large-volume arteriovenous malformations. **J Neurosurg** **129**:1660–1662, 2018 (Letter)
13. Tripathi M, Aziz TZ: Expected fate of radiofrequency lesioning: a silent death or a cold-blooded murder. **Stereotact Funct Neurosurg** **96**:274–275, 2018
14. Tripathi M, Kumar N, Mukherjee KK: Pushing the limits of the Leksell stereotactic frame for spinal lesions up to C3: fixation at the maxilla. **Acta Neurochir (Wien)** **158**:1691–1695, 2016
15. Yagnick NS, Singh R, Tripathi M, Mohindra S, Deora H, Suri A, et al: The need for grass root innovation in developing countries: case for stationary binder clips in scalp hemostasis. **World Neurosurg** **121**:222–226, 2019

Disclosures

The authors report no conflict of interest.

Correspondence

Manjul Tripathi: drmanjultripathi@gmail.com.

INCLUDE WHEN CITING

Published online July 5, 2019; DOI: 10.3171/2019.3.JNS19576.

Response

No response was received from the authors of the original article.

©AANS 2019, except where prohibited by US copyright law

Temporal lobe epilepsy: open or stereotactic surgery?

TO THE EDITOR: We read with great interest the study reported by Lee et al.⁴ that concerns stereoelectroencephalography-guided radiofrequency thermocoagulation (SEEG-guided RFTC) in mesial temporal lobe epilepsy (MTLE) (Lee CY, Li HT, Wu T, et al: Efficacy of limited hippocampal radiofrequency thermocoagulation for mesial temporal lobe epilepsy. *J Neurosurg* [epub ahead of print October 26, 2018. DOI: 10.3171/2018.4.JNS184]). Between the date of acceptance (April) and the date of publication (October) of their article, our group published two studies on a very similar subject,^{1,5} and our conclusions are divergent. Whereas Lee et al. concluded that SEEG-guided RFTC is a possible alternative to surgery in MTLE, we concluded that anterior temporal lobectomy (ATL) in this disease has a better outcome.^{1,5} It would, therefore, be of interest to analyze the reason for this.

To summarize, in a meta-analysis that included 296 patients who had undergone SEEG-guided RFTC, we failed to demonstrate a better outcome in the 39 patients with MTLE than in those with other etiologies, and the outcomes in the overall population (seizure-free rate: 23%; responder rate: 58%)¹ were much worse than those usually obtained after ATL.⁷ Furthermore, in a controlled study conducted on 70 patients with MTLE, significantly better outcomes were found in the ATL group than in the SEEG-guided RFTC group ($p < 0.001$).⁵

The divergence in the conclusions of our studies^{1,5} and those of Lee et al.⁴ are not explained by differences in lesion volume, which are in the same range. Their small study size could contribute to it, but we believe the main explanation is the duration of follow-up. During the first year after the procedure, the efficacy of SEEG-guided RFTC has been shown to dramatically decrease.^{2,3} This is the reason why the meta-analysis and the controlled study only included patients having at least a year of follow-up.^{1,5} In the study by Lee et al., the minimum follow-up by design was 6 months, and only two of the seven patients had a follow-up that was greater than 1 year; therefore, it would be interesting to have data from a more long-term follow-up for these patients.⁴

More generally, it is notable that the current recommendations are to immediately perform ATL in MTLE without SEEG when phase I presurgical investigations are congruent.⁷ Performing SEEG-guided RFTC for MTLE in patients who will subsequently require an ATL due to poor outcomes will increase the risk of surgery (stereotactic procedure and surgery vs surgery alone) even if SEEG and SEEG-guided RFTC are safe procedures.^{1,6} We, therefore, believe that on the basis of the present literature, there is no reason to propose SEEG-guided RFTC as an alternative to ATL in routine clinical practice.

Pierre Bourdillon, MD, MSc¹⁻⁴
Sylvain Rheims, MD, PhD^{1,2,5}

Jean Isnard, MD, PhD¹

Marc Guénot, MD, PhD^{1,2,6}

¹Hospital Pierre Wertheimer, Hospices Civils de Lyon, Lyon, France

²University of Lyon, Faculty of Medicine Claude Bernard, Lyon, France

³Sorbonne University, Paris, France

⁴Brain and Spine Institute, INSERM U1127, CNRS, UMR7225, Paris, France

⁵TIGER, Neuroscience Research Center of Lyon, INSERM U1028, CNRS 5292 Lyon, France

⁶NEUROPAIN Team, Lyon Neuroscience Research Center, INSERM U1028, CNRS 5292, Lyon, France

Acknowledgments

We thank Dr. Philip Robinson (DRCI, Hospices Civils de Lyon, France) for help in manuscript preparation and English editing.

References

1. Bourdillon P, Cucherat M, Isnard J, Ostrowsky-Coste K, Catenioix H, Guénot M, et al: Stereo-electroencephalography-guided radiofrequency thermocoagulation in patients with focal epilepsy: a systematic review and meta-analysis. *Epilepsia* **59**:2296–2304, 2018
2. Bourdillon P, Isnard J, Catenioix H, Montavont A, Rheims S, Ryvlin P, et al: Stereo electroencephalography-guided radiofrequency thermocoagulation (SEEG-guided RF-TC) in drug-resistant focal epilepsy: results from a 10-year experience. *Epilepsia* **58**:85–93, 2017
3. Cossu M, Fuschillo D, Casaceli G, Pelliccia V, Castana L, Mai R, et al: Stereoelectroencephalography-guided radiofrequency thermocoagulation in the epileptogenic zone: a retrospective study on 89 cases. *J Neurosurg* **123**:1358–1367, 2015
4. Lee CY, Li HT, Wu T, Cheng MY, Lim SN, Lee ST: Efficacy of limited hippocampal radiofrequency thermocoagulation for mesial temporal lobe epilepsy. *J Neurosurg* [epub ahead of print October 26, 2018. DOI: 10.3171/2018.4.JNS184]
5. Moles A, Guénot M, Rheims S, Berthiller J, Catenioix H, Montavont A, et al: SEEG-guided radiofrequency coagulation (SEEG-guided RF-TC) versus anterior temporal lobectomy (ATL) in temporal lobe epilepsy. *J Neurol* **265**:1998–2004, 2018
6. Mullin JP, Shriver M, Alomar S, Najm I, Bulacio J, Chauvel P, et al: Is SEEG safe? A systematic review and meta-analysis of stereo-electroencephalography-related complications. *Epilepsia* **57**:386–401, 2016
7. Ryvlin P, Cross JH, Rheims S: Epilepsy surgery in children and adults. *Lancet Neurol* **13**:1114–1126, 2014

Disclosures

The authors report no conflict of interest.

Correspondence

Pierre Bourdillon: pierre.bourdillon@neurochirurgie.fr.

INCLUDE WHEN CITING

Published online February 22, 2019; DOI: 10.3171/2018.11.JNS183107.

Response

No response was received from the authors of the original article.

©AANS 2019, except where prohibited by US copyright law

How does size ratio affect the clinical result of open surgery for cerebral aneurysms?

TO THE EDITOR: I read with interest the article by Matsukawa et al.¹ (Matsukawa H, Kamiyama H, Kinoshita Y, et al: Morphological parameters as factors of 12-month neurological worsening in surgical treatment of patients with unruptured saccular intracranial aneurysms: importance of size ratio. *J Neurosurg* [epub ahead of print September 21, 2018. DOI: 10.3171/2018.4.JNS173221]). They reported on their large cohort of patients with surgically treated unruptured saccular intracranial aneurysms (USIAs). They found that the ratio between the aneurysm diameter and the parent artery diameter was a factor more related to neurological worsening at 12 months after surgery than the absolute size of the aneurysm. As the article mentioned, studies based on computed flow dynamics analysis showed that size ratio and aspect ratio are related to aneurysm bleeding or to the possibility of sufficient coil embolization. However, in most cases that involve surgical clipping, we can observe the structures around the aneurysm. The patency of the related arteries should have been confirmed after application of clips or after revascularization, using Doppler sonography and/or indocyanine green video-angiography, which they should also have used. The authors speculated that a larger size ratio resulted in a need for greater manipulation of structures around the aneurysms, which caused neurological worsening. However, as they described, the size ratio is highly correlated with the maximal size of the aneurysm, since the ratio is derived by simply dividing the diameter of the aneurysm by the diameter of the parent artery. Considering previous reports noting risk factors related to surgical clipping, Matsukawa and colleagues' results may also indicate that large aneurysms pose a higher risk of surgical morbidity. Or, owing to their good clinical outcome, the number of cases of neurological worsening was so small that the statistical power was not sufficient enough to extract the true primary risk factor after multiple regression analyses. In addition, according to their results, the specific anatomical location of the aneurysm and the presence of a postoperative ischemic lesion significantly contributed to the neurological worsening at 12 months. Generally speaking, distal basilar aneurysms are considered to be difficult-to-treat lesions due to the risk of infarction of the thalamoperforating artery. Also, large paraclinoid-segment aneurysms may pose the surgical risk of occluding the anterior choroidal artery. I suppose that there is multicollinearity between the aneurysm's location and postoperative ischemic lesion causing neurological worsening at 12 months.

Toshikazu Kimura, MD, PhD

Japanese Red Cross Medical Center, Tokyo, Japan

References

1. Matsukawa H, Kamiyama H, Kinoshita Y, Saito N, Hatano Y, Miyazaki T, et al: Morphological parameters as factors

of 12-month neurological worsening in surgical treatment of patients with unruptured saccular intracranial aneurysms: importance of size ratio. *J Neurosurg* [epub ahead of print September 21, 2018. DOI: 10.3171/2018.4.JNS173221]

Disclosures

The author reports no conflict of interest.

Correspondence

Toshikazu Kimura: tkim-ky@umin.ac.jp.

INCLUDE WHEN CITING

Published online November 23, 2018; DOI: 10.3171/2018.10.JNS182796.

Response

We appreciate the valuable comments from Dr. Kimura regarding our article.

First, as Dr. Kimura suggested, the patency of the related arteries should have been confirmed after application of clips or after revascularization by performing Doppler sonography and/or indocyanine green video-angiography; however, each of these monitoring tools has its limitations in detecting blood flow insufficiency in the perforating arteries. A perfect method for monitoring blood flow disturbance in the perforating arteries has not been established. In fact, even with confirmation of the patency of perforating arteries with either of the aforementioned intraoperative monitoring modalities and without causing direct injury to perforating arteries, an unexpected infarction due to blood flow reduction in the parent artery and/or perforating arteries branching close to the aneurysm can still occur.¹⁻³ We strongly agree with these facts.

Second, Dr. Kimura stated, "Considering previous reports noting risk factors related to surgical clipping, Matsukawa and colleagues' results may also indicate that large aneurysms pose a higher risk of surgical morbidity." In this regard, we have performed multivariate analysis for 12-month neurological worsening based on the size ratio, maximum size, dome-to-neck ratio, and aspect ratio, as well other factors as covariates. After adjustment of these aneurysm morphological characteristics, size ratio was still a 12-month negative prognostic factor (Table 3 in our article). Therefore, we consider size ratio to be a risk factor for 12-month neurological worsening in surgically treated patients with USIAs, although the size ratio should be further studied in a large, prospective, observational cohort.

Third, Dr. Kimura suggested that there is multicollinearity between the aneurysm location and the presence of a postoperative ischemic lesion causing neurological worsening at 12 months. Because this is true, as stated above, we have performed multivariate analysis using aneurysm location and postoperative ischemic lesion as covariates (Table 3 in our article).

Hidetoshi Matsukawa, MD, PhD

Rokuya Tanikawa, MD

Stroke Center, Teishinkai Hospital, Sapporo, Japan

References

1. Friedman JA, Pichelmann MA, Piepgras DG, Atkinson JL, Maher CO, Meyer FB, et al: Ischemic complications of sur-

- gery for anterior choroidal artery aneurysms. **J Neurosurg** **94**:565–572, 2001
2. Sakuma J, Suzuki K, Sasaki T, Matsumoto M, Oinuma M, Kawakami M, et al: Monitoring and preventing blood flow insufficiency due to clip rotation after the treatment of internal carotid artery aneurysms. **J Neurosurg** **100**:960–962, 2004
 3. Yaşargil MG, Yonas H, Gasser JC: Anterior choroidal artery aneurysms: their anatomy and surgical significance. **Surg Neurol** **9**:129–138, 1978

INCLUDE WHEN CITING

Published online November 23, 2018; DOI: 10.3171/2018.10.JNS182830.

©AANS 2019, except where prohibited by US copyright law