Neurosurgical Forum LETTERS TO THE EDITOR

Is 7-Tesla MRI necessary in the assessment of microstructural injury to visual pathways due to pituitary adenomas?

TO THE EDITOR: We read with great interest the article by Rutland et al.⁵ (Rutland JW, Padormo F, Yim CK, et al: Quantitative assessment of secondary white matter injury in the visual pathway by pituitary adenomas: a multimodal study at 7-Tesla MRI. *J Neurosurg* [epub ahead of print January 18, 2019. DOI: 10.3171/2018.9.JNS182022]). The authors for the first time investigated the microstructural damage to the optic tracts and radiations in patients with pituitary macroadenomas using 7-T MRI, adding significant insight to the transformative studies on the changes of visual tract structures and function. However, several issues should be addressed.

First, the necessity and feasibility of adopting 7-T MRI is debatable. The optic radiation is located deep in the brain surrounded by multiple other white matter tracts. It is therefore difficult to visualize the microstructure of the optic radiation and calculate its characteristics using conventional MRI.² With higher spatial resolution, ultra-high-field MRI was able to resolve between the visual pathway and neighboring white matter tracts.⁴ However, 7-T MRI is not unique in this capacity. Diffusion tensor imaging (DTI) using a 3-T MRI scanner was also reported to be able to quantitatively measure the optic pathways, including the optic chiasm, optic tract, and the optic radiation.² Until 2017, there were only about 60 7-T MRI systems throughout the world.⁴ It is doubtful that 7-T MRI will be widely used in the next decade because of challenges such as magnetic field inhomogeneity, patient safety, comfort issues, and cost.³ On the contrary, 3-T MRI, which has been applied for clinical use for decades, may be more economical and feasible for use in radiological studies of visual pathways.

Second, the algorithm for the measurement of visual pathways deserves discussion. Quantitative parameters of diffusion-weighted MRI (DWI) could reveal detailed white matter microstructural changes. However, it is important to note that there are prominent variations in scalar diffusion imaging measurements within the white matter tracts.¹ The tract-averaged approach employed in this article, which averages the scalar values in a tract dissection

into a single estimate, might lead to inaccurate results by ignoring potential variations. The along-tract analysis by Colby et al.,¹ which provided extensible end-to-end along-tract workflow, was confirmed to be more accurate than the tract-averaged approach. Therefore, the results might be improved with the along-tract approach, and further studies with this approach are warranted.

Third, the effects of anatomically correlated gyral bias need to be discussed. Gyral bias has been reported in many gyral blades across the brain, and it appears to be challenging for the current fiber-tracking algorithms, including both deterministic and probabilistic, to terminate on gyral crowns, even for the high-resolution diffusion data sets.⁶ Thus, this gyral bias will lead to an artificial increase in the mean values of DWI parameters within tracts and a significant deviation of group analysis results. Furthermore, in this study, the diffusion indices were calculated using only 20 points located along the optic tract to the optic radiation, so the results of this method might be largely deviated by the tracts' natural variations. Optimal methods for data acquisition and analysis should be employed to take full advantage of ultra-high-field MRI on measuring microstructures along the entire visual pathway.

Xiaopeng Guo, MD

Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China
China Pituitary Disease Registry Center, Chinese Pituitary Adenoma Cooperative Group, Beijing, China

Shun Yao, MD

The First Affiliated Hospital, Sun Yat-sen University, Guangzhou, Guangdong, China

Bing Xing, MD

Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China
China Pituitary Disease Registry Center, Chinese Pituitary Adenoma Cooperative Group, Beijing, China

References

- Colby JB, Soderberg L, Lebel C, Dinov ID, Thompson PM, Sowell ER: Along-tract statistics allow for enhanced tractography analysis. *Neuroimage* **59**:3227–3242, 2012
- Kamali A, Hasan KM, Adapa P, Razmandi A, Keser Z, Lincoln J, et al: Distinguishing and quantification of the human visual pathways using high-spatial-resolution diffusion tensor tractography. *Magn Reson Imaging* **32**:796–803, 2014
- Karamat MI, Darvish-Molla S, Santos-Diaz A: Opportuni-

ties and challenges of 7 Tesla magnetic resonance imaging: a review. **Crit Rev Biomed Eng** 44:73–89, 2016

4. Kraff O, Quick HH: 7T: Physics, safety, and potential clinical applications. **J Magn Reson Imaging** 46:1573–1589, 2017
5. Rutland JW, Padormo F, Yim CK, Yao A, Arrighi-Allisan A, Huang KH, et al: Quantitative assessment of secondary white matter injury in the visual pathway by pituitary adenomas: a multimodal study at 7-Tesla MRI. **J Neurosurg** [pub ahead of print January 18, 2019. DOI: 10.3171/2018.9.JNS182022]
6. Schilling K, Gao Y, Janve V, Stepniewska I, Landman BA, Anderson AW: Confirmation of a gyral bias in diffusion MRI fiber tractography. **Hum Brain Mapp** 39:1449–1466, 2018

Disclosures

The authors report no conflict of interest.

Correspondence

Bing Xing: xingbingemail@aliyun.com.

INCLUDE WHEN CITING

Published online May 17, 2019; DOI: 10.3171/2019.2.JNS19413.

Response

We thank Drs. Guo, Yao, and Xing for their interest in our recent publication.

First, we acknowledge that quantitative examination of the optic pathway in the context of pituitary adenoma has been performed by previous groups using 3-T MRI.^{3,4} These preliminary investigations were referenced in the article, and their findings were compared with the results of the present study. The primary contributions of the present study were to acquire DW images of the optic pathway in high spatial resolution using ultra-high-field 7-T MRI and to correlate microstructural findings to detailed neuro-ophthalmological evaluation in order to elucidate the structural substrates that contribute to functional loss in patients with optic chiasm compression by pituitary adenoma. We believe that ultra-high-field examination of the optic tracts is especially important, as the tracts are completely surrounded by cerebrospinal fluid before entering the brain,² and partial volume effects due to the larger voxels employed by previous investigations likely resulted in significant artifact. Therefore, while 1.5- and 3-T MRI scanners are indeed more numerous than 7-T scanners, we believe that the physical advantages conferred by ultra-high-field MRI make application of this technology to discern the subtle microstructural effects of chiasm compression in the human visual system clinically relevant. As of January 2018, there were approximately 65 7-T scanners installed worldwide and 79 7-T scanners ordered for installation. Siemens Healthcare has introduced the clinically targeted Terra system for which the company obtained 510K approval from the FDA in October 2017. By January 2018, 9 Terra systems had been installed worldwide and 22 were ordered for installation, indicating rapid adoption. We believe that ultra-high-field MRI and advanced imaging modalities will become increasingly pervasive in the coming years and greatly benefit in vivo characterization of skull base pathology.

Second, we agree that a spatially normalized evaluation of optic pathway microstructure is an important line

of inquiry and may capture regional deviation in white matter integrity that could be overlooked by examining the tract as a whole. For this reason, we chose to calculate white matter microstructure at 20 equivalent points along both the optic tracts and optic radiations. These results can be seen in Fig. 4 of our article. By plotting these data on a spatially normalized axis, we observed that patients and controls exhibited similar spatial patterns of diffusion characteristics within the tracts and radiations; however, patients were shifted inferiorly on the fractional anisotropy axis and inferiorly on the mean diffusivity axis, indicating overall reduction in white matter integrity. This analysis is homologous to the along-tract analysis employed by Colby et al.,¹ and we believe that within-tract properties of the optic tracts and radiations were appropriately described in the present study.

Finally, we agree that the effects of gyral bias are an important consideration and an inherent limitation of DWI-based fiber tracking. The following methodological factors were employed in the present study, which we believe helped to mitigate the effects of gyral bias. First, constrained spherical deconvolution (CSD),⁶ a higher-order fiber orientation algorithm, was used for fiber reconstruction. While not immune to the effects of gyral bias, the CSD method has been shown to provide a considerable reduction in gyral bias (1%–20%).⁵ Second, the 7-T DWI sequence in the present study permitted higher spatial resolution (1.05 mm isotropic) than traditional 1.5- or 3-T studies that typically have 2- to 3-mm DWI voxel sizes.⁵ The high spatial resolution in the present study allowed more accurate identification of fiber insertion points into the cortex, thereby minimizing the effects of gyral bias. Lastly, the DWI acquisition employed in the present study was a high-angular-resolved DWI (HARDI) sequence. HARDI techniques can resolve more fiber angles along the gray matter/white matter boundary, reducing the effects of the convoluted cerebral cortex.⁵ We believe that this advanced acquisition, combined with CSD and high-spatial resolution DWI, minimized the inherent effects of gyral bias and allowed appropriate fiber termination in the primary visual cortex. Despite these efforts to minimize gyral bias, we recognize this effect as a technical limitation of DWI and acknowledge the importance of developing next-generation fiber-tracking algorithms that can overcome the complex anatomical interface of the white and gray matter and further reduce the effects of gyral bias.

John W. Rutland, BA
Bradley N. Delman, MD, MS
Raj K. Shrivastava, MD
Priti Balchandani, PhD

Icahn School of Medicine at Mount Sinai, New York, NY

References

1. Colby JB, Soderberg L, Lebel C, Dinov ID, Thompson PM, Sowell ER: Along-tract statistics allow for enhanced tractography analysis. **Neuroimage** 59:3227–3242, 2012
2. Dasenbrock HH, Smith SA, Ozturk A, Farrell SK, Calabresi PA, Reich DS: Diffusion tensor imaging of the optic tracts in multiple sclerosis: association with retinal thinning and visual disability. **J Neuroimaging** 21:e41–e49, 2011

3. Lilja Y, Gustafsson O, Ljungberg M, Starck G, Lindblom B, Skoglund T, et al: Visual pathway impairment by pituitary adenomas: quantitative diagnostics by diffusion tensor imaging. *J Neurosurg* **127**:569–579, 2017
4. Phal PM, Steward C, Nicholas AD, Kokkinos C, Desmond PM, Danesh-Meyer H, et al: Assessment of optic pathway structure and function in patients with compression of the optic chiasm: a correlation with optical coherence tomography. *Invest Ophthalmol Vis Sci* **57**:3884–3890, 2016
5. Schilling K, Gao Y, Janve V, Stepniewska I, Landman BA, Anderson AW: Confirmation of a gyral bias in diffusion MRI fiber tractography. *Hum Brain Mapp* **39**:1449–1466, 2018
6. Tournier JD, Calamante F, Connelly A: Robust determination of the fibre orientation distribution in diffusion MRI: non-negativity constrained super-resolved spherical deconvolution. *Neuroimage* **35**:1459–1472, 2007

Disclosures

Dr. Priti Balchandani is a named inventor on patents relating to MRI and RF pulse design. The patents have been licensed to GE Healthcare, Siemens AG, and Philips International. Dr. Balchandani receives royalty payments related to these patents.

INCLUDE WHEN CITING

Published online May 17, 2019; DOI: 10.3171/2019.3.JNS19487.

©AANS 2020, except where prohibited by US copyright law

Stimulating cholinergic neurons causes urinary incontinence after DBS of the GPi

TO THE EDITOR: In their case report, Liu et al.¹ reported on a woman with urinary incontinence following deep brain stimulation of the globus pallidus internus (GPi) (Liu FT, Lang LQ, Zhou RY, et al: Urinary incontinence following deep brain stimulation of the globus pallidus internus: case report. *J Neurosurg* [epub ahead of print February 22, 2019. DOI: 10.3171/2018.11.JNS181513]). In this report, the authors suggested that the urinary incontinence was induced by fibers projecting from the cortex to pontine micturition centers. Although this hypothesis is plausible, we believe that the incontinence was actually caused by the stimulation of cholinergic neurons.

Importantly, the patient's urinary incontinence disappeared after stimulating more dorsal contacts, suggesting that the problem was probably due to the most ventral contacts. Importantly, the most ventral contacts were observed outside of and interior to the GPi. Instead, we believe that these ventral contacts were associated with cholinergic neurons in the nucleus basalis of Meynert (NBM). Thus, stimulation of cholinergic neurons may actually underlie the induced urinary incontinence. Importantly, urinary incontinence is primarily treated by anticholinergic drugs, strengthening the case for a link between cholinergic stimulation and incontinence.^{2,3}

Furthermore, in this report, the incontinence was only associated with the left electrode. Although the reason for this location bias is unknown, it may be because the cholinergic neurons of the NBM are dispersed in this location. Indeed, although the NBM appears to be large in

brain atlases, targeting NBM cholinergic neurons, even intentionally, is difficult. In this report, the NBM may have been targeted only by chance.

Wei Liu, MD

Weihai Municipal Hospital, Capital Medical University, Shandong, China

Jian-Guo Zhang, MD, PhD

Beijing Tian-Tan Hospital, Beijing Neurosurgical Institute, Capital Medical University, Beijing, China

Acknowledgments

National Natural Science Foundation of China, No. 61761166004.

References

1. Liu FT, Lang LQ, Zhou RY, Feng R, Hu J, Wang J, et al: Urinary incontinence following deep brain stimulation of the globus pallidus internus: case report. *J Neurosurg* [epub ahead of print February 22, 2019. DOI: 10.3171/2018.11.JNS181513]
2. Lukacz ES, Santiago-Lastra Y, Albo ME, Brubaker L: Urinary incontinence in women. *JAMA* **318**:1592–1604, 2017
3. Wood LN, Anger JT: Urinary incontinence in women. *BMJ* **349**:g4531, 2014

Disclosures

The authors report no conflict of interest.

Correspondence

Jian-Guo Zhang: zjguo73@126.com.

INCLUDE WHEN CITING

Published online July 26, 2019; DOI: 10.3171/2019.3.JNS19587.

Response

We are very glad to receive the letter from Dr. Liu and Dr. Zhang in response to our case report. In their letter, they suggested that the incontinence might be caused by the stimulation of cholinergic neurons. However, we believe that this hypothesis is a long way from being proven.

As far as we know, the basal ganglia is involved in the micturition control pathways,¹ which have been discussed widely, but the detailed mechanism is still not completely clear. However, it might be a bit arbitrary to declare that the GPi stimulation–induced urinary incontinence was associated with cholinergic neurons in the NBM.

We suggest that Dr. Liu should provide more solid evidence. First, is the NBM involved in the normal micturition control? What is the function of the NBM in the cases of urinary incontinence or in patients with Parkinson disease?² Second, what are the functions of the NBM and the micturition reflexes in the cases with NBM stimulation? If the NBM is proved to be involved in the micturition reflexes, the authors should then prove that the fibers connecting the NBM pass by the neighboring region of the GPi, or that there are connections between the GPi and the NBM. After proving that, the authors should then prove that the responsible fibers were dopaminergic. Because the brain penetrance of most of the anticholinergic drugs in treating urinary incontinence was so low,³ the evidence listed by our colleagues was really weak. Meanwhile, the

electrical stimulation is nonselective in that the declaration of the dopaminergic involvement of the NBM in GPI-induced micturition needs more detailed evidence.

In our case, the woman developed urinary incontinence after a long period of stimulation of the GPI. After turning the stimulation off or stimulating more dorsal contacts, the urinary incontinence disappeared. In the whole process, the motor symptoms were well controlled. In this case, we suggested there were at least two possible reasons accounting for this phenomenon. First, the stimulation of GPI in our case spread to the neighboring regions ventral to the GPI, in which the micturition pathway might be stimulated. Second, the stimulation of the GPI itself might induce the urinary incontinence, and the responsible subregion of the GPI and related connectivity should be further explored. As reported in a previous study,⁴ the anatomical and functional connection of the GPI in micturition reflexes should be further explored.

In conclusion, the detailed mechanisms between GPI stimulation and urinary incontinence are still unclear. Investigators should focus on the roles of GPI and neighboring regions in micturition reflexes.

Feng-Tao Liu, MD
Jian-Jun Wu, MD

Huashan Hospital, Fudan University, Shanghai, China

References

1. Blanco L, Ros CM, Tarragon E, Fernandez-Villalba E, Herrero MT: Functional role of Barrington's nucleus in the micturition reflex: relevance in the surgical treatment of Parkinson's disease. **Neuroscience** 266:150–161, 2014
2. Lee DJ, Milosevic L, Gramer R, Sasikumar S, Al-Ozzi TM, De Vloo P, et al: Nucleus basalis of Meynert neuronal activity in Parkinson's disease. **J Neurosurg** [epub ahead of print February 22, 2019]. DOI: 10.3171/2019.3.JNS182386
3. Palma JA, Kaufmann H: Treatment of autonomic dysfunction in Parkinson disease and other synucleinopathies. **Mov Disord** 33:372–390, 2018
4. Yao J, Zhang Q, Liao X, Li Q, Liang S, Li X, et al: A corticopontine circuit for initiation of urination. **Nat Neurosci** 21:1541–1550, 2018

INCLUDE WHEN CITING

Published online July 26, 2019; DOI: 10.3171/2019.3.JNS19813.

©AANS 2020, except where prohibited by US copyright law

Risk of new seizure occurrence following cranioplasty procedures

TO THE EDITOR: The article by Yeap et al.⁷ was read with great interest (Yeap MC, Chen CC, Liu ZH, et al: Postcranioplasty seizures following decompressive craniectomy and seizure prophylaxis: a retrospective analysis at a single institution. *J Neurosurg* 131:936–940, September 2019). The authors highlight an important clinical issue on the prophylactic use of antiepileptics in cranioplasty surgery, a practice that seems to be efficacious in the prevention of new-onset seizures. According to Yeap and

colleagues, the incidence of and the various risk factors associated with postcranioplasty seizure have not been analyzed previously.⁷ However, the study by Yeap et al. does not represent the first study to assess the efficacy of antiepileptic drugs for seizure prophylaxis in cranioplasty. In 2018, Yao et al. performed a meta-analysis in which pooled incidences of postcranioplasty seizures ranged from 1.6% to 29%, early cranioplasty within 6 months of decompressive craniectomy was associated with an increased incidence of seizures, and the prophylactic use of antiepileptic drugs (AEDs) effectively reduced seizures.⁶ In 2013, Lee et al. analyzed 243 cranioplasties, and the incidence of new-onset postcranioplasty seizure was 14.81%,⁴ but these authors did not recommend routine prophylactic antiepileptic medications in view of the highly variable incidence of and various risk factors for seizures, that is, a history of previous trauma, precranioplasty neurological deficit, and intracranial hemorrhage. Yeap et al. also observed intraoperative blood loss, cirrhosis of the liver, and shunt-dependent hydrocephalus as risk factors.⁷ However, Krause-Titz observed that a poor neurological status, shunt-dependent hydrocephalus, and an advanced age correlate with new-onset seizure.² Sobani et al. observed that 15.6% of cases had seizures postcranioplasty.⁵

The incidence of new-onset seizures following cranioplasty remains highly variable, as the primary pathological indications for decompressive craniectomy remain highly heterogeneous, as do the seizure diagnostic criteria. As in the study by Liang et al.,³ Chen and colleagues¹ utilized both electroencephalography and clinical diagnostic criteria. In their randomized controlled study, Chen et al. observed that the incidence of epileptic seizures was significantly higher in patients who had not received antiepileptic prophylaxis treatment than in those who received AED treatment (28.6% vs 5.9%, respectively).¹

Yeap et al. should be commended for separately assessing early- and late-onset postcranioplasty seizures as both have different mechanisms of pathogenesis and treatment protocols.⁷ These authors proved that the prophylactic use of antiepileptic medications will lead to improvement in the overall neurological outcome and minimize morbidity. Similarly, Chen et al. also recommended the early use of AEDs to effectively reduce the occurrence of seizures.¹

Guru Dutta Satyarthee, MD, MCh

Neurosciences Centre, AIIMS, New Delhi, India

Luis R. Moscote-Salazar, MD

Latin-American Neurotrauma and Neurocritical Care (Red Latino),
Bogota, Colombia

References

1. Chen F, Duan Y, Li Y, Han W, Shi W, Zhang W, et al: Use of an antiepileptic drug to control epileptic seizures associated with cranioplasty: a randomized controlled trial. **Int J Surg** 40:113–116, 2017
2. Krause-Titz UR, Warneke N, Freitag-Wolf S, Barth H, Mehdorn HM: Factors influencing the outcome (GOS) in reconstructive cranioplasty. **Neurosurg Rev** 39:133–139, 2016
3. Liang S, Ding P, Zhang S, Zhang J, Zhang J, Wu Y: Prophylactic levetiracetam for seizure control after cranioplasty: a multicenter prospective controlled study. **World Neurosurg** 102:284–292, 2017

4. Lee L, Ker J, Quah BL, Chou N, Choy D, Yeo TT: A retrospective analysis and review of an institution's experience with the complications of cranioplasty. **Br J Neurosurg** 27:629–635, 2013
5. Sobani ZA, Shamim MS, Zafar SN, Qadeer M, Bilal N, Mur-taza SG: Cranioplasty after decompressive craniectomy: an institutional audit and analysis of factors related to complications. **Surg Neurol Int** 2:123, 2011
6. Yao Z, Hu X, You C: The incidence and treatment of seizures after cranioplasty: a systematic review and meta-analysis. **Br J Neurosurg** 32:489–494, 2018
7. Yeap MC, Chen CC, Liu ZH, Hsieh PC, Lee CC, Liu YT, et al: Postcranioplasty seizures following decompressive craniectomy and seizure prophylaxis: a retrospective analysis at a single institution. **J Neurosurg** 131:936–940, 2019

Disclosures

The authors report no conflict of interest.

Correspondence

Guru Dutta Satyarthee: duttaguru2002@yahoo.com.

INCLUDE WHEN CITING

Published online December 13, 2019; DOI: 10.3171/2019.9.JNS182980.

Response

We thank Drs. Satyarthee and Moscote-Salazar for their great commendation. Indeed, the incidence of postcranioplasty seizure remains high and is associated with a poor outcome and longer hospital stays. In the literature, as mentioned by Drs. Satyarthee and Moscote-Salazar, some authors have reported a high risk of postcraniotomy seizure.^{2–5} However, when we launched our analysis, there was still no study that had demonstrated the effects of prophylactic AED treatments. We showed that new-onset seizures occurred in 26.5% of patients and that early seizures occurred in 7.4% of patients. Our protocol was actually very simple and effective. For patients who had no history of seizure, prophylactic AEDs were administered for 1 week. Management was based on common strategies for posttraumatic seizures or postcraniotomy prophylaxis. Recently, as presented in another of our studies,¹ patients who had a history of previous seizure and who had already taken AEDs, the administration of AEDs was changed to an intravenous formula perioperatively and then shifted back to an oral form at least 3 days later. The results reduced

the postcranioplasty early recurrent seizure rate (20.5% vs 8.9%). However, because our two studies were retrospective in nature, there are some limitations. First, we could not know how long the AEDs should have been used or how effective they were. In Liang and colleagues' series,³ prophylactic AEDs were used for 2 weeks. In our experience, only 17.4% of patients who experienced recurrent seizures after cranioplasty experienced early recurrent seizures within 1 week postoperatively, at a median of 2 days (IQR 1–4 days).¹ Therefore, in the future, a prospective multicenter study assessing early-onset, late-onset, and recurrent seizures should be performed.

Again, thank you for your great interest and commendations regarding our work.

Ching-Chang Chen, MD
Po-Hsun Tu, MD

Chang Gung Memorial Hospital, Linkou Medical Center, Chang Gung Medical College & University, Taoyuan City, Taiwan

References

1. Chen CC, Yeap MC, Liu ZH, Hsieh PC, Chen CT, Liu YT, et al: A novel protocol to reduce early seizures after cranioplasty: a single-center experience. **World Neurosurg** 125:e282–e288, 2019
2. Lee L, Ker J, Quah BL, Chou N, Choy D, Yeo TT: A retrospective analysis and review of an institution's experience with the complications of cranioplasty. **Br J Neurosurg** 27:629–635, 2013
3. Liang S, Ding P, Zhang S, Zhang J, Zhang J, Wu Y: Prophylactic levetiracetam for seizure control after cranioplasty: a multicenter prospective controlled study. **World Neurosurg** 102:284–292, 2017
4. Walcott BP, Kwon CS, Sheth SA, Fehnel CR, Koffie RM, Asaad WF, et al: Predictors of cranioplasty complications in stroke and trauma patients. **J Neurosurg** 118:757–762, 2013
5. Zanaty M, Chalouhi N, Starke RM, Clark SW, Bovenzi CD, Saigh M, et al: Complications following cranioplasty: incidence and predictors in 348 cases. **J Neurosurg** 123:182–188, 2015

INCLUDE WHEN CITING

Published online December 13, 2019; DOI: 10.3171/2019.10.JNS192730.

©AANS 2020, except where prohibited by US copyright law