Neurosurgical Forum

LETTERS TO THE EDITOR

Tumor extension determines approach in insular gliomas

TO THE EDITOR: We read with great interest the series by Przybylowski et al.³ on the relative morbidity of the transsylvian versus transcortical approach to insular gliomas (Przybylowski CJ, Baranoski JF, So VM, et al: Surgical morbidity of transsylvian versus transcortical approaches to insular gliomas. *J Neurosurg* [epub ahead of print April 5, 2019. DOI: 10.3171/2018.12.JNS183075]). At the outset we congratulate the authors for reporting such a large series of insular gliomas. They also need to be commended for using volumetric assessment for both low-grade and high-grade gliomas as well as diffusion-weighted imaging in all postoperative MRI scans to determine infarcts in these cases, something that we believe is often underreported. As reported by them, these lesions are notoriously difficult to excise without subjecting the patients to risks of deficits. Both the approaches described have been used in this series for all 4 zones, although the inferior posterior area (zone III) was predominately treated via the transcortical approach ($p < 0.01$) and the inferior anterior area (zone IV) was approached predominately via the transsylvian approach ($p = 0.17$). Herein lies the conundrum—should location of the tumor decide the approach, or are both approaches equivalent in their results, apart from the superior posterior quadrant (zone II), as concluded in this article?

Sylvian fissure opening may not be easy in all cases of insular gliomas, particularly when the opercula are involved by the tumor. Most neurosurgeons would agree that it is practically impossible to open the sylvian fissure distally, rendering visualization of zone IV extremely difficult, as duly noted by the authors themselves.² An exception is generally observed with exophytic insular gliomas, which tend to autodissect the sylvian fissure.¹ Therefore, the transcortical approach is the only option left in the majority of the sizable insular gliomas. We as well as other authors have often observed that removal of the middle part of the superior temporal gyrus, especially on the nondominant side, provides excellent exposure of zone II.

When going transsylvian, we believe a wide sylvian split is necessary to allow the surgeon to work inside the insula without having to retract the opercula. Although the strategy of sectoral opening of the sylvian fissure used by

the authors seems appealing, a wide exposure and subsequent multiangle maneuverability of the tumor has been advocated by others too.^{4,5}

The authors mention that they use awake craniotomy to map the periinsular cortices during the transcortical approach, whereas all transsylvian approaches are performed under general anaesthesia. We believe that awake craniotomy and intraoperative monitoring (clinical and electrophysiological) are desirable in all patients with insular glioma. This point has been strongly emphasized by the Duffau group too.² In our practice, awake craniotomy and intraoperative monitoring are used in all patients with tumor on the dominant side, irrespective of whether the transsylvian or transopercular approach is used. On the nondominant side, we have observed that subcortical stimulation with the patient off muscle relaxants under general anaesthesia is a reasonable approach when awake craniotomy is not feasible for any reason. High-grade gliomas with extensive infiltration and accompanying brain bulge may limit awake craniotomy use in some cases. We also agree that the high-grade gliomas tend to be infiltrative, particularly of the lenticulostriate perforators. Moreover, a propensity to bleed during surgery also increases the risk of inadvertent vascular injury while trying to establish hemostasis (either by coagulation or by trying to remove the bleeding tumor).

So, the type of insular glioma accessed by the two approaches are actually different, and hence their comparison may not be appropriate. Many of these tumors are beyond the insula by the time they reach a neurosurgeon; more often than not, transcortical remains the only possible approach. Moreover, as Duffau et al.² have shown, the transcortical approach may allow better seizure freedom and fewer retraction-related injuries, apart from less vascular manipulation. On the other hand, small (Yaşargil type 3A) tumors can be very well addressed by the transsylvian route. We think tumors larger than this are best approached by a transcortical approach. In our experience, the transsylvian approach in larger tumors not only leads to insufficient tumor exposure but also endangers the vascular structures. Moreover, it has been our experience that eventually another corridor (transcortical) becomes necessary to visualize deeper areas of tumor extension to avoid undue opercular retraction. However, we find one advantage of early sylvian fissure split (deeper sphenoidal part) even when using a transcortical approach, which includes

the ability to identify the medial-most lenticulostriate perforator extrapially early on in the surgery. Thereafter we proceed with transcortical tumor decompression and often try to visualize (or determine the location of) the lenticulostriate arteries through the opened-up sylvian fissure intermittently as we go along. This appears much safer to us than visualizing the lenticulostriate arteries through the tumor, which acts as a resection limit medially.

Therefore, according to our understanding, the surgical approach is often dictated by the tumor size and extension rather than the surgeon's preference.

Harsh Deora, MCh, DNB

National Institute of Mental Health and Neurosciences (NIMHANS),
Bangalore, India

Kuntal Kanti Das, MCh

Awadhesh Jaiswal, MCh

Sanjay Behari, MCh, DNB

Sanjay Gandhi Post Graduate Institute of Medical Education and
Research, Lucknow, India

References

1. Das KK, Singh S, Deora H, Khatri D, Mehrotra A, Srivastava AK, et al: Microsurgical excision of giant dominant lobe insular cavernoma presenting acutely: Sometimes you win, sometimes you learn. **Interdiscip Neurosurg** **18**:100475, 2019
2. Duffau H, Capelle L, Lopes M, Faillot T, Sichez JP, Fohanno D: The insular lobe: physiopathological and surgical considerations. **Neurosurgery** **47**:801–811, 2000
3. Przybylowski CJ, Baranoski JF, So VM, Wilson J, Sanai N: Surgical morbidity of transsylvian versus transcortical approaches to insular gliomas. **J Neurosurg** [epub ahead of print April 5, 2019. DOI: 10.3171/2018.12.JNS183075]
4. Rey-Dios R, Cohen-Gadol AA: Technical nuances for surgery of insular gliomas: lessons learned. **Neurosurg Focus** **34**(2):E6, 2013
5. Yaşargil MG: **Microneurosurgery: Microsurgical Anatomy of the Basal Cisterns and Vessels of the Brain**. Stuttgart: Georg Thieme, 1984

Disclosures

The authors report no conflict of interest.

Correspondence

Kuntal Kanti Das: kkdas@sgpgi.ac.in.

INCLUDE WHEN CITING

Published online August 23, 2019; DOI: 10.3171/2019.5.JNS191308.

Response

We thank Deora and colleagues for their letter to the editor in response to our publication.

We agree with the authors that gliomas confined purely to the anterior insular region can be safely removed through the transsylvian route. As intraoperative mapping techniques have advanced, our practice has evolved to favor the awake transcortical approach for insular glioma resection. Overall, we agree that operating through windows of nonfunctional cortex allows for safer and larger exposures of these lesions, often leading to greater extent of resection without increased neurological morbidity.^{1–3}

As demonstrated in our study, we have preferred the transcortical approach for larger insular lesions, as the authors imply. However, we have also demonstrated that posterior insular location is an important factor that favors the transcortical approach. Because retraction of the frontal operculum or sacrifice of superficial sylvian veins is often required to gain adequate access to the posterior insula via the transsylvian route, we now use the transcortical approach for all posterior insular gliomas, independent of tumor size. As such, tumor location, in addition to tumor size, should continue to influence the neurosurgeon's decision on how to access insular gliomas.

Colin J. Przybylowski, MD

Nader Sanai, MD

Barrow Neurological Institute, St. Joseph's Hospital and Medical Center,
Phoenix, AZ

References

1. Benet A, Hervey-Jumper SL, Sánchez JJ, Lawton MT, Berger MS: Surgical assessment of the insula. Part 1: surgical anatomy and morphometric analysis of the transsylvian and transcortical approaches to the insula. **J Neurosurg** **124**:469–481, 2016
2. Duffau H, Moritz-Gasser S, Gatignol P: Functional outcome after language mapping for insular World Health Organization Grade II gliomas in the dominant hemisphere: experience with 24 patients. **Neurosurg Focus** **27**(2):E7, 2009
3. Sanai N, Polley MY, Berger MS: Insular glioma resection: assessment of patient morbidity, survival, and tumor progression. **J Neurosurg** **112**:1–9, 2010

INCLUDE WHEN CITING

Published online August 23, 2019; DOI: 10.3171/2019.5.JNS191361.

©AANS 2020, except where prohibited by US copyright law

Harvey Cushing's legacy

TO THE EDITOR: We read with great enthusiasm the article by Al-Mefty et al.¹ on Harvey Cushing's surgical neurology (Al-Mefty O, Laws ER, Popp AJ: Surgical neurology: Harvey Cushing's endangered legacy. *J Neurosurg* [epub ahead of print May 3, 2019. DOI: 10.3171/2019.1.JNS182290]). While the historical analysis of famous physicians was brilliant, we felt that the paper could have benefitted from analyzing how scientific societies develop a great neurosurgeon and relating historical findings to the current state of neurosurgery.¹ The authors reflect on neurosurgery as it was envisioned by our forebears. The historical accounts of many famous neurosurgeons were discussed, and the paper ends by warning against the diminution of the neurosurgeon to a mere technician.

We believe that the paper could have been further developed by evaluating the patterns of ingenuity that made neurosurgery flourish. The paper brilliantly recounts the achievements of Pierre Paul Broca. However, by mentioning his involvement in the *Société Anatomique de Paris*, the authors could have helped the reader understand how academic meetings can serve to develop physicians and in time result in medical breakthroughs. It was in just such

an environment that Broca and other notable physicians such as Jean-Martin Charcot (the father of neurology) were able to create a culture of academia. By emphasizing this, the authors could have helped current neurosurgeons comprehend the relevance of academic societies today.

The paper emphasizes a holistic approach to neurosurgery, and while we completely agree, we want to highlight the advantages of subspecialization in neurosurgery: improved patient mortality and morbidity.⁴ The “staggering expenditure on healthcare, coupled with technological advances” is also quoted as the cause of the diminishing role of the neurosurgeon. However, it is unclear if the paper references neurosurgeons in the United States or worldwide. Consider, for instance, that Japan and the United States each have a unique healthcare footprint.⁵ Moreover, modern technological advances such as MRI provide neurosurgeons with valuable information concerning brain pathologies. Spinal cord stimulation systems can help to reduce the use of opioids,^{2,7} and deep brain stimulation can help in the treatment of Parkinson’s disease.⁶ Therefore, we consider technological advances and subspecialization favorable to neurosurgery.

In our neurosurgery department, we herald Cushing’s introduction of the silver clip, so we would like to highlight here the gravity and ingenuity of such a simple and effective hemostatic method that we still use today.³

Google Trends shows that Harvey Cushing’s contributions are still remembered not only by the medical community but also by an international audience as Germany holds the highest interest while the United States is barely in 13th place. Moreover, searches for his name have increased by 150% over the last 10 years.

Regardless of our assessment, the study by Al-Mefty et al. offers a crucial reminder to the neurosurgical community to reflect on the principles that guide its everyday practice, as the past can often inspire the future. Harvey Cushing’s legacy should serve to instill the core values of neurosurgery as we all aim to uphold this very special branch of medicine.

Tomasz Szmuda, MD, PhD
Shan Ali

Paweł Słoniewski, MD, PhD
Medical University of Gdansk, Poland

Acknowledgments

Shan Ali would like to thank Tomasz Szmuda for mentoring him through medical school and ingraining in him the core values of neurosurgery. Tomasz Szmuda would like to thank Paweł Słoniewski for his continuous mentorship.

References

1. Al-Mefty O, Laws ER, Popp AJ: Surgical neurology: Harvey Cushing’s endangered legacy. **J Neurosurg** [epub ahead of print May 3, 2019]. DOI: 10.3171/2019.1.JNS182290
2. Deer T, Kim C, Bowman R, Ranson M, Douglas CS, Tolentino W: Spinal cord stimulation as a method of reducing opioids in severe chronic pain: a case report and review of the literature. **W J Med J** 106:56–59, 2010
3. Duane W: A modification of the McKenzie silver clip. **J Neurosurg** 7:92–93, 1950
4. McCutcheon BA, Hirshman BR, Gabel BC, Heffner MW, Marcus LP, Cole TS, et al: Impact of neurosurgeon spe-

cialization on patient outcomes for intracranial and spinal surgery: a retrospective analysis of the Nationwide Inpatient Sample 1998–2009. **J Neurosurg** 128:1578–1588, 2018

5. Ruggles BM, Xiong A, Kyle B: Healthcare coverage in the US and Japan. **Nursing (Lond)** 49:56–60, 2019
6. Schiefer TK, Matsumoto JY, Lee KH: Moving forward: advances in the treatment of movement disorders with deep brain stimulation. **Front Integr Neurosci** 5:69, 2011
7. Verrills P, Sinclair C, Barnard A: A review of spinal cord stimulation systems for chronic pain. **J Pain Res** 9:481–492, 2016

Disclosures

The authors report no conflict of interest.

Correspondence

Shan Ali: shanali@gumed.edu.pl.

INCLUDE WHEN CITING

Published online September 27, 2019; DOI: 10.3171/2019.6.JNS191711.

Response

We welcome Dr. Szmuda and colleagues’ comments and are very pleased that our article has inspired thoughts and discussion about the future direction of neurosurgery, as we had hoped. Indeed, neurosurgery has flourished in a scientific environment and will continue to do so with the interaction and progress of other disciplines. This places a heavy responsibility on the neurosurgical leaders at academic medical centers to innovate and advance the field and inspire their trainees beyond the practice of daily care. We hold high and espouse the neurosurgeon-scientist and believe this breed to be vital to the survival and advancement of our field. Being a scientist and understanding the physiology, anatomy, pathology, and other aspects related to the nervous system in health and disease were mandated by Cushing as a prerequisite to the practice and development of neurosurgery, as we quoted in our article:

To successfully cope with the many operative problems offered by the various disorders of the nervous system, a man, after a thorough training in pathology and medicine (in its broadest sense) must study, not only in the neurological clinic but also in the laboratory, the pathology of these afflictions in their histological and—what is more important—in their experimental aspects.¹

Moreover, prior to his contributions to the surgical aspect of neurosurgery, in 1901, Cushing invested time in Europe in the laboratory of Dr. Hugo Kronecker in Berne, Switzerland, and Angelo Mosso in Italy, researching brainstem control of systemic blood pressure.³ Cushing revealed in the laboratory what is coined as the “Cushing reflex” way before his design of a surgical instrument, the “Cushing forceps.” He saw such efforts as pivotal in developing neurosurgery:

The real leaders of today in surgery who owe their place not to any special brilliancy in operative manipulations, but to their laborious investigations of certain problems of disease, whereby has been disclosed a rational mechanical basis for a surgical therapy which can then be safely and successfully adapted by their many followers.²

We believe that the early founders of our specialty were knowledgeable scientists who collaborated with the lumi-

naries around them. We maintain that the neurosurgeon-scientist plays a critical role in the future of our specialty. We are blessed that our specialty attracts the finest minds in medicine that can carry on the torch. And indeed, the neurosurgeon is fortunate to benefit from the surrounding multidimensional advances.

Neurosurgery has risen from, benefitted from, and contributed to the magnificent advancement in technology and neuroscience. With this naturally comes growth, branching, and multidisciplinary collaboration. Our tree now holds more fruit. We should resist cutting off the branches, which will become unviable fragments of narrow interest, with the potential danger of becoming limited not only to procedures, but even to a specific procedure. A carpenter would not be fulfilled by only hammering nails while solely crafting chairs.

Ossama Al-Mefty, MD

Edward R. Laws Jr., MD

Brigham & Women's Hospital, Boston, MA

A. John Popp, MD

Albany Medical College, Albany, NY

Acknowledgments

We thank Dr. Linda Bi for editing assistance.

References

1. Cushing H: The special field of neurological surgery. **Bull Johns Hopkins Hosp** 16:77–87, 1905
2. Cushing H: The special field of neurological surgery; five years later. **Bull Johns Hopkins Hosp** 21:325–339, 1910
3. Greenblatt SH, Smith DC: The emergence of Cushing's leadership: 1901 to 1920, in Greenblatt SH, Dagi TF, Epstein MH (eds): **A History of Neurosurgery in Its Scientific and Professional Contexts**. Park Ridge, IL: American Association of Neurological Surgeons, 1997, pp 167–190

INCLUDE WHEN CITING

Published online September 27, 2019; DOI: 10.3171/2019.8.JNS191768.

©AANS 2020, except where prohibited by US copyright law

Multimodal management for benign skull base meningiomas

TO THE EDITOR: We read with great interest the article by Kim et al.,³ in which they discussed multimodal management of petroclival meningiomas (Kim JW, Jung HW, Kim YH, et al: Petroclival meningiomas: long-term outcomes of multimodal treatments and management strategies based on 30 years of experience at a single institution. *J Neurosurg* [epub ahead of print May 10, 2019. DOI: 10.3171/2019.2.JNS182604]). The current authors' preference is for multimodal management, with planned subtotal resection followed by adjuvant stereotactic radiosurgery (SRS).

During the past 10 years, our unit has developed an optimally invasive skull base surgery approach for large benign tumors. The philosophy of this approach is planned

subtotal resection followed by SRS using Gamma Knife surgery (GKS), with the aim to reduce morbidity and mortality. GKS is usually performed between 3 and 6 months after microsurgical resection, without a “wait-and-scan” follow-up period of the remnant tumor. In relation to this recently published paper by Kim and coauthors, we also apply this approach to other skull base meningiomas, such as the clinoid and cavernous sinus ones.²

In the particular case of petroclival meningiomas radical removal remains challenging, primarily due to brainstem adherence or invasion, and encasement of basilar artery/perforators, as well as cranial nerves (CNs) V, VII, VIII, IX, X, and XI.⁶ In these cases we prefer to use the supracerebellar infratentorial (extended retrosigmoid) approach. The extent of tumor removal frequently depends on the extent of clival component, invasion of cavernous sinus, and adherence to brainstem pia and the basilar artery or its branches. What is deemed risky to remove is left behind for subsequent GKS-SRS. In our series of combined approaches for benign skull base lesions, we have not had any mortality or major neurological morbidity. Single-fraction GKS can be safely applied on residual tumors,¹ even in cases with brainstem contact, due to low prescription marginal doses of 12 Gy, which allow for high rates of tumor control.⁷ In cases in which the limiting factor is the optic apparatus, due to the steep gradient of GKS it is possible to keep the dose below 8 Gy.^{4,5}

In the particular case of clinoid and cavernous sinus meningiomas, the surgical challenge is to preserve the carotid artery and its branches, recover visual function, and limit oculomotor CN palsies. We do not routinely remove the intracavernous sinus part for these tumors. We also perform chiasmopexy, by interposing a small fat graft between the optic nerve, chiasm, and residual intracavernous sinus tumor component, allowing safe GKS on the remnant. The same strategy is applied to pituitary stalk whenever needed, to minimize the risk of pituitary insufficiency.

In conclusion, we are completely in agreement and we congratulate Kim and coauthors for a very nice paper, which will contribute, in the future, to a major paradigm shift in the treatment of these difficult skull base tumors.

Constantin Tuleasca, MD, PhD¹⁻⁵

Lorenzo Giammattei, MD¹

Roy Thomas Daniel, MBBS, MCh^{1,2}

Marc Levivier, MD, PhD, IFAANS^{1,2}

¹Neurosurgery Service and Gamma Knife Center, Lausanne University Hospital, Lausanne, Switzerland

²Faculty of Biology and Medicine, University of Lausanne, Lausanne, Switzerland

³Signal Processing Laboratory, Ecole Polytechnique Fédérale de Lausanne, Lausanne, Switzerland

⁴Faculté de Médecine, Sorbonne Université, Paris, France

⁵Service de Neurochirurgie, Assistance Publique-Hôpitaux de Paris, Hôpitaux Universitaires Paris-Sud, Centre Hospitalier Universitaire Bicêtre, Paris, France

Acknowledgments

We acknowledge Lausanne University Hospital and University of Lausanne. Constantin Tuleasca gratefully acknowledges receipt of a “Young Researcher in Clinical Research Grant”

(Jeune Chercheur en Recherche Clinique) from the University of Lausanne (UNIL), Faculty of Biology and Medicine (FBM), and the Lausanne University Hospital (CHUV).

References

1. Faramand A, Kano H, Niranjan A, Johnson SA, Hassib M, Park KJ, et al: Cranial nerve outcomes after primary stereotactic radiosurgery for symptomatic skull base meningiomas. *J Neurooncol* **139**:341–348, 2018
2. Giammattei L, Starnoni D, Levivier M, Messerer M, Daniel RT: Surgery for clinoidal meningiomas: case series and meta-analysis of outcomes and complications. *World Neurosurg* **129**:e700–e717, 2019
3. Kim JW, Jung HW, Kim YH, Park CK, Chung HT, Paek SH, et al: Petroclival meningiomas: long-term outcomes of multimodal treatments and management strategies based on 30 years of experience at a single institution. *J Neurosurg* [epub ahead of print May 10, 2019. DOI: 10.3171/2019.2.JNS182604]
4. Leavitt JA, Stafford SL, Link MJ, Pollock BE: Long-term evaluation of radiation-induced optic neuropathy after single-fraction stereotactic radiosurgery. *Int J Radiat Oncol Biol Phys* **87**:524–527, 2013
5. Pollock BE, Link MJ, Leavitt JA, Stafford SL: Dose-volume analysis of radiation-induced optic neuropathy after single-fraction stereotactic radiosurgery. *Neurosurgery* **75**:456–460, 2014
6. Seifert V: Clinical management of petroclival meningiomas and the eternal quest for preservation of quality of life: personal experiences over a period of 20 years. *Acta Neurochir (Wien)* **152**:1099–1116, 2010
7. Tulasca C, Leroy HA, Régis J, Levivier M: Gamma Knife radiosurgery for cervical spine lesions: expanding the indications in the new era of Icon. *Acta Neurochir (Wien)* **158**:2235–2236, 2016

Disclosures

The authors report no conflict of interest.

Correspondence

Constantin Tulasca: constantin.tulasca@gmail.com.

INCLUDE WHEN CITING

Published online November 15, 2019; DOI: 10.3171/2019.6.JNS191682.

Response

It is our great pleasure to send this response to the letter from Dr. Tulasca. As mentioned in our article, the treatment modality of petroclival meningioma is still under debate—and for certain surgical approaches such as the transpetrosal approach, which allows shorter distance to the tumor, broadening of the surgical field, and minimizing brain retraction, complete resection of the lesion appears to be beneficial. However, considering the changes in the treatment paradigm, it is important for surgeons to devise the most optimal treatment plan to enhance the patients' clinical outcome postoperatively. Hence, we believe that multimodal treatments including adjuvant SRS or radiation therapy can improve the patients' clinical outcome by reducing the risks associated with surgery alone.

In response to adjuvant GKS, during the period of 3 to 6 months after the resection, we perform fractionated or single-fraction GKS based on the postoperative imaging follow-up. As Dr. Tulasca mentioned, single-fraction GKS can be used even if the residual tumor is in contact

with the brainstem. However, if major CN pathways are compromised by the tumor, we consider fractionated GKS. Although it is not set as a routine, usually for lesions in contact with the optic nerve we limit the maximum amount of radiation to 8 Gy and for fractionated GKS we use 5.5–5.7 Gy at each session. For single-fraction GKS, small tumors are controlled with 14 Gy and large tumors are treated with 11 Gy.

As to Dr. Tulasca's remark on cavernous and clinoid meningiomas, we are also expanding our indications for multimodal treatments not confined to the petroclival area, and we are observing promising results.

In conclusion, we would like to acknowledge Dr. Tulasca for the thorough response to our work. We believe such reviews can facilitate the adoption of multimodal treatment in petroclival meningioma and furthermore in difficult skull base tumors.

Sang Hyung Lee, MD, PhD

Seoul Metropolitan Government–Seoul National University Boramae Medical Center, Seoul, Korea

Jin Wook Kim, MD, PhD

Hyun-Tai Chung, PhD

Seunghoon Lee, MD

Seoul National University Hospital, Seoul, Korea

INCLUDE WHEN CITING

Published online November 15, 2019; DOI: 10.3171/2019.8.JNS191793.

©AANS 2020, except where prohibited by US copyright law

Preoperative embolization of brain arteriovenous malformations

TO THE EDITOR: We read with great interest the article by Dr. Donzelli and colleagues³ that describes their experience with preoperative embolization of brain arteriovenous malformations (AVMs) (Donzelli GF, Nelson J, McCoy D, et al: The effect of preoperative embolization and flow dynamics on resection of brain arteriovenous malformations. *J Neurosurg* [epub ahead of print May 17, 2019. DOI: 10.3171/2019.2.JNS182743]). This retrospective study found that preoperative embolization was associated with longer median resection times and that it did not affect intraoperative blood loss or the modified Rankin Scale score.

Although we admit that it is very difficult to stratify data in terms of safety and effectiveness of preoperative embolization because of the heterogeneous landscape of brain AVMs, our extensive experience has progressively led us to select only grade III Spetzler-Martin AVMs for preoperative embolization.^{2,5} An exception would be medium-eloquent lesions (grade III+, S2V0E1), for which a conservative treatment is recommended because the surgical risk is similar to that of high-grade AVMs.^{4,5} Apart from the obvious need for a detailed preoperative angiographic study of the AVM that must imperatively be completed through superselective microcatheterization of the nidus, it is our opinion that 3 other essential aspects

must be considered before drawing definitive conclusions about the effectiveness and safety of preoperative embolization—namely, 1) type of embolic material used; 2) time frame between embolization sessions; and 3) time to resection after embolization. In the article by Dr. Donzelli et al., *N*-butyl 2-cyanoacrylate was the most common embolic agent used (53% of cases); however, reports in the literature support the superiority of low-density ethylene vinyl alcohol (EVOH) copolymer with respect to efficacy of embolization, especially for deeper arterial feeders.^{1,5} Next, our experience also leads us to recommend a time frame of 10–15 days between embolization sessions to accommodate the well-known risk of dynamic recruitment of new arterial feeders. Furthermore, we suggest that resection be performed within 7 days of embolization, because longer time frames may nullify the valuable effects of preoperative staged embolization or even negatively affect surgery.

Given the above, it is reasonable to assume that a detailed analysis of these factors within this large series can lead to slightly different interpretations of the effectiveness of preoperative embolization of brain AVMs. Therefore, in contrast to the conclusion reached by Donzelli et al., we are firmly convinced that preoperative embolization may be capable of downgrading the AVM from a functional vascular standpoint rather than a volumetric one, only in select grade III AVMs, which would ultimately make the surgery easier and safer. Nonetheless, we completely agree with Dr. Donzelli and colleagues on the utility of the iFlow parametric color-coding quantitative technique for preoperatively assessing AVM flow. We hope that the authors will further analyze data pertaining to these significant aspects in their large and therefore precious case series.

Sabino Luzzi, MD, PhD^{1,2}

Mattia Del Maestro, MD²

Renato Galzio, MD^{1,2}

¹University of Pavia, Italy

²Fondazione IRCCS Policlinico San Matteo, Pavia, Italy

References

1. Crowley RW, Ducruet AF, Kalani MYS, Kim LJ, Albuquerque FC, McDougall CG: Neurological morbidity and mortality associated with the endovascular treatment of cerebral arteriovenous malformations before and during the Onyx era. **J Neurosurg** **122**:1492–1497, 2015
2. Del Maestro M, Luzzi S, Gallieni M, Trovarelli D, Giordano AV, Gallucci M, et al: Surgical treatment of arteriovenous malformations: role of preoperative staged embolization. **Acta Neurochir Suppl** **129**:109–113, 2018
3. Donzelli GF, Nelson J, McCoy D, McCulloch CE, Hetts SW, Amans MR, et al: The effect of preoperative embolization and flow dynamics on resection of brain arteriovenous malformations. **J Neurosurg** [epub ahead of print May 17, 2019. DOI: 10.3171/2019.2.JNS182743]
4. Lawton MT: Spetzler-Martin Grade III arteriovenous malformations: surgical results and a modification of the grading scale. **Neurosurgery** **52**:740–749, 2003
5. Luzzi S, Del Maestro M, Bongetta D, Zoia C, Giordano AV, Trovarelli D, et al: Onyx embolization before the surgical treatment of grade III Spetzler-Martin brain arteriovenous malformations: single-center experience and technical nuances. **World Neurosurg** **116**:e340–e353, 2018

Disclosures

The authors report no conflict of interest.

Correspondence

Sabino Luzzi: sabino.luzzi@unipv.it.

INCLUDE WHEN CITING

Published online December 6, 2019; DOI: 10.3171/2019.6.JNS191541.

Response

We thank Drs. Luzzi, Del Maestro, and Galzio for their interest in our article and for their thoughtful comments. It is our hope that our study will spark discussion regarding the selection of patients with AVM for embolization. Our study is retrospective and by no means definitive, with the possibility that we did not collect and incorporate all pertinent variables associated with surgical outcomes in our analysis. Nonetheless, we have attempted to address our correspondents' key concerns below.

The correspondents note that their experience has led them to select only grade III Spetzler-Martin AVMs for embolization. When we restricted our models to grade III Spetzler-Martin AVMs ($n = 58$), we found a trend toward decreased operative time (–21 minutes; 95% CI –80 to +38 minutes, $p = 0.486$), and a trend toward decreased blood loss (proportional increase 0.84; 95% CI 0.54 to 1.31, $p = 0.436$). Although these results do not meet any reasonable threshold of statistical significance, the direction of effect of these coefficients is in agreement with the correspondents' observations.

The correspondents also note that the embolization material should be considered, citing the superiority of EVOH. To address this, we compared outcomes for cases that used EVOH as the embolic material ($n = 12$) to *N*-butyl 2-cyanoacrylate ($n = 71$; the most common material used in our series). We found a trend toward shorter resection times for EVOH (–24 minutes; 95% CI –107 to +58 minutes; $p = 0.562$), as well as less blood loss (proportional increase 0.75; 95% CI 0.41 to 1.38, $p = 0.359$). Again, the coefficients' direction of effect was in agreement with the correspondents' observations, but we lack the statistical power to make any strong claims. Unfortunately, we were unable to explore the role of time between embolizations and time from embolization to surgery due to limitations of the data set.

The use of preoperative embolization in general is poorly studied whether it is for AVMs, tumors, or other vascular anomalies. Such practices are inconsistent and driven largely by personal experience rather than formal study. Given emerging interest in primary endovascular treatment of brain AVMs, however, there is promise for better guidance to the neurointerventional community. Further study would help identify subsets of patients for whom preoperative embolization may be of benefit as well as the materials and techniques that are best suited for this procedure. We believe that this selection is critical, especially because our study suggests that, as uniformly applied, preoperative embolization may not have a clearly beneficial effect on surgical outcomes. As a start, the community would benefit from improving embolization nomenclature to better define both the purpose and meth-

ods used in a treatment. Our ability to better communicate such surgical subclassification strategies helps bring into focus our goals as they relate to research and clinical care.

Grace F. Donzelli, MD

Jeffrey Nelson, MS

Helen Kim, PhD

Daniel L. Cooke, MD

UCSF Medical Center, San Francisco, CA

INCLUDE WHEN CITING

Published online December 6, 2019; DOI: 10.3171/2019.8.JNS191659.

©AANS 2020, except where prohibited by US copyright law

Do bacteria contribute to formation and rupture of intracranial aneurysms?

TO THE EDITOR: We read the article by Aboukais et al.¹ with great interest (Aboukais R, Loiez C, Leclerc X, et al: Absence of bacteria in intracranial aneurysms. *J Neurosurg* [epub ahead of print March 1, 2019. DOI: 10.3171/2018.12.JNS183044]).

First, we would like to congratulate the authors for trying to identify other inherent risks factors associated with intracranial aneurysm (IA) formation and rupture.

Aboukais et al. conducted a prospective study of 30 French patients treated between January and July 2018, including analysis of specimens from 10 ruptured and 20 unruptured IA walls obtained intraoperatively after microsurgical clipping. In addition, the authors compared their results with those of previous Finnish studies^{4–6} in which analyses of odontogenic bacteria showed an association of oral infection and saccular IA formation and rupture.^{4,5} However, some methodologic and analytical issues should be discussed.

The data in the study by Aboukais et al. may have been compromised due to a low number of cases (n = 30) managed in only one center, in comparison with the 70 cases examined in the Finnish study.⁴ Moreover, Aboukais et al. did not use a control group to compare the ruptured and unruptured IA data for the study patients to the general population.

These authors argue that the higher prevalence of bacteria found in the Finnish study aneurysms is due to more frequent tooth brushing in this population. However, to our knowledge, there are no supporting data for this assertion in the literature. Nevertheless, Aboukais et al. suggest that brushing teeth regularly prevents periodontitis and that possible microtrauma formation caused by brushing is not sufficient to explain the transitory bacteremia, and therefore it is likely that these microtraumas bias inflammatory changes and facilitate the movement of bacteria into the blood. This explanation, in our opinion, may be due to an ecological inference fallacy, because periodontitis infection can change the mouth microbiota and induce systemic inflammatory changes. However, we believe that

Aboukais et al. reached this false-negative finding because their investigation only analyzed 2 bacteria species, *Escherichia coli* and *Staphylococcus aureus*, neither of which is commonly found in the oral cavity. In our experience, other pathogens such as *Fusobacterium* and *Treponema* are prevalent in the oral cavity.

Several factors can influence IA formation and rupture, and studies like those we discuss here can give rise to a new understanding of the underlying pathophysiology.^{4,5} Periodontitis and periapical lesions facilitate chronic inflammatory changes, and inflammation modifies the microenvironment of the oral cavity. Bacteria may then migrate via blood circulation and be deposited into the cerebral vessels, promoting new inflammatory responses and adjusting the vessel wall quality. Therefore, chronic periodontal inflammation by bacteria can lead to systemic inflammation that could be a cause of IA rupture because of wall changes such as apoptosis, infiltration of T cells and macrophages, and complement activation.⁴ This condition is in addition to the traditional risk factors for IA formation, including being younger than 50 years and female, hyperlipidemia, smoking, aneurysm size ratio, polycystic kidney disease, old cerebral infarct, multiple aneurysms, Marfan and collagen syndromes, systemic arterial hypertension, family history, and Japanese or Finnish ethnicity.^{2,3}

Therefore, it is necessary to perform further comparative studies to establish the exact roles and relationships of odontogenic bacteria, oral infection, and IA to understand the pathophysiology of aneurysm rupture and formation. In any case, the authors should be congratulated for continuing this interesting debate and for bringing to light these thoughtful insights into the genesis of IA.

Nicollas Nunes Rabelo, MD

Renan Salomão Rodrigues, MD

Arthur Araújo Massoud Salame, MD

Paulo Henrique Braz-Silva, DDS, PhD

Manoel Jacobsen Teixeira, MD, PhD

Eberval Gadelha Figueiredo, MD, PhD

University of São Paulo, São Paulo, Brazil

References

1. Aboukais R, Loiez C, Leclerc X, Bourgeois P, Wallet F, Menovsky T, et al: Absence of bacteria in intracranial aneurysms. *J Neurosurg* [epub ahead of print March 1, 2019. DOI: 10.3171/2018.12.JNS183044]
2. Boucher Y, Matossian L, Rilliard F, Machtou P: Radiographic evaluation of the prevalence and technical quality of root canal treatment in a French subpopulation. *Int Endod J* 35:229–238, 2002
3. Bourgeois D, Bouchard P, Mattout C: Epidemiology of periodontal status in dentate adults in France, 2002–2003. *J Periodontol Res* 42:219–227, 2007
4. Pyysalo MJ, Mishra PP, Sundström K, Lehtimäki T, Karhunen PJ, Pessi T, et al: Increased tooth brushing frequency is associated with reduced gingival pocket bacterial diversity in patients with intracranial aneurysms. *PeerJ* 7:e6316, 2019
5. Pyysalo MJ, Pyysalo LM, Pessi T, Karhunen PJ, Lehtimäki T, Oksala N, et al: Bacterial DNA findings in ruptured and unruptured intracranial aneurysms. *Acta Odontol Scand* 74:315–320, 2016

6. Pyysalo MJ, Pyysalo LM, Pessi T, Karhunen PJ, Öhman JE: The connection between ruptured cerebral aneurysms and odontogenic bacteria. *J Neurol Neurosurg Psychiatry* 84:1214–1218, 2013

Disclosures

The authors report no conflict of interest.

Correspondence

Eberval Gadelha Figueiredo: ebgedelha@yahoo.com.

INCLUDE WHEN CITING

Published online October 4, 2019; DOI: 10.3171/2019.5.JNS191267.

Response

No response was received from the authors of the original article.

©AANS 2020, except where prohibited by US copyright law

Are disproportionate subarachnoid spaces and stroke history predictors of external lumbar drainage outcome in iNPH?

TO THE EDITOR: The article by El Ahmadieh et al.¹ pursues the aim to find predictors of one of the most practiced and accurate tests to select idiopathic normal pressure hydrocephalus (iNPH) patients for shunt surgery (El Ahmadieh TY, Wu EM, Kafka B, et al: Lumbar drain trial outcomes of normal pressure hydrocephalus: a single-center experience of 254 patients. *J Neurosurg* [epub ahead of print January 4, 2019. DOI: 10.3171/2018.8.JNS181059]).^{5,7} The issue is important because, although external lumbar drainage (ELD) is becoming less demanding for patients and for the healthcare system as well, it is not risk free, and it involves discomfort and requires hospitalization with the ensuing costs,² hence the value in finding methods able to overcome the need for this invasive test. However, the study by El Ahmadieh et al.¹ presents several limitations that make their results difficult for the reader to understand and, as they stand, of moderate clinical utility.

According to the classification of iNPH,⁸ it is questionable to label the study population of El Ahmadieh et al.¹ as probable iNPH patients because of the presence in some patients of neurodegenerative diseases. Also, a history of brain surgery may be suggestive of a secondary origin for some of the patients' hydrocephalus.⁶

The reader is left unaware of the breakdown of potential predictors between the ELD responsive and unresponsive groups. This is a critical point because, for instance, it cannot be verified where the 29% of patients affected by neurodegenerative disorders fall, particularly the 25 patients with a diagnosis of Alzheimer's disease who are not expected to pass an ELD test⁹ and should have been excluded from the study.

The authors found that a negative history of stroke and disproportionate subarachnoid spaces are predictors

of successful ELD, but the reader cannot see how many patients with none, one, or both have either a positive or negative test outcome. In other words, it is impossible to derive the accuracy of the method, including its sensitivity, specificity, and predictive values. Even if the predictors were reliable, and we believe they are not for the above reasons, they are unlikely to be a real game changer in the management of iNPH, because of the weak correlation between them and the outcome ($r^2 = 0.12$).

A much more important step forward would be the development of a bedside clinical prediction tool allowing the clinician to identify a subset of iNPH patients with a very high a priori probability to have a negative or positive ELD trial. Those patients with a very high likelihood of having a negative ELD outcome could be confidently screened out and spared from undergoing both ELD and subsequent surgery. Likewise, those patients identified as having a very high a priori probability of a positive ELD outcome could be safely assigned to surgery without undergoing ELD beforehand. Eventually, this would avoid undue discomfort and complication risks associated with ELD³ and surgery⁴ for a subset of iNPH patients, with obvious savings for healthcare providers. This would really move things along.

Pasquale Gallina, MD^{1,2}

Giancarlo Lastrucci, MD¹

Saverio Caini, MD³

Berardino Porfirio, MD^{2,4}

¹Florence School of Neurosurgery, University of Florence, Italy

²Careggi University Hospital, Florence, Italy

³Cancer Risk Factors and Lifestyle Epidemiology Unit, Cancer Research and Prevention Institute (ISPRO), Florence, Italy

⁴University of Florence, Italy

References

1. El Ahmadieh TY, Wu EM, Kafka B, Caruso JP, Neeley OJ, Plitt A, et al: Lumbar drain trial outcomes of normal pressure hydrocephalus: a single-center experience of 254 patients. *J Neurosurg* [epub ahead of print January 4, 2019. DOI: 10.3171/2018.8.JNS181059]
2. Gallina P, Lastrucci G, Caini S, Di Lorenzo N, Porfirio B, Scollato A: Accuracy and safety of 1-day external lumbar drainage of CSF for shunt selection in patients with idiopathic normal pressure hydrocephalus. *J Neurosurg* [epub ahead of print November 1, 2018. DOI: 10.3171/2018.6.JNS18400]
3. Governale LS, Fein N, Logsdon J, Black PM: Techniques and complications of external lumbar drainage for normal pressure hydrocephalus. *Neurosurgery* 63 (4 Suppl 2):379–384, 2008
4. Halperin JJ, Kurlan R, Schwab JM, Cusimano MD, Gronseth G, Gloss D: Practice guideline: Idiopathic normal pressure hydrocephalus: Response to shunting and predictors of response: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology* 85:2063–2071, 2015 (Erratum in *Neurology* 86:793, 2016)
5. Marmarou A, Bergsneider M, Klinge P, Relkin N, Black PM: The value of supplemental prognostic tests for the preoperative assessment of idiopathic normal-pressure hydrocephalus. *Neurosurgery* 57 (3 Suppl):S17–S28, 2005
6. Marmarou A, Bergsneider M, Relkin N, Klinge P, Black PM: Development of guidelines for idiopathic normal-pressure hy-

drocephalus: introduction. **Neurosurgery** 57 (3 Suppl):S1–S3, 2005

7. Marmarou A, Young HF, Aygok GA, Sawauchi S, Tsuji O, Yamamoto T, et al: Diagnosis and management of idiopathic normal-pressure hydrocephalus: a prospective study in 151 patients. **J Neurosurg** 102:987–997, 2005
8. Relkin N, Marmarou A, Klinge P, Bergsneider M, Black PM: Diagnosing idiopathic normal-pressure hydrocephalus. **Neurosurgery** 57 (3 Suppl):S4–S16, 2005
9. Silverberg GD, Mayo M, Saul T, Fellmann J, Carvalho J, McGuire D: Continuous CSF drainage in AD: results of a double-blind, randomized, placebo-controlled study. **Neurology** 71:202–209, 2008

Disclosures

The authors report no conflict of interest.

Correspondence

Berardino Porfirio: berardino.porfirio@unifi.it.

INCLUDE WHEN CITING

Published online June 28, 2019; DOI: 10.3171/2019.2.JNS19268.

Response

We thank Dr. Porfirio and colleagues from Florence, Italy, for their interest and thorough review of our article, which brings up several important points and encourages further discussion of the topic. We also value their experience and previous contributions to the literature.^{2,3} The main goal of this study was to review our institutional experience with lumbar drainage (LD) in iNPH patients and assess outcomes and complication rates. We had a special interest in looking at predictors of passing an LD trial in our patient population and comparing them with predictors reported in the literature. It was not our intention to create a universal predictive model that would enable physicians to overcome the need for an LD trial. While such a model would provide an invaluable tool that could help avoid unnecessary invasive testing, the current body of literature supports a trial of temporary CSF diversion to confirm appropriate response prior to permanent shunt placement.¹ A reliable predictive model may be achieved in the future by conducting well-designed, multicenter prospective studies, but our study represents a retrospective review of more than 10 years of experience at our center.

Regarding the question about the legitimacy of the “probable” iNPH diagnosis in our patient population, the iNPH guidelines clearly state that there should be “no other neurological, psychiatric, or general medical conditions that are sufficient to explain the presenting symptoms.”²⁴ Therefore, the mere presence of a neurodegenerative disease, which is to be expected in a patient population with a mean age of 77 years, or a history of brain surgery, does not exclude patients from a probable iNPH diagnosis if their symptoms are not explained by these conditions. These patients are typically referred to the neurosurgery clinic for iNPH evaluation after an extensive workup, performed by their neurologist, that exhausts all other possibilities as a potential cause of their presenting symptoms. We believe that our study population represents a real-life iNPH patient population that carries an inherent challenge of harboring other comorbidities that could be contribut-

ing to the clinical picture. In our experience, the perfect iNPH patient who has no other medical comorbidities and no overlapping symptoms is a rare phenomenon.

We agree with the authors that patients with Alzheimer’s disease are not ideal candidates for an LD trial. Their chances of passing an LD trial and benefiting from a subsequent shunt are low overall but not zero, based on our experience. In our study, 25 patients had a working diagnosis of Alzheimer’s disease in addition to progressive symptoms and new imaging findings suggestive of iNPH that were deemed worthy of investigation by the neurologist. Of these 25 patients, 4 passed the LD trial, and 3 of them eventually underwent shunt placement and experienced an improvement in their symptoms. One patient elected not to proceed with a shunt procedure. The outcome of shunting in our patient population required a separate analysis and is currently under review in another medical journal. With these numbers, 16% (4/25) of patients with a clinical diagnosis of Alzheimer’s disease in this study passed the LD trial. Therefore, excluding these patients, as suggested by the authors, would have eliminated the chance of picking up those few patients who could potentially benefit from a shunt. Whether that is worth the risk of an LD trial remains a judgment call to be made by the physician and the patient or family after a thorough discussion of expectations and after weighing the risks of the procedure against the potential benefits. Of course, it would be of interest to further study patients with various other clinical diagnoses who present with symptoms of iNPH, though sample sizes would need to be adequate for meaningful analyses.

The development of a bedside predictive model that allows physicians to screen out patients who are unlikely to benefit from CSF diversion is a challenging task with the currently available predictors and levels of evidence. Until that can be achieved, the LD trial remains a useful, reliable, and relatively safe tool that can predict shunt responsiveness in patients with iNPH. Neurodegenerative diseases are not an absolute contraindication for an LD trial, and select patients with findings highly suggestive of iNPH should be given the chance to pass or fail an LD trial. A case-by-case approach to such patients and extensive patient and family counseling are encouraged.

Tarek Y. El Ahmadieh, MD

Salah G. Aoun, MD

Eva M. Wu, MD

DaiWai M. Olson, PhD, RN

C. Munro Cullum, PhD

H. Hunt Batjer, MD

Jonathan A. White, MD

The University of Texas Southwestern Medical Center, Dallas, TX

References

1. Chotai S, Medel R, Herial NA, Medhkour A: External lumbar drain: a pragmatic test for prediction of shunt outcomes in idiopathic normal pressure hydrocephalus. **Surg Neurol Int** 5:12, 2014
2. Gallina P, Lastrucci G, Caini S, Di Lorenzo N, Porfirio B, Scollato A: Accuracy and safety of 1-day external lumbar drainage of CSF for shunt selection in patients with idiopath-

- ic normal pressure hydrocephalus. **J Neurosurg** [epub ahead of print November 16, 2018; DOI: 10.3171/2018.6.JNS18400]
3. Gallina P, Savastano A, Becattini E, Orlandini S, Scollato A, Rizzo S, et al: Glaucoma in patients with shunt-treated normal pressure hydrocephalus. **J Neurosurg** 129:1078–1084, 2018
 4. Relkin N, Marmarou A, Klinge P, Bergsneider M, Black PM: Diagnosing idiopathic normal-pressure hydrocephalus. **Neurosurgery** 57 (3 Suppl):S4–S16, 2005

INCLUDE WHEN CITING

Published online June 28, 2019; DOI: 10.3171/2019.2.JNS19293.

©AANS 2020, except where prohibited by US copyright law

The future of patient safety in neurological surgery

TO THE EDITOR: We read with great interest the article by Sankaran et al.¹ on the current and future directions of patient safety in neurological surgery (Sankaran S, Andrews JP, Chicas M, et al. Patient safety movement in neurological surgery: the current state and future directions. *J Neurosurg*. 2020;132(1):313–323). The paper highlights the promise of technological advancements such as the adoption of electronic medical records, neurosurgical simulators, and artificial intelligence (AI) concerning patient safety. While we agree that these developments have merit, we feel that the adoption of this technology in medicine may pose a threat to patient safety. Moreover, the seemingly simple issue of work-hour restrictions helped us commend the advantages but also the limitations that evidence-based medicine can offer.

Artificial intelligence and its various medical applications need to be examined for inconsistencies. Currently, the application of AI to medicine is limited by its sample pool. Thus, machine learning algorithms can only be applied on patients of a certain age group, ethnicities, and anatomical variations. If the same neural network is applied in a different hospital or minority population, the analysis and prediction of the AI would be completely biased. Therefore, we call on the scientific community to develop robust evaluation methods *prior* to the implementation of new technology² and encourage new information-sharing systems to be realized while protecting patient anonymity. With recent information breaches such as the 2,200,000 personal medical records hacked from *21st Century Oncology*,³ we realize that the adoption of new technology raises concerns of privacy, data usage policy, security, oversight, and the lack of traditional medical relationships. These factors have substantial potential to lead to worse patient outcomes in the medical, social, and economic sectors. Therefore, along with the current limitations of AI, we are cautious to paint an optimistic view of the future that embraces technology.

The paper indicates that resident work-hour restrictions may have actually harmed patient safety rather than improved it. In this case, evidence-based medicine eloquently provided clarity to a seemingly obvious assumption that working fewer hours results in better patient safety as the

physician is more alert and rested. However, we feel that a healthy distance is required from evidence-based medicine as it might also hinder medical innovation. When William T. G. Morton used ether to provide general anesthesia to his dental patients in 1846, he had meager evidence (mostly from experiments on his dog and himself) to support ether's anesthetic properties.⁴ Furthermore, plastic surgery, neurosurgery, and arterial ligation techniques were all pioneered during the American Civil War, a time when physicians enjoyed considerable leeway to test new ideas that did not necessarily have concrete evidence.⁵ Therefore, while we herald medicine's current devotion to evidence-based medicine, we ask that physicians are allowed sufficient latitude to experiment to pave safer procedures for patients.

Both the physician and the patient share a future held in optimism; thus, we commend Sujatha Sankaran and colleagues for raising these pertinent issues of patient safety in their paper. We hope this letter prompts physicians to keep cognizant of the current issues in medicine so that our field will enjoy a more fruitful future.

Tomasz Szmuda, MD, PhD

Shan Ali

Paweł Słoniewski, MD, PhD

Medical University of Gdańsk, Poland

References

1. Sankaran S, Andrews JP, Chicas M, et al. Patient safety movement in neurological surgery: the current state and future directions. *J Neurosurg*. 2020;132(1):313–323.
2. Hamet P, Tremblay J. Artificial intelligence in medicine. *Metabolism*. 2017;69S:S36–S40.
3. Walter K. 21st Century Oncology notifies 2.2 million of hacking, data breach. WPEC. Accessed March 22, 2020. <https://cbs12.com/news/local/21st-century-oncology-notifies-22-million-of-hacking-data-breach>
4. Robinson DH, Toledo AH. Historical development of modern anesthesia. *J Invest Surg*. 2012;25:141–149.
5. Reilly RF. Medical and surgical care during the American Civil War, 1861–1865. *Baylor Univ Med Cent Proc*. 2016;29:138–142.

Disclosures

The authors report no conflict of interest.

Correspondence

Shan Ali: shanali@gumed.edu.pl.

INCLUDE WHEN CITING

Published online April 10, 2020; DOI: 10.3171/2019.11.JNS193005.

Response

We greatly appreciate the perspective of Szmuda et al. regarding our paper on the current and future directions of patient safety in neurosurgery. We agree that AI offers a tremendous opportunity for improving the quality, safety, and efficiency of care but that there is a risk of overhype and unanticipated consequences. It will be important to subject AI-based interventions to rigorous evaluation. We also agree with the notion that evidence-based medicine has limitations and may hinder innovation if applied indiscriminately to areas that do not lend themselves to

simple evidence-based analysis. We suggest that more research and a wider focus on improving patient safety in neurosurgery will benefit both practitioners and patients immensely.

Sujatha Sankaran, MD
John P. Andrews, MD
Madeline Chicas, MHA
Robert M. Wachter, MD
Mitchel S. Berger, MD

University of California, San Francisco, CA

INCLUDE WHEN CITING

Published online April 10, 2020; DOI: 10.3171/2020.3.JNS193031.

©AANS 2020, except where prohibited by US copyright law

Meralgia paresthetica: what to do?

TO THE EDITOR: I read the interesting article by Malessy et al. describing the technique and results of dynamic decompression of the lateral femoral cutaneous nerve (LFCN) to treat meralgia paresthetica (MP) (Malessy MJA, Eekhof J, Pondaag W. Dynamic decompression of the lateral femoral cutaneous nerve to treat meralgia paresthetica: technique and results. *J Neurosurg.* 2019;131(5):1552–1560).¹ The article describes a layer of fascia superficial to the LFCN and another layer deeper to the nerve. These were previously clearly described as the LFCN canal by our group and independently confirmed by Xu et al.^{2,3} The LFCN canal is an anatomical fact that cannot be ignored and should be the basis of any surgical approach. It clearly defines the planes that need to be decompressed in surgery on the LFCN without the need for dynamic testing. In addition, many patients with MP have the nerve in proximity to or riding over the iliac crest at the region of the anterior superior iliac spine (ASIS). The technique described by Malessy et al. does not address this patient population since it does not tell us how to deal with the bone. The dynamic decompression as described raises some concerns. Most MP patients have a high body mass index, so taking that heavy lower limb and moving it in flexion, abduction, and extension seems impractical and not reproducible and presents the risks of breaking sterility and inflicting injury to the person while lifting up the leg. The LFCN canal has well-defined anterior and posterior walls that must be decompressed regardless of the findings with limb mobilization.² The LFCN transposition as described by Hanna addresses the issue of proximity to bone by mobilizing the LFCN medially for about 2 cm, which will eliminate the nerve rubbing against the bone.^{4–6} This situation was not observed in the authors' series either because of ethnic differences or more likely due to a lack of intraoperative measurements prior to decompression. While proximity of the LFCN to the ASIS is rare in human cadavers, it is very common in MP patients. This means it is part of the pathology and needs to be addressed by transposition. It is likely that Malessy and colleagues actually performed transpositions, but it is hard to confirm

since no measurements were taken. This is very obvious in their Fig. 1A, where the nerve is very close to the ASIS (labeled 2) prior to deep decompression, then is clearly medialized after deep decompression in Fig. 1C.¹

Amgad Hanna, MD

University of Wisconsin, Madison, WI

References

1. Malessy MJA, Eekhof J, Pondaag W. Dynamic decompression of the lateral femoral cutaneous nerve to treat meralgia paresthetica: technique and results. *J Neurosurg.* 2019;131(5):1552–1560.
2. Hanna A. The lateral femoral cutaneous nerve canal. *J Neurosurg.* 2017;126:972–978.
3. Xu Z, Tu L, Zheng Y, et al. Fine architecture of the fascial planes around the lateral femoral cutaneous nerve at its pelvic exit: an epoxy sheet plastination and confocal microscopy study. *J Neurosurg.* 2019;131(6):1860–1868.
4. Hanna A. Transposition of the lateral femoral cutaneous nerve. *J Neurosurg.* 2019;130(2):496–501.
5. Hanna A, Hanna B. Response To: Anatomical considerations on transposition of the lateral femoral cutaneous nerve. *Clin Anat.* 2018;31(8):1222–1224.
6. Hanna AS. Lateral femoral cutaneous nerve transposition: renaissance of an old concept in the light of new anatomy. *Clin Anat.* 2017;30(3):409–412.

Disclosures

The author reports no conflict of interest.

Correspondence

Amgad Hanna: ah2904@yahoo.com.

INCLUDE WHEN CITING

Published online March 20, 2020; DOI: 10.3171/2019.10.JNS192708.

Response

We have read with interest the letter from Dr. Hanna. Since our publication in 2018, in which we reported the outcome of dynamic decompression of the LFCN in idiopathic MP, the total number of surgically treated patients with sufficient follow-up has amounted to 30. In the vast majority of patients, the pain completely disappeared and skin sensation recovered to (near) normal. As described, we first decompress the LFCN with the leg in a neutral position. Subsequently, we move the leg in extension and abduction. In doing so, we observed that fibers located dorsal to the LFCN tighten and cause compression at various locations. These could be fibers of the musculotendinous origin of the sartorius muscle at the level of the ASIS, the iliac fascia proximal to the ASIS, or the lateral aspect of the fascia of the sartorius muscle distal to the ASIS. Additionally cutting these fibers, which we call “dynamic decompression,” is likely the distinctive factor that positively distinguishes our results from those of others who perform the same decompression but only with the leg in a neutral position.^{1,2} The good results following dynamic decompression encourages us to continue using this technique.

Dr. Hanna does not move the leg out of fear of infection and physical strain and therefore misses the observation of the dynamics of compression of the LFCN. This is unfortunate because this relatively easy and simple procedure

takes only a few minutes to perform, is reproducible, and does not break sterility. Additionally, it is not a burdensome procedure regardless of body mass index. More importantly, moving the leg provides important information.

In our consecutive series, the entrapment site of the LFCN was consistently found medial to the ASIS. By observing the LFCN during limb movement at the end stage of dynamic decompression, it becomes clear that “riding” over the ASIS does not occur, nor did we see “rubbing against the bone.” Complete dorsal decompression and release induce a posterior shifting of the LFCN. We did not perform medial transposition as was first described by Keegan and Holyoke in the *Journal of Neurosurgery*³ and now proposed again by Dr. Hanna.⁴ In view of our observations and results, there is no need for medial transposition. Fixation of the LFCN in this position with a stitch is, in view of the dynamics, in fact contraindicated.

Martijn J. A. Malessy, MD, PhD^{1,2}

Job Eekhof, MD, PhD²

Willem Pondaag, MD, PhD^{1,2}

¹Leiden University Medical Center, Leiden, The Netherlands

²Alrijne Medical Center, Leiden, The Netherlands

References

1. Siu TL, Chandran KN. Neurolysis for meralgia paresthetica: an operative series of 45 cases. *Surg Neurol*. 2005;63(1):19–25.
2. Son BC, Kim DR, Kim IS, et al. Neurolysis for meralgia paresthetica. *J Korean Neurosurg Soc*. 2012;51(6):363–366.
3. Keegan JJ, Holyoke EA. Meralgia paresthetica. An anatomical and surgical study. *J Neurosurg*. 1962;19(4):341–345.
4. Hanna A. Transposition of the lateral femoral cutaneous nerve. *J Neurosurg*. 2019;130(2):496–501.

INCLUDE WHEN CITING

Published online March 20, 2020; DOI: 10.3171/2020.2.JNS192753.

©AANS 2020, except where prohibited by US copyright law