Neurosurgical Forum LETTERS TO THE EDITOR

The third ventricle bowing and ETV success

TO THE EDITOR: We read with great interest the article by Gürbüz et al.¹ (Gürbüz MS, Dağçınar A, Bayri Y, et al. Anterior third ventricular height and infundibulochiasmatic angle: two novel measurements to predict clinical success of endoscopic third ventriculostomy in the early postoperative period. *J Neurosurg.* 2020;132[6]:1764–1772). Bowing of the third ventricle has been increasingly recognized as a way to predict endoscopic third ventriculostomy (ETV) success preoperatively.^{2,3} Using the signs of anterior and inferior bowing of the third ventricle, Gürbüz et al. used the two novel parameters of anterior third ventricular height (TVH) and infundibulochiasmatic (IC) angle to predict ETV success in both pre- and postoperative stages. We congratulate the authors on their excellent work, which proved that the change in morphology of the third ventricle can predict ETV success. In our department, we have used the sign of bowing of the third ventricle floor (TVFB) to preoperatively select ETV candidates since 2013. The ETV success rate in our series was 81%, which also proved that TVFB was an efficient method to select ETV candidates.⁴

However, although third ventricle bowing is useful to predict ETV success in both the pre- and postoperative period, its limitations should be remembered. In their studies, Foroughi et al.³ and Dlouhy et al.² found that 46% and 33% of hydrocephalic patients without bowing of the third ventricle could still be successfully treated with ETV. Hence, it seems that in some patients, even without bowing of the third ventricle, ETV can be successful and changes in the anterior TVH and IC angle will not be present. Third ventricle bowing has been anecdotally suggested as predictive of more intraventricularly obstructive types of hydrocephalus, which can achieve higher success rates with ETV;^{2,5} however, when the obstructions appear in the cisterns, such as the basal cistern and supratentorial cisterns, bowing of the third ventricle may not be present. Rekate explored the sites of obstruction to CSF circulation that caused hydrocephalus and found that these sites included not only the aqueduct and the outlet of the fourth ventricle, but also the basal cisterns blocking the CSF pathway.^{6,7} In some types of communicating hydrocephalus caused by trauma, meningitis, and hemorrhage, there is always some extent of adhesive obstruction in the basal

cisterns. And while bowing of the third ventricle may not be present in these patients with extraventricular intracisternal obstruction, ETV remains effective by creating an internal bypass.^{8,9} Furthermore, Singh et al. found that in some cases of hydrocephalus, even when MR ventriculography depicted CSF patent from the fourth ventricle into the cisternal spaces without intraventricular obstructions, ETV can still be successful (70%).⁹ Therefore, for hydrocephalic patients with third ventricle bowing preoperatively, the methods by Gürbüz et al. were very useful to predict ETV success. But for hydrocephalic patients without preoperative third ventricle bowing, other complementary preoperative prediction systems, such as the ETV success score,¹⁰ may be required to select ETV candidates. Other supplementary methods such as observing signs of periventricular infiltration of CSF, such as tension of the ventricles, fissures, sulci, and gyri of the brain, may help to determine ETV success in the early postoperative period in these patients.

Qiguang Wang, MD
Si Zhang, PhD
Yan Ju, PhD

West China Hospital of Sichuan University, Chengdu, Sichuan, China

References

1. Gürbüz MS, Dağçınar A, Bayri Y, et al. Anterior third ventricular height and infundibulochiasmatic angle: two novel measurements to predict clinical success of endoscopic third ventriculostomy in the early postoperative period. *J Neurosurg.* 2020;132(6):1764–1772.
2. Dlouhy BJ, Capuano AW, Madhavan K, et al. Preoperative third ventricular bowing as a predictor of endoscopic third ventriculostomy success. *J Neurosurg Pediatr.* 2012;9(2):182–190.
3. Foroughi M, Wong A, Steinbok P, et al. Third ventricular shape: a predictor of endoscopic third ventriculostomy success in pediatric patients. *J Neurosurg Pediatr.* 2011;7(4):389–396.
4. Wang Q, Cheng J, Zhang S, et al. Prediction of endoscopic third ventriculostomy (ETV) success with preoperative third ventricle floor bowing (TVFB): a supplement to ETV success score. *Neurosurg Rev.* Published online November 6, 2020. doi:10.1007/s10143-019-01178-1
5. Kehler U, Regelsberger J, Gliemroth J, Westphal M. Outcome prediction of third ventriculostomy: a proposed hydrocephalus grading system. *Minim Invasive Neurosurg.* 2006;49(4):238–243.
6. Rekate HL. Comments on the article by D. Greitz “Paradigm

shift in hydrocephalus research in legacy of Dandy's pioneering work: rationale for third ventriculostomy in communicating hydrocephalus." *Childs Nerv Syst.* 2007;23(11):1227–1231.

7. Rekate HL. Circuit diagram of the circulation of cerebrospinal fluid. 1989. *Pediatr Neurosurg.* 1994;21(4):248–253.
8. Hailong F, Guangfu H, Haibin T, et al. Endoscopic third ventriculostomy in the management of communicating hydrocephalus: a preliminary study. *J Neurosurg.* 2008;109(5):923–930.
9. Singh I, Haris M, Husain M, et al. Role of endoscopic third ventriculostomy in patients with communicating hydrocephalus: an evaluation by MR ventriculography. *Neurosurg Rev.* 2008;31(3):319–325.
10. Labidi M, Lavoie P, Lapointe G, et al. Predicting success of endoscopic third ventriculostomy: validation of the ETV Success Score in a mixed population of adult and pediatric patients. *J Neurosurg.* 2015;123(6):1447–1455.

Disclosures

The authors report no conflict of interest.

Correspondence

Yan Ju: juyan0806@126.com.

INCLUDE WHEN CITING

Published online October 16, 2020; DOI: 10.3171/2020.5.JNS202016.

Response

We thank the authors for their interest in and comments on our study. As emphasized in their comments, in the study by Foroughi et al., in which the authors assessed the convexity of the lamina terminalis and convexity of the floor of the third ventricle, ETV success was found in 46% of patients without both features.¹ We should note that “without both features” does not at all mean “no change in the third ventricle shape.” Although every area of the third ventricle goes through some changes, the recognizable ones occur mainly in three directions: anterior-superior wall (lamina terminalis), inferior wall (tuber cinereum), and anterior-inferior wall (infundibular recess and optic recess). In reviewing the literature, we found that the changes in all three of these directions have not been included in any study but ours. We endeavored to cover the changes happening in these three directions. Seventy procedures (89.7%) in our study were successful, revealing that our success rate seems relatively higher. It is possibly due to the strict criteria for selecting the study population, which consisted of adult patients with only intraventricular obstructive hydrocephalus. As stated in the literature,^{2–4} patients with intraventricular obstructive hydrocephalus can obtain higher success rates with ETV.

In their valuable comments, Wang and colleagues⁴ stressed that ETV still helps in some patients with intracisternal extraventricular obstruction (“some types of communicating hydrocephalus caused by trauma, meningitis, and hemorrhage”) without prominent changes in the third ventricle walls. Dlouhy et al. found that the patients with third ventricular bowing had a threefold higher chance of successful ETV, although 33% of the patients without bowing still had successful ETV.² In such cases,

ETV surely may help, although its success rate decreases.⁵ We accept that observing no visible change in the shape of the anterior portion of the third ventricle does not absolutely mean that the procedure will fail. However, in a structurally normal ventricular system with normal CSF production, it is not possible to see normal intraventricular pressure when there is an obstruction. In obstructive hydrocephalus, configurational changes will necessarily happen as a result of high pressure. As an exception, high pressure inside the third ventricle with no configurational changes in its walls may only be caused by a loss of elasticity (resultant stiffness) in the third ventricle walls following an infection, trauma, or hemorrhage.^{4–6}

As a general rule, the ETV success score and other methods such as observing signs of periventricular infiltration of CSF, such as tension of the ventricles, fissures, sulci, and gyri of the brain, may help to predict ETV success.⁷ Nevertheless, configurational changes, including the ones described in our study, help keep the success rates higher in ETV, particularly in obstructive hydrocephalus cases. Further studies are needed to figure out the negative and positive predictive values of the mentioned configurational features.

Mehmet Sabri Gürbüz, MD

Istanbul Medeniyet University, Medical School, Istanbul, Turkey

Aşkın Şeker, MD

Altınbaş University, Medical School, Istanbul, Turkey

References

1. Foroughi M, Wong A, Steinbok P, et al. Third ventricular shape: a predictor of endoscopic third ventriculostomy success in pediatric patients. *J Neurosurg Pediatr.* 2011;7(4):389–396.
2. Dlouhy BJ, Capuano AW, Madhavan K, et al. Preoperative third ventricular bowing as a predictor of endoscopic third ventriculostomy success. *J Neurosurg Pediatr.* 2012;9(2):182–190.
3. Kehler U, Regelsberger J, Gliemroth J, Westphal M. Outcome prediction of third ventriculostomy: a proposed hydrocephalus grading system. *Minim Invasive Neurosurg.* 2006;49(4):238–243.
4. Wang Q, Cheng J, Zhang S, et al. Prediction of endoscopic third ventriculostomy (ETV) success with preoperative third ventricle floor bowing (TVFB): a supplement to ETV success score. *Neurosurg Rev.* Published online November 6, 2019. doi:10.1007/s10143-019-01178-1
5. Singh I, Haris M, Husain M, et al. Role of endoscopic third ventriculostomy in patients with communicating hydrocephalus: an evaluation by MR ventriculography. *Neurosurg Rev.* 2008;31(3):319–325.
6. Rekate HL, Nadkarni TD, Wallace D. The importance of the cortical subarachnoid space in understanding hydrocephalus. *J Neurosurg Pediatr.* 2008;2(1):1–11.
7. Labidi M, Lavoie P, Lapointe G, et al. Predicting success of endoscopic third ventriculostomy: validation of the ETV Success Score in a mixed population of adult and pediatric patients. *J Neurosurg.* 2015;123(6):1447–1455.

INCLUDE WHEN CITING

Published online October 16, 2020; DOI: 10.3171/2020.8.JNS202042.

©AANS 2021, except where prohibited by US copyright law

Is time-weighted PaO₂ during the hyperacute phase of aneurysmal subarachnoid hemorrhage really helpful in usual care settings?

TO THE EDITOR: We have read the article by Fukuda et al.¹ with great interest (Fukuda S, Koga Y, Fujita M, et al. Hyperoxemia during the hyperacute phase of aneurysmal subarachnoid hemorrhage is associated with delayed cerebral ischemia and poor outcome: a retrospective observational study [published online November 15, 2019]. *J Neurosurg*. doi:10.3171/2019.9.JNS19781). Previously, we have examined the association between respiratory management and neurological outcome in patients with subarachnoid hemorrhage.^{2,3} The authors of the present study showed that the time-weighted average PaO₂ during the first 24 hours (TWA_{24h}-PaO₂) was associated with delayed cerebral ischemia (DCI) and poor neurological outcomes. TWA_{24h}-PaO₂ is a marker of the degree and duration of hyperoxemia, and there should be increased attention on TWA_{24h}-PaO₂ during the early brain injury phase.⁴ However, we have two questions about this study.

First, we believe that there is a problem with the covariate selection for multivariable logistic regression analyses. The authors selected covariates using *p* values from the univariate analysis and stepwise procedure. Consequently, the Hunt-Hess grade was excluded from their multivariable logistic regression analyses. The Hunt-Hess or Hunt-Kosnik grade is a clinically important factor in patients with subarachnoid hemorrhage. It helps physicians determine treatment strategies⁵ and estimate neurological outcome; therefore, it has been included as a covariate to adjust multivariable logistic regression analysis in previous studies.^{2,6}

Second, it seems difficult for physicians to integrate the concept of TWA_{24h}-PaO₂ into clinical practice. During the hyperacute phase of aneurysmal subarachnoid hemorrhage, physicians have many treatment priorities, including blood pressure management to prevent rebleeding and maintain cerebral perfusion, analgesedative optimization, and preparation for aneurysmal treatment. If a real-time calculation system for TWA_{24h}-PaO₂ were available, it would be possible to manage this factor. We believe that the appropriate management of PaO₂ (or SpO₂) would lead to the appropriate management of TWA_{24h}-PaO₂ in usual care settings. Therefore, we request that the authors provide the cutoff values of TWA_{24h}-PaO₂ between the non-DCI and DCI groups or the favorable and poor outcome groups. We hypothesize that maintaining PaO₂ under these cutoff levels can result in the maintenance of TWA_{24h}-PaO₂ within a safe range.

We look forward to the authors' reply.

Tomoya Okazaki, MD
Shota Yokoyama, MD
Kenya Kawakita, MD, PhD
Yasuhiro Kuroda, MD, PhD
Kagawa University Hospital, Kagawa, Japan

References

1. Fukuda S, Koga Y, Fujita M, et al. Hyperoxemia during the hyperacute phase of aneurysmal subarachnoid hemorrhage is associated with delayed cerebral ischemia and poor outcome: a retrospective observational study [published online November 15, 2019]. *J Neurosurg*. doi:10.3171/2019.9.JNS19781
2. Yokoyama S, Hifumi T, Kawakita K, et al. Early hyperoxia in the intensive care unit is significantly associated with unfavorable neurological outcomes in patients with mild-to-moderate aneurysmal subarachnoid hemorrhage. *Shock*. 2019;51:593–598.
3. Yokoyama S, Hifumi T, Okazaki T, et al. Association of abnormal carbon dioxide levels with poor neurological outcomes in aneurysmal subarachnoid hemorrhage: a retrospective observational study. *J Intensive Care*. 2018;6:83.
4. Okazaki T, Kuroda Y. Aneurysmal subarachnoid hemorrhage: intensive care for improving neurological outcome. *J Intensive Care*. 2018;6:28.
5. de Oliveira Manoel AL, Goffi A, Marotta TR, et al. The critical care management of poor-grade subarachnoid haemorrhage. *Crit Care*. 2016;20:21.
6. Jeon SB, Choi HA, Badjatia N, et al. Hyperoxia may be related to delayed cerebral ischemia and poor outcome after subarachnoid haemorrhage. *J Neurol Neurosurg Psychiatry*. 2014;85:1301–1307.

Disclosures

The authors report no conflict of interest.

Correspondence

Tomoya Okazaki: tomoyaokazaki4028@gmail.com.

INCLUDE WHEN CITING

Published online April 10, 2020; DOI: 10.3171/2020.1.JNS20177.

Response

We appreciate Dr. Okazaki's comments regarding our study. The issue of the covariate selection for the multivariate analysis was discussed among the authors. Because this study was conducted retrospectively, we wanted to avoid selecting variables intentionally for the multivariate analysis. Therefore, although it may not have been the optimal method, we selected variables with *p* values < 0.1 to enter into the stepwise selection procedure. Yokoyama et al.¹ reported that the Hunt-Hess grade is an important indicator of both clinical outcome and the adverse effect of hyperoxemia on clinical outcome in patients with SAH. The Hunt-Hess grade was excluded from the multivariate analysis in our study either by the stepwise procedure or because the *p* value was < 0.1 in the univariate analysis. Therefore, we reanalyzed the data to include the Hunt-Hess grade in the multivariate analysis. However, it did not change the result, and the TWA_{24h}-PaO₂ was consistently a significant risk factor for DCI (OR 1.08, 95% CI 1.00–1.16, *p* = 0.046) and poor neurological outcome (OR 1.15, 95% CI 1.05–1.26, *p* = 0.003), whereas the Hunt-Hess grade was not significantly associated with DCI (*p* = 0.75) or poor outcome (*p* = 0.16).

Dr. Okazaki also raised the issue of the applicability of TWA-PaO₂ to clinical practice. Although TWA-PaO₂ might not be applicable in a clinical setting, we wanted to investigate the effect of sustained hyperoxemia on patients with subarachnoid hemorrhage. Moreover, we thought that TWA-PaO₂ would provide a more reliable result than

PaO₂ measured at a single time point. In fact, we found that TWA-PaO₂ was more significantly related to DCI and poor outcome than either maximum PaO₂ or initial PaO₂.

We agree that maintaining PaO₂ below the appropriate cutoff level would allow the effective management of TWA_{24h}-PaO₂. We also examined the cutoff values for TWA_{24h}-PaO₂ between the DCI and non-DCI groups and between the favorable and poor outcome groups. We entered several cutoff values for TWA_{24h}-PaO₂ into the stepwise selection procedure for the multivariate logistic regression analyses to explore the cutoff values that were most significantly associated with DCI or poor neurological outcome. The cutoff value that led TWA_{24h}-PaO₂ to be most significantly associated with DCI or a poor neurological outcome was 190 or 160 mm Hg, respectively. However, because TWA_{24h}-PaO₂ can be affected by both hyperoxemia and hypoxemia, PaO₂ under these cutoff values would not necessarily represent the safety range in the clinical setting. Clinicians should also consider that the optimal cutoff value for PaO₂ might vary among individuals (e.g., with a history of smoking or chronic pulmonary disease, baseline cardiac function, and the severity of subarachnoid hemorrhage).

Shinya Fukuda, MD¹

Yasutaka Koga, MD, PhD¹

Motoki Fujita, MD, PhD²

Eiichi Suehiro, MD, PhD^{1,2}

Kotaro Kaneda, MD, PhD¹

Yasutaka Oda, MD, PhD²

Hideyuki Ishihara, MD, PhD²

Michiyasu Suzuki, MD, PhD²

Ryosuke Tsuruta, MD, PhD^{1,2}

¹Yamaguchi University Hospital, Ube, Yamaguchi, Japan

²Yamaguchi University Graduate School of Medicine, Ube, Yamaguchi, Japan

References

1. Yokoyama S, Hifumi T, Kawakita K, et al. Early hyperoxia in the intensive care unit is significantly associated with unfavorable neurological outcomes in patients with mild-to-moderate aneurysmal subarachnoid hemorrhage. *Shock*. 2019;51:593–598.

INCLUDE WHEN CITING

Published online April 10, 2020; DOI: 10.3171/2020.2.JNS20202.

©AANS 2021, except where prohibited by US copyright law