Neurosurgical Forum

LETTERS TO THE EDITOR

Pain outcomes for trigeminal neuralgia

TO THE EDITOR: We read with great interest the multicenter retrospective study by Mureb et al.¹ on radiosurgery for type 1 trigeminal neuralgia (TN) published in the *Journal of Neurosurgery* in July 2020 (Mureb M, Golub D, Benjamin C, et al. Earlier radiosurgery leads to better pain relief and less medication usage for trigeminal neuralgia patients: an international multicenter study. *J Neurosurg*. Published online July 3, 2020. doi:10.3171/2020.4.JNS192780). In this study the authors concentrated on analyzing pain outcomes, focusing on the time interval between the diagnosis of TN and radiosurgery.

It is well known that in TN unresponsive to pain relief medications, radiosurgical treatment results in a high initial success rate, with 80%–90% pain relief at 90 days.^{2,3} The authors showed that patients who received radiosurgery within 4 years of TN diagnosis achieved significantly faster pain relief and higher pain relief rates at 90 days than those who received radiosurgery after 4 years; more specifically, pain relief occurred at a median of 21 days versus 30 days after treatment, and pain relief at 90 days was recorded in 83.8% versus 73.7% of cases.

On the contrary, the frequent lack of success in obtaining durable pain relief is probably one of the main weaknesses of this treatment. Why the rate of pain recurrence is so high is still unclear. At 5 years, the following rates of maintained pain relief were reported: Regis et al., 64.9%;² Kondziolka et al., 46%.⁴ The authors investigated another detail of this phenomenon, that earlier radiosurgical treatment (within 4 years from TN diagnosis), as well as a higher maximum dose, is predictive of patients being pain free at treatment follow-up. This finding confirms the results already published by Lee et al.⁵ and Mousavi et al.⁶ in case series with smaller cohorts of patients.

Recently, several studies focused on brain MRI in patients affected by TN. Both gray and white matter changes over time were analyzed. Obermann et al.⁷ and Li et al.⁸ demonstrated some changes in gray matter volume and cortical thickness in several brain regions (primary and secondary somatosensory areas, insula, cingulate, hippocampus, amygdala, and thalamus) involved in pain perception and modulation in patients with chronic TN. Furthermore, DeSouza et al.⁹ conducted white matter analysis

and reported some abnormalities in both the trigeminal nerves and the central nervous system. These investigators reported abnormalities in the areas involved in the sensory and cognitive-affective dimensions of pain, attention, and motor function, including the corpus callosum, cingulum, posterior corona radiata, and superior longitudinal fasciculus.

The authors suggested that the duration of the TN may be a critical factor associated with these brain alterations.^{7,9}

It could be hypothesized that neuromodulation from radiosurgical treatment differs in patients with a more recent TN diagnosis compared with patients with an older diagnosis, possibly because of the aforementioned cerebral modifications that occur in patients with long-lasting TN. Further studies should therefore be conducted in this field.

We congratulate the authors for their interesting paper.

Overall, we agree with the authors that radiosurgical treatment should be recommended sooner rather than later in patients with type 1 medically refractory TN to improve treatment effectiveness and to avoid the need for increases in the numbers and/or dose frequencies of drugs (and the resulting well-known negative side effects) to achieve adequate pain relief.

Luigi Albano, MD

Lina Raffaella Barzaghi, MD

Pietro Mortini, MD

IRCCS San Raffaele Scientific Institute and Vita-Salute University, Milan, Italy

References

1. Mureb M, Golub D, Benjamin C, et al. Earlier radiosurgery leads to better pain relief and less medication usage for trigeminal neuralgia patients: an international multicenter study. *J Neurosurg*. Published online July 3, 2020. doi:10.3171/2020.4.JNS192780
2. Regis J, Tulasca C, Resseguier N, et al. The very long-term outcome of radiosurgery for classical trigeminal neuralgia. *Stereotact Funct Neurosurg*. 2016;94(1):24–32.
3. Tulasca C, Regis J, Sahgal A, et al. Stereotactic radiosurgery for trigeminal neuralgia: a systematic review. *J Neurosurg*. 2018;130(3):733–757.
4. Kondziolka D, Zorro O, Lobato-Polo J, et al. Gamma Knife stereotactic radiosurgery for idiopathic trigeminal neuralgia. *J Neurosurg*. 2010;112(4):758–765.
5. Lee JY, Sandhu S, Miller D, et al. Higher dose rate Gamma Knife radiosurgery may provide earlier and longer-lasting pain relief for patients with trigeminal neuralgia. *J Neurosurg*. 2015;123(4):961–968.

6. Mousavi SH, Niranjan A, Akpinar B, et al. A proposed plan for personalized radiosurgery in patients with trigeminal neuralgia. *J Neurosurg*. 2018;128(2):452–459.
7. Obermann M, Rodriguez-Raecke R, Naegel S, et al. Gray matter volume reduction reflects chronic pain in trigeminal neuralgia. *Neuroimage*. 2013;74:352–358.
8. Li M, Yan J, Li S, et al. Reduced volume of gray matter in patients with trigeminal neuralgia. *Brain Imaging Behav*. 2017;11(2):486–492.
9. DeSouza DD, Hodaie M, Davis KD. Abnormal trigeminal nerve microstructure and brain white matter in idiopathic trigeminal neuralgia. *Pain*. 2014;155(1):37–44.

Disclosures

The authors report no conflict of interest.

Correspondence

Luigi Albano: albanoluigi@hsr.it.

INCLUDE WHEN CITING

Published online August 28, 2020; DOI: 10.3171/2020.7.JNS202729.

Response

No response was received from the authors of the original article.

©AANS 2021, except where prohibited by US copyright law

Earlier radiosurgery is related to better outcome in trigeminal neuralgia

TO THE EDITOR: We read with great interest the international multicenter study published by Mureb and coworkers (Mureb M, Golub D, Benjamin C, et al. Earlier radiosurgery leads to better pain relief and less medication usage for trigeminal neuralgia patients: an international multicenter study. *J Neurosurg*. Published online July 3, 2020. doi:10.3171/2020.4.JNS192780).¹ Trigeminal neuralgia (TN) is a common chronic pain disorder, and the first-line treatment for this condition is medical therapy.^{1–3} Despite favorable pain control in a short time period with medical therapy, in about two-thirds of cases patients continue to experience pain at long-term follow-up.³ At this stage, all the available therapeutic options are usually managed by neurosurgeons.^{2,3} To date, there are no homogeneous treatment algorithms, and therapeutic modalities are administered based on surgeon and patient preference, and facility availability at the referring center.³ For these reasons, it is still not possible to individualize therapies based on a patient's characteristics.^{2,3}

Microvascular decompression (MVD) continues to represent the reference surgical therapy, and it is able to provide faster pain relief rates. However, at the same time MVD represents the most “anatomy-preserving” technique and the most invasive one.^{2–5} Radiosurgery, especially Gamma Knife radiosurgery (GKRS), represents the less invasive ablative therapy with lower rates of complications and better long-term results of pain control.^{2–6} Based

on these findings, one of the actual main concerns in TN management is which therapy has to be administered first.²

We would like to congratulate the authors for the quality of their article and the reported results. They found that a pain history of less than 4 years resulted in a shorter interval to pain relief, lower complication rates, and longer duration of pain relief in a series of 404 patients who underwent GKRS as the first surgical therapy after medical failure.¹ Moreover, the authors stressed the capability of GKRS in decreasing pain medication utilization.¹ This study represents the largest one analyzing these aspects in TN management.¹ These findings further confirm that GKRS has to be considered earlier in the management of TN.²

A long history of TN is related to microstructural nervous changes that appear to negatively affect pain response to surgical management.^{2,7} Another important point that was not stressed by the authors is that TN shares with epilepsy and chronic pain the so-called “kindling phenomenon” of second-order neurons into the brainstem.^{2,7} This phenomenon represents the factor that explains why therapies targeting first-order sensory neurons may not fully express their therapeutic potential.^{2,7} The duration of the history of pain is strongly related to the fiber synaptic reorganization of the “kindling phenomenon.”^{2,7}

As we recently reported, GKRS has to be considered early in the treatment algorithm of TN because of its characteristics, i.e., the high possibility to control pain in the first stages of pain and the ability to improve a patient's quality of life.^{1–8} More research on the optimal timing of each treatment modality and tailored treatment algorithms is warranted to determine the best treatment strategy for each patient suffering from this disabling disease.

Alfio Spina, MD

Nicola Boari, MD

Pietro Mortini, MD

I.R.C.C.S. San Raffaele Scientific Institute, Vita-Salute San Raffaele University, Milan, Italy

References

1. Mureb M, Golub D, Benjamin C, et al. Earlier radiosurgery leads to better pain relief and less medication usage for trigeminal neuralgia patients: an international multicenter study. *J Neurosurg*. Published online July 3, 2020. doi: 10.3171/2020.4.JNS192780
2. Spina A, Boari N, Gagliardi F, et al. Gamma Knife radiosurgery for trigeminal neuralgia: when? *Neurosurg Rev*. 2019; 42(2):599–601.
3. Spina A, Mortini P, Alemanno F, et al. Trigeminal neuralgia: toward a multimodal approach. *World Neurosurg*. 2017;103: 220–230.
4. Nanda A, Javalkar V, Zhang S, Ahmed O. Long term efficacy and patient satisfaction of microvascular decompression and gamma knife radiosurgery for trigeminal neuralgia. *J Clin Neurosci*. 2015;22(5):818–822.
5. Spina A, Boari N, Gagliardi F, et al. The emerging role of gamma knife radiosurgery in the management of glossopharyngeal neuralgia. *Neurosurg Rev*. 2019;42(1):31–38.
6. Gagliardi F, Spina A, Bailo M, et al. Effectiveness of Gamma Knife Radiosurgery in improving psychophysical performance and patient's quality of life in idiopathic trigeminal neuralgia. *World Neurosurg*. 2018;110:e776–e785.

7. Lee CC, Chen CJ, Chong ST, et al. Early stereotactic radiosurgery for medically refractory trigeminal neuralgia. *World Neurosurg.* 2018;112:e569–e575.
8. Mousavi SH, Niranjana A, Huang MJ, et al. Early radiosurgery provides superior pain relief for trigeminal neuralgia patients. *Neurology.* 2015;85(24):2159–2165.

Disclosures

The authors report no conflict of interest.

Correspondence

Alfio Spina: spina.alfio@hsr.it.

INCLUDE WHEN CITING

Published online November 13, 2020; DOI: 10.3171/2020.9.JNS203342.

Response

No response was received from the authors of the original article.

©AANS 2021, except where prohibited by US copyright law

Zero complications in SEEG: a goal to pursue

TO THE EDITOR: We read with great interest the study recently published by Rollo et al.,¹ demonstrating the safety of stereo-electroencephalography (SEEG) implantation independently from the angle of entry through the skull (Rollo PS, Rollo MJ, Zhu P, et al. Oblique trajectory angles in robotic stereo-electroencephalography. *J Neurosurg.* Published online August 14, 2020. doi:10.3171/2020.5.JNS20975). This represents the largest series of SEEG electrodes (2040 electrodes) for which implantation accuracy was assessed.

We would like to express our appreciation of the seizure outcomes and technical aspects of the authors' SEEG implantation. They reported that 69.5% of the patients undergoing an SEEG-guided resection had a favorable outcome (Engel class I), which is similar to that of other large studies.^{2,3}

The SEEG method of Rollo et al. entails the use of volumetric contrast-enhanced CT scans coregistered with

FIG. 1. Lateral (A) and anteroposterior (B) schemata of a right-sided SEEG implantation following the Talairach classic methodology reconstructed from tele-stereoscopic ventriculography and angiography. The surgical procedure was performed at our center in 1996. We chose 3 electrodes as examples to illustrate the Talairach concept of orthogonality. Electrode T is a classic orthogonal one, where orthogonality is referred to the grids and the frame planes. Due to its position (perisylvian region), electrode T is orthogonal to the bone. Electrode Q is, again, a classic orthogonal one, even though it is perpendicular only to the double grid and to the frame. This concept is well illustrated in panel C, in which the surgeon implants the bolt orthogonal to the double grid (and not to the bone). Electrode Y is an oblique one, since its implantation was performed with the use of a special support for the grid. We stress the concept that orthogonal trajectories were perpendicular to the sagittal plane and not to the bone. Another concept which is nicely illustrated through panels A and B is that brain regions seated above the higher electrodes (electrodes P, Q, M, E, and F) are undersampled. New methods have helped overcome this issue with optimization of the traditional orthogonal implantation (it is worth mentioning that the positioning of an oblique electrode in the style of Talairach was difficult and time consuming). Figure is available in color online only.

MRI. A recent report highlighted that using MRI as the vascular imaging modality led to a higher number of symptomatic hemorrhages, in comparison with CT angiography and catheter angiography.⁴ Similarly, in our experience, no implantation-related symptomatic hemorrhages occurred with our recent workflow (5181 electrodes).^{3,5}

The application accuracy of implantation is comparable to that in previous reports,^{2,6} and such a rigorous technique reflects on the absence of complications in such a large surgical series. Of note, we do not agree that O-arm–based registration is less accurate, as we have reported even better accuracy on both phantom and clinical conditions with an optimized frameless registration tool,⁷ obtaining results comparable to those of frame-based implantation.⁸

The hypothesis that the entry angle affects errors at the entry point⁹ is confirmed by Rollo et al., with an entry point deviation of 0.45 mm for every 30° angle increase, but the authors highlighted that neither complications nor seizure control was affected by the angle increase because the multivariate model parameters indicated an effect that is irrelevant in practice. Further along this line, the authors suggested that the paradigm of orthogonal trajectories, typical of the classic Talairach methodology, could be mitigated. On this topic, we ought to clarify the traditional concept of Talairach orthogonality. Due to technical constraints, most trajectories were perpendicular to the sagittal plane of the frame and, as a consequence, to the brain midline.¹⁰ Higher trajectories were therefore not perpendicular to the skull, even if orthogonal to the midline (Fig. 1).

To conclude, we appreciate the rigorous surgical technique along with the clinical results, but, most importantly, we strongly agree with the provoking statement, “Why not zero?” related to the possibility of bringing the complication rate in SEEG down to 0%. Zero complications is a goal to pursue, the horizon of our surgical practice, given the data reported in recent large SEEG series.^{1,3}

Michele Rizzi, MD¹

Laura Castana, MD¹

Piergiorgio d’Orio, MD^{1,2}

Francesco Cardinale, MD¹

¹“Claudio Munari” Center for Epilepsy Surgery, ASST GOM Niguarda Hospital, Milan, Italy

²CNR Neuroscience Institute, Parma, Italy

References

1. Rollo PS, Rollo MJ, Zhu P, et al. Oblique trajectory angles in robotic stereo-electroencephalography. *J Neurosurg*. Published online August 14, 2020. doi:10.3171/2020.5.JNS20975
2. Gonzalez-Martinez J, Bulacio J, Thompson S, et al. Technique, results, and complications related to robot-assisted stereoelectroencephalography. *Neurosurgery*. 2016;78(2):169–180.
3. Cardinale F, Rizzi M, Vignati E, et al. Stereoelectroencephalography: retrospective analysis of 742 procedures in a single centre. *Brain*. 2019;142(9):2688–2704.
4. McGovern RA, Butler RS, Bena J, et al. Incorporating new technology into a surgical technique: the learning curve of a single surgeon’s stereo-electroencephalography experience. *Neurosurgery*. 2020;86(3):E281–E289.
5. Cardinale F, Pero G, Quilici L, et al. Cerebral angiography for multimodal surgical planning in epilepsy surgery: de-

scription of a new three-dimensional technique and literature review. *World Neurosurg*. 2015;84(2):358–367.

6. De Benedictis A, Trezza A, Carai A, et al. Robot-assisted procedures in pediatric neurosurgery. *Neurosurg Focus*. 2017;42(5):E7.
7. Cardinale F, Rizzi M, D’Orio P, et al. A new tool for touch-free patient registration for robot-assisted intracranial surgery: application accuracy from a phantom study and a retrospective surgical series. *Neurosurg Focus*. 2017;42(5):E8.
8. Sharma JD, Seunarine KK, Tahir MZ, et al. Accuracy of robot-assisted versus optical frameless navigated stereo-electroencephalography electrode placement in children. *J Neurosurg*. 2019;23(3):297–302.
9. Iordanou JC, Camara D, Ghatan S, et al. Approach angle affects accuracy in robotic stereoelectroencephalography lead placement. *World Neurosurg*. 2019;128:e322–e328.
10. Musolino A, Tournoux P, Missir O, et al. Methodology of “in vivo” anatomical study and stereo-electroencephalographic exploration in brain surgery for epilepsy. *J Neuroradiol*. 1990;17(2):67–102.

Disclosures

Dr. Rizzi: consultant for Wise srl. Dr. Cardinale: consultant (paid expert testimony) to Renishaw Mayfield, manufacturer of the Neuromate robotic system, until February 2019.

Correspondence

Laura Castana: laura.castana@ospedaleniguarda.it.

INCLUDE WHEN CITING

Published online December 25, 2020; DOI: 10.3171/2020.9.JNS203440.

Response

We thank Drs. Rizzi, Castana, d’Orio, and Cardinale for their insightful comments. The Claudio Munari Center for Epilepsy Surgery has pioneered many innovations in SEEG, and the papers they cite are an excellent source of reading for the serious student of this technique.

Low complication rates and the journey to “zero” in all of neurosurgery can be directly related to standardization of optimal techniques and their meticulous implementation.¹ To prevent hemorrhagic complications in SEEG, visualization of vessels, trajectory optimization, robust sub-millimeter registration, and obsessive implantation technique are crucial constituents. Higher complication rates likely result from registration errors, inadvertently mobile constituents of the stereotactic system, and inadequate attention to detail.

SEEG techniques at our center have been invariant since our adoption of robotic stereotaxy in 2013. This use of volumetric CT scans combined with contrasted MRI scans and skull fiducials for registration forms the basis for our work. This perspective is also borne out by experience at other programs.^{2,3} We find the O-arm solution proposed by Cardinale et al. with the Renishaw robot intriguing given their excellent results.² This approach also used skull fiducials and not a scalp surface-based registration, which, in our opinion, is crucial to enable accurate registration. Notably, the integration of the O-arm with digital subtraction angiography (DSA) generates high-fidelity, interactive 3D visual renderings of cerebral vasculature to help guide SEEG placement and minimize complications.^{4,5} This approach does require an additional invasive procedure and,

coupled with 6 O-arm cone-beam CT image acquisitions, is more resource intensive than some centers may be able to tolerate. We accomplish similar safety outcomes at our center where DSA is not incorporated.

We agree that the traditional Talairach-based SEEG technique does indeed employ trajectories at the vertex that are oblique rather than perpendicular to the skull while still orthogonal to the midline—this is distinct from the radial implantations that modern stereotactic frames and robotic techniques can enable. However, the point we make is the distinction between obligatory obliquity (Talairach frame) and optional obliquity (robotic SEEG as implemented here). The latter is motivated by comprehensive sampling of the putative epileptogenic zone with the least number of electrode passes necessary for this purpose. With such an approach, seizure and functional localization are readily achieved. Our outcomes, both from a safety perspective and in regard to seizure freedom, reflect this.

Patrick S. Rollo, BA^{1,2}

Cihan M. Kadipasaoglu, MD, PhD^{1,2}

Nitin Tandon, MD¹⁻³

¹McGovern Medical School at UTHHealth Houston, TX

²Texas Institute for Restorative Neurotechnologies, University of Texas Health Science Center at Houston, TX

³Memorial Hermann Hospital, Texas Medical Center, Houston, TX

References

1. Tandon N, Tong BA, Friedman ER, et al. Analysis of morbidity and outcomes associated with use of subdural grids vs stereoelectroencephalography in patients with intractable epilepsy. *JAMA Neurol*. 2019;76(6):672–681.
2. Cardinale F, Rizzi M, D'Orio P, et al. A new tool for touch-free patient registration for robot-assisted intracranial surgery: application accuracy from a phantom study and a retrospective surgical series. *Neurosurg Focus*. 2017;42(5):E8.
3. McGovern RA, Butler RS, Bena J, Gonzalez-Martinez J. Incorporating new technology into a surgical technique: the learning curve of a single surgeon's stereo-electroencephalography experience. *Clin Neurosurg*. 2020;86(3):E281–E289.
4. Cardinale F, Cossu M, Castana L, et al. Stereoelectroencephalography: surgical methodology, safety, and stereotactic application accuracy in 500 procedures. *Neurosurgery*. 2013;72(3):353–366.
5. Cardinale F, Rizzi M, Vignati E, et al. Stereoelectroencephalography: retrospective analysis of 742 procedures in a single centre. *Brain*. 2019;142(9):2688–2704.

INCLUDE WHEN CITING

Published online December 25, 2020; DOI: 10.3171/2020.10.JNS203452.

©AANS 2021, except where prohibited by US copyright law

The importance of orthogonal implantation in SEEG: historical considerations

TO THE EDITOR: In a recent publication, Rollo et al.¹ statistically compared the anatomical precision in oblique and “traditional orthogonal SEEG” implantations (Rollo

PS, Rollo MJ, Zhu P, et al. Oblique trajectory angles in robotic stereo-electroencephalography. *J Neurosurg*. Published online August 14, 2020. doi:10.3171/2020.5.JNS20975). We would like to clarify specific aspects of this paper by focusing on the historical perspective of the stereo-electroencephalography (SEEG) methodology and the rationale for the so-called orthogonal implantation. The narrative about the novelty of implantation angles does not correspond to any historical background. The term “traditional SEEG” is troublesome, as SEEG has changed only in its technique, not in its principles and methodology. We emphasize that SEEG procedures are part of a method. The attempt to separate the complementary elements of an anatomico-electro-clinical method will bring detrimental misunderstandings.

There is no “traditional SEEG” or neo-SEEG, but rather and simply, SEEG. There are historical reasons for this argument: The SEEG method has been devised by the tight collaboration between Jean Bancaud (neurologist/neurophysiologist) and Jean Talairach (neurosurgeon/anatomist). With growing experience in interpreting SEEG, Bancaud progressively adjusted the implantation types to individual clinical scenarios, to systematically organize and define an SEEG method. In parallel, Talairach implemented the technique accordingly. A “traditional SEEG” with use of “orthogonal trajectories guided by semiology and arteriography” never existed. The method was thoroughly described and explained in detail in previous publications.²⁻⁵ Unfortunately, the references used by the authors are rather anecdotal and incomplete by not providing the landscape of the method's developmental process. In fact, the refinement process of the SEEG method was quite the opposite to what was described: as Talairach and colleagues had designed different angled double grids and multiple attachments possible for them on his frame, the oblique orientations were prevailing in the early years of SEEG. Because one of their main preoccupations in the 1960s was to search for a centrencephalic origin of generalized epilepsies, the deepest leads were often in the thalamus that they reached through oblique trajectories. The trajectories were later refined towards orthogonal orientations, not because of any inaccuracy of the trajectories or any technical limitations, but for an optimal interpretation of the intracerebral recordings and a better understanding of the epileptogenic zone organization. This why the term “traditional SEEG” and its association with orthogonal trajectories are incorrect and misleading.

The notion of “dogma” is problematic. The point missed is that SEEG is not a depth electrode recording technique but an anatomico-electro-clinical method that takes into consideration the 3D organization of the epileptogenic zone, which can only be delineated by the analysis of the temporal-spatial evolution of the epileptogenic activity. For an SEEG evaluation to be accurately interpreted, implantation must not be planned to reach some “specific” areas at the tip of the electrodes but prearranged to allow a rational deciphering of distinctive electrophysiological patterns from a coherent ensemble of trajectories that are based on validated anatomico-functional criteria. This is the reason why relevant anatomico-electro-clinical SEEG data, in general provided by orthogonal electrodes, have led to unique latero-medial tracks as a general basic principle.

Obviously, individual cases may require oblique trajectories, but more as an exception rather than a general rule.⁶

Jorge A. Gonzalez-Martinez, MD, PhD

Patrick Y. Chauvel, MD

University of Pittsburgh Medical Center, Pittsburgh, PA

University of Pittsburgh Medical School, Pittsburgh, PA

References

1. Rollo PS, Rollo MJ, Zhu P, et al. Oblique trajectory angles in robotic stereo-electroencephalography. *J Neurosurg*. Published online August 14, 2020. doi:10.3171/2020.5.JNS20975.
2. Talairach J, David M, Tournoux P. *L'exploration chirurgicale stéréotaxique de lobe temporal dans l'épilepsie temporale*. Masson, libraires de l'Académie de médecine; 1958.
3. Talairach J, Bancaud J, Bonis A, et al. Functional stereotaxic investigations in epilepsy. Methodological remarks concerning a case. Article in French. *Rev Neurol (Paris)*. 1961;105:119–130.
4. Bancaud J, Angelergues R, Bernouilli C, et al. Functional stereotaxic exploration (SEEG) of epilepsy. *Electroencephalogr Clin Neurophysiol*. 1970;28(1):85–86.
5. Talairach J, Tournoux P, Musolino A, Missir O. Stereotaxic exploration in frontal epilepsy. *Adv Neurol*. 1992;57:651–688.
6. Chauvel P, Gonzalez-Martinez J, Bulacio J. Presurgical intracranial investigations in epilepsy surgery. *Handb Clin Neurol*. 2019;161:45–71.

Disclosures

Dr. Gonzalez-Martinez reports ownership in Neurologic Solutions and being a consultant to Zimmer Biomet.

Correspondence

Jorge A. Gonzalez-Martinez: gonzalezjo@upmc.edu.

INCLUDE WHEN CITING

Published online February 5, 2021; DOI: 10.3171/2020.10.JNS203393.

Response

We agree entirely and wholeheartedly that SEEG is not a method; it represents an amalgamation of noninvasive data, the implantation strategy, methods and style for the interpretation of EEG data, and the design of the optimal intervention to control epilepsy in each case. Indeed, advances have occurred in all of these aspects of SEEG, and thanks to both Patrick Chauvel and Jorge Gonzalez-Martinez, we have a much better understanding of the anatomic-electro-clinical correlations of particular epilepsy syndromes,¹ safer and more effective localization of seizures using robotic (whether oblique or not) SEEG,² and more sophisticated derivations of epileptogenicity³ than we had in the last century. Furthermore, in specific scenarios, laser interstitial thermal ablation⁴ represents an improvement over both radiofrequency lesioning and traditional microsurgical resections.

In the current era we are seeing an explosion of the use of SEEG, driven to a large extent by ready access to robotic technology. Some of this, sadly, occurs with an incomplete appreciation of the crucial dependence of successful SEEG on the careful study and characterization of the epilepsy. We completely agree that any “attempt to separate the complementary elements of an anatomic-electro-clinical method” will result in suboptimal and even undesirable

outcomes. That is far from how we view and use SEEG and not the purpose of designing new implantation strategies. There is no doubting the genius and creativity that Talairach and Bancaud displayed in creating the field. We thank Drs. Chauvel and Gonzalez-Martinez for highlighting their desire to record from the thalamus in the early years, something we were not aware of. Yet, we would argue that Bancaud’s development of “implantation types to individual clinical scenarios, to systematically organize and define an SEEG method” were doubtless limited by the technology of the time. Our use of the term “traditional,”⁵ is used specifically and narrowly to illustrate the technical distinctions between our approach and what has been in actual practice for more than 50 years. Restrictive constraint of surgical plans, with fidelity to the status quo, using orthogonal (and this includes oblique obliquity of trajectories that access the superior medial frontal and parietal lobes) trajectories seems unnecessary to us. We show that optional oblique approaches allow for sampling regions not routinely studied using traditional trajectories, and that these are often involved in seizure onsets. It is worth noting that even within the French and Italian schools, there is a tacit understanding that the insula is best sampled with oblique electrodes (e.g., X, Y, X', Y'). Our point is that these need not be the only exceptions.

Ultimately, the style of SEEG implantation should be measured against patient outcomes. Three factors are relevant here: complication rates, localization (SEEG success) rates, and seizure-free (surgery success) rates. We advocate in our paper for efforts to reach a zero clinically relevant complication rate in SEEG coupled with high efficacy. We had a 75% definitive surgical rate, of which 69.5% were seizure free at 1 year, results that are very favorable relative to prior series using principally orthogonal implants.^{5,6} We propose that adoption of some of the technical details as elaborated in our publication may, in fact, result in safer and more effective SEEG. Indeed, without our own “rational deciphering of distinctive electrophysiological patterns from a coherent ensemble of trajectories,” we would not have had such results.

Malum consilium quod mutari non potest, i.e., “Bad is the plan that cannot change,” said Publilius Syrus, in *Maxims*, in the 1st century AD. Changes to established and proven methods are hard, yet inevitable and necessary. We humbly submit that the ingenuity of the epilepsy community, both in the technical and cognitive aspects of SEEG implantation and data interpretation, will inevitably lead to changes in methods developed 60 years ago, and that our paper presages many such changes in the future.

Nitin Tandon, MD¹⁻³

Patrick S. Rollo, BA^{1,2}

¹McGovern Medical School at UTHealth Houston, TX

²Texas Institute for Restorative Neurotechnologies, UTHealth, Houston, TX

³Memorial Hermann Hospital, Texas Medical Center, Houston, TX

References

1. Bonini F, McGonigal A, Trébuchon A, et al. Frontal lobe seizures: from clinical semiology to localization. *Epilepsia*. 2014;55(2):264–277.
2. Mullin JP, Shriver M, Alomar S, et al. Is SEEG safe? A systematic review and meta-analysis of stereo-electroen-

- cephalography-related complications. *Epilepsia*. 2016;57(3):386–401.
3. Grinenko O, Li J, Mosher JC, et al. A fingerprint of the epileptogenic zone in human epilepsies. *Brain*. 2018;141(1):117–131.
 4. Donos C, Breier J, Friedman E, et al. Laser ablation for mesial temporal lobe epilepsy: Surgical and cognitive outcomes with and without mesial temporal sclerosis. *Epilepsia*. 2018;59(7):1421–1432.
 5. Steriade C, Martins W, Bulacio J, et al. Localization yield and seizure outcome in patients undergoing bilateral SEEG exploration. *Epilepsia*. 2019;60(1):107–120.
 6. McGovern RA, Ruggieri P, Bulacio J, et al. Risk analysis of hemorrhage in stereo-electroencephalography procedures. *Epilepsia*. 2019;60(3):571–580.

INCLUDE WHEN CITING

Published online February 5, 2021; DOI: 10.3171/2020.11.JNS203745.

©AANS 2021, except where prohibited by US copyright law

Validating the 5-factor modified frailty index

TO THE EDITOR: I read with interest the article by Khalafallah et al.¹ (Khalafallah AM, Huq S, Jimenez AE, et al. The 5-factor modified frailty index: an effective predictor of mortality in brain tumor patients. *J Neurosurg*. Published online August 14, 2020. doi:10.3171/2020.5.JNS20766) and their related article in *Neurosurgery*.² A considerably simpler tool compared with others^{3,4} to guide prehabilitation and enhance recovery after surgery has great potential in improving outcomes for patients with a brain tumor. I congratulate the authors on their work in validating the 5-factor modified frailty index (mFI-5). However, there are two important aspects of the study that were not discussed by the authors but require careful consideration when integrating the mFI-5 into clinical care.

Khalafallah et al. evaluated 90-day mortality in a cohort of brain tumor patients who had undergone surgery. With the study design used by these authors, the included patients had already been assessed to be suitable candidates for surgery. Using this tool to inform decisions regarding prehabilitation in patients who would otherwise be offered surgery is reasonable.² However, findings from this study do not directly support the use of mFI-5 to aid decision-making in patients for whom the balance of risks for surgery is more challenging, because the nonoperative proportion of this patient group was not included. Furthermore, though a thoughtful suggestion, prehabilitation may not modify the risks of unfavorable outcomes. The potential for benefitting from prehabilitation may also be associated with the mFI-5 score; however, the risk factors contributing to the higher 90-day mortality may not be modifiable by prehabilitation.

Brain tumors are a heterogeneous group of diseases associated with a range of prognoses. It is not surprising that 83% (n = 44/53) of the deaths within 90 days occurred in patients with high-grade glioma or brain metastases (Table 3 in the Khalafallah et al. study report¹). In the multiple logistic regression for 90-day mortality, the large number

of variables included resulted in fewer than 4 events per variable. While the low number of events per variable does not dismiss the results, the risk of model overfitting and the close association between tumor diagnosis and the primary outcome may call into question the generalizability of the mFI-5. One way to examine this would be to stratify the analyses by tumor type, negating the need to include tumor diagnosis as a confounder and allowing comparison of mFI-5 performance by tumor type. Another way, which no doubt the authors are seeking to do, is to validate their findings in another cohort with stratification.

Validating a simple tool such as mFI-5 to identify patients who can benefit from perioperative optimization is desirable. Khalafallah et al. have contributed a transparent study with results that justify further examination of mFI-5 in brain tumor patients.^{1,2} We should strive to adopt the most rigorous approach to validate risk stratification strategies for our patients.

Michael T. C. Poon, MBChB, MSc, MRCS(Ed)

Usher Institute, University of Edinburgh, United Kingdom

Royal Infirmary of Edinburgh, United Kingdom

Acknowledgments

Cancer Research UK Brain Tumour Centre of Excellence Award (C157/A27589).

References

1. Khalafallah AM, Huq S, Jimenez AE, et al. The 5-factor modified frailty index: an effective predictor of mortality in brain tumor patients. *J Neurosurg*. Published online August 14, 2020. doi:10.3171/2020.5.JNS20766
2. Huq S, Khalafallah AM, Jimenez AE, et al. Predicting postoperative outcomes in brain tumor patients with a 5-factor modified frailty index. *Neurosurgery*. Published online August 17, 2020. doi:10.1093/neuros/nyaa335
3. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373–383.
4. Velanovich V, Antoine H, Swartz A, et al. Accumulating deficits model of frailty and postoperative mortality and morbidity: its application to a national database. *J Surg Res*. 2013;183(1):104–110.

Disclosures

The author reports no conflict of interest.

Correspondence

Michael T. C. Poon: mpoon@ed.ac.uk.

INCLUDE WHEN CITING

Published online November 13, 2020; DOI: 10.3171/2020.8.JNS203241.

Response

We thank Dr. Poon for his thoughtful response to our recent articles published in *Journal of Neurosurgery* and *Neurosurgery*.

We agree with the author's point that our study results only demonstrate the utility of the mFI-5 in risk stratifying patients who have already been defined as suitable candidates for surgery. However, the specificity of this application of the mFI-5 should not necessarily hamper its imple-

mentation into clinical workflows, particularly following rigorous external validation of this index of frailty. As has been discussed by previous authors, frailty indices like the Charlson Comorbidity Index (CCI), the 11-factor modified frailty index (mFI-11), and the mFI-5 are meant to serve as useful, practical tools for preoperative risk assessment and patient stratification, though these indices represent only a single part of an extensive workup incorporating age and other relevant risk factors.¹⁻⁵ We do not believe that preoperative risk assessment could ever be reduced to a single metric, but we do believe that the mFI-5, as an easy-to-calculate, objective measure of patient frailty that has demonstrated significant associations with many different postoperative outcomes in a variety of surgical specialties, shows promise as an addition to the neurosurgeon's current armament of preoperative risk assessment tools. In this sense, the mFI-5 may still contribute meaningfully in the risk stratification of patients in whom the "balance of risks for surgery is more challenging" if utilized alongside other important metrics such as American Society of Anesthesiology (ASA) class and Karnofsky Performance Status (KPS). As noted in our study limitations, further prospective studies examining the prognostic utility of the mFI-5 among brain tumor patients will be necessary to rigorously demonstrate the usefulness of this index in aiding clinical decision-making.

Regarding the author's point about the utility of prehabilitation, we agree that prehabilitation may not fully modify the risks of unfavorable outcomes. We mentioned prehabilitation as one possible way in which the mFI-5 could be incorporated into clinical workflows, and we make no strong claim as to its efficacy in optimizing surgical outcomes specifically among brain tumor patients. Though the author makes a reasonable comment in questioning the efficacy of prehabilitation given the dearth of research investigating this intervention among brain tumor patients specifically, it is important to note that prehabilitation has demonstrated improved postoperative outcomes among both abdominal surgery and lumbar spine surgery patients.^{6,7} Further research beyond the scope of the present studies will be necessary to determine the importance and efficacy of prehabilitation in optimizing outcomes among brain tumor patients specifically.

The author also presents valid concerns regarding model generalizability and overfitting. Ultimately, we believe these concerns would be best addressed through external validation of the mFI-5 and our multivariate models using larger, multicenter brain tumor patient cohorts. We agree that mFI-5 performance within each tumor type will likely be a fruitful avenue for future research, and we have recently pursued this line of inquiry among patients with pituitary adenomas.⁸ Our rationale for including brain tumor diagnosis in the present studies was specifically to investigate whether mFI-5 would remain associated with postoperative outcomes even when controlling for diagnosis. Thus, the fact that we observed a significant association between mFI-5 score and 90-day postoperative mortality even when controlling for diagnosis may suggest that the mFI-5 provides important prognostic information independent of brain tumor type. Again, we agree with the author that specific concerns regarding model overfitting

must be assessed via external validation before the mFI-5 may be fully utilized in clinical settings.

We hope our study results may serve as an impetus for prospective research examining the utility of the mFI-5 in aiding neurosurgeons in preoperatively risk stratifying brain tumor patients. We thank Dr. Poon for engaging with our work and for emphasizing the importance of rigorous external validation in the development of surgical risk stratification tools.

Adham M. Khalafallah, MD

Adrian E. Jimenez, BS

Debraj Mukherjee, MD, MPH

Johns Hopkins University School of Medicine, Baltimore, MD

References

1. Ali R, Schwab JM, Nerenz DR, et al. Use of the modified frailty index to predict 30-day morbidity and mortality from spine surgery. *J Neurosurg Spine*. 2016;25(4):537-541.
2. Grossman R, Mukherjee D, Chang DC, et al. Preoperative Charlson comorbidity score predicts postoperative outcomes among older intracranial meningioma patients. *World Neurosurg*. 2011;75(2):279-285.
3. Tsiouris A, Hammoud ZT, Velanovich V, et al. A modified frailty index to assess morbidity and mortality after lobectomy. *J Surg Res*. 2013;183(1):40-46.
4. Weaver DJ, Malik AT, Jain N, et al. The modified 5-item frailty index: a concise and useful tool for assessing the impact of frailty on postoperative morbidity following elective posterior lumbar fusions. *World Neurosurg*. 2019;124:e626-e632.
5. Youngerman BE, Neugut AI, Yang J, et al. The modified frailty index and 30-day adverse events in oncologic neurosurgery. *J Neurooncol*. 2018;136(1):197-206.
6. Howard R, Yin YS, McCandless L, et al. Taking control of your surgery: impact of a prehabilitation program on major abdominal surgery. *J Am Coll Surg*. 2019;228(1):72-80.
7. Nielsen PR, Andreasen J, Asmussen M, Tønnesen H. Costs and quality of life for prehabilitation and early rehabilitation after surgery of the lumbar spine. *BMC Health Serv Res*. 2008;8:209.
8. Khalafallah AM, Shah PP, Huq S, et al. The 5-factor modified frailty index predicts health burden following surgery for pituitary adenomas. *Pituitary*. 2020;23(6):630-640.

INCLUDE WHEN CITING

Published online November 13, 2020; DOI: 10.3171/2020.9.JNS203271.

©AANS 2021, except where prohibited by US copyright law

Multidisciplinary meetings for neuro-oncology services

TO THE EDITOR: Our group read with great enthusiasm the article published by Khalafallah et al.¹ focusing on the value of multidisciplinary meetings (MDMs) or tumor boards as an essential component of a comprehensive neuro-oncology service (Khalafallah AM, Jimenez AE, Romo CG, et al. Quantifying the utility of a multidisciplinary neuro-oncology tumor board. *J Neurosurg*. Published online September 18, 2020. doi:10.3171/2020.5.JNS201299). We share the opinion that MDMs are one of the most sig-

nificant advances in neuro-oncology—alongside or even more so than the technological and therapeutic advances—responsible for improved patient outcomes.^{2,3}

In the United Kingdom, this move toward an MDM-based structure was driven by the National Institute for Health and Clinical Excellence document *Improving Outcomes for People with Brain and Other CNS Tumours* published in June 2006.⁴ The team consists of a designated lead, accountable for the overall service and core members with subspecialist expertise in neuro-oncology. These include brain tumor neurosurgeons, neuro-oncologists, neurologists, neuroradiologists, neuropathologists, neurotherapists (speech and language, physical and occupational), clinical nurse specialists, palliative care physicians, and coordinators with access to extended members such as neuropsychologists as needed. The critical initial management plan for nearly all brain tumor patients was made in a structured, consistent, and “semi-elective” way by a group of subspecialist clinicians, rather than by a single individual, often in a “semi-emergency” setting, as was the case prior to publication of this document. The document did acknowledge that emergency decision-making outside of the MDM was still warranted for a very small minority of patients presenting acutely, for example, in a coma or with hydrocephalus; they, however, still needed to be discussed in the MDM postsurgery to plan ongoing treatment.^{2,3}

As highlighted by Khalafallah et al.,¹ the MDM is a perfect example of an expert team working together with a single goal of providing the best care possible for each individual patient. In addition, the MDM is an optimal setting, not only to train the next generation of clinicians but also for individual specialists to learn from colleagues. The MDM also provides a powerful platform to develop a cohesive collaborative academic approach translating into a significant research output.

Given our experience with more than 15 years of MDM working, at our institution, we have extended this concept beyond the first presentation time point to every stage of the patient journey. We believe that our patients deserve access to the full MDM expertise not just at diagnosis but throughout their care. Thus, before each neuro-oncology follow-up clinic, we hold an MDM attended by the core members, where all the clinical information and imaging are reviewed and a collective decision on further management is made. This MDM clinic approach has recently been recognized and advocated by the Tessa Jowell Brain Cancer Mission as an integrated care system to be utilized for brain tumor patients at every stage of their care.⁵

The current COVID-19 pandemic has presented us with immense challenges, and many of our patients may have limited access to face-to-face consultations or contact. Perhaps, at no other time has the value of MDM working been more evident, to support both our patients and each other.^{6,7}

José P. Lavrador, MD

Graeme Pang, MBChB

Francesco Vergani, PhD, FRCS

Ranjeev Bhangoo, FRCS(Eng), FRCS(SN)

Richard Gullan, FRCP, FRCS

Keyoumars Ashkan, FRCP, FRCS

King's College Hospital NHS Foundation Trust, London, United Kingdom

References

1. Khalafallah AM, Jimenez AE, Romo CG, et al. Quantifying the utility of a multidisciplinary neuro-oncology tumor board. *J Neurosurg*. Published online September 18, 2020. doi:10.3171/2020.5.JNS201299
2. deSouza RM, Shaweis H, Han C, et al. Has the survival of patients with glioblastoma changed over the years? *Br J Cancer*. 2016;114(2):146–150.
3. Rittman T, Corns R, Kumar A, et al. Is referral to the neuro-oncology MDT safe? *Br J Neurosurg*. 2012;26(3):321–324.
4. National Institute for Health and Clinical Excellence. *Improving Outcomes for People with Brain and Other CNS Tumours*. Published June 2006. Accessed February 1, 2021. <https://www.nice.org.uk/guidance/csg10/resources/improving-outcomes-for-people-with-brain-and-other-central-nervous-system-tumours-update-27841361437>
5. Tessa Jowell Brain Cancer Mission. Tessa Jowell Centre Designation. Accessed February 1, 2021. <https://www.tessajowellbraincancermission.org/strategic-programmes/tessa-jowell-badges-for-neurocentres/>
6. Ashkan K, Jung J, Velicu AM, et al. Neurosurgery and coronavirus: impact and challenges—lessons learnt from the first wave of a global pandemic. *Acta Neurochir (Wien)*. 2021; 163(2):317–329.
7. Ghimire P, Lavrador JP, Onyiriuka L, et al. Patient-reported experience measure for neuro-oncology telephone clinics during the COVID-19 pandemic. *Clin Oncol (R Coll Radiol)*. 2021;33(1):e87.

Disclosures

The authors report no conflict of interest.

Correspondence

Graeme Pang: graeme.pang@nhs.net.

INCLUDE WHEN CITING

Published online February 26, 2021; DOI: 10.3171/2020.12.JNS204247.

Response

We thank Lavrador et al. for their thoughtful response to our recent article.

We agree with the authors that multidisciplinary neuro-oncology tumor boards (MDTBs) are an important advancement in the care of neuro-oncology patients. While the safety and utility of MDTBs are beginning to be established across many medical specialties, research has not yet rigorously quantified the effect of MDTB referral on important postoperative outcomes such as discharge disposition, readmission, and overall survival.^{1–4} Furthermore, the effect of MDTB referral on peri- and postoperative patient morbidity measured using general quality-of-life (QOL) tools such as the PROMIS-29 (Patient-Reported Outcomes Measurement Information System) or more specialized instruments such as the NfBrSI-24 (24-item National Comprehensive Cancer Network/Functional Assessment of Cancer Therapy Brain Symptom Index) has also not been studied.^{5,6} Such future research efforts will be important in fully understanding the role of MDTBs in optimizing patient outcomes and coordination of multidisciplinary care. With increasing focus on high-value care across subspecialties, it is our opinion that these investigations will be necessary to justify the continuation and expansion of these collaborative care teams, especially during periods such as the ongoing COVID-19 pandemic, which impose severe limitations on healthcare resources.^{7–9}

We hope our study serves as an impetus for further prospective research examining the utility of the MDTBs in affecting important postoperative outcomes such as postoperative complications, QOL metrics, and overall survival. We thank Lavrador et al. for engaging with our work and for emphasizing the importance of interdisciplinary collaboration through MDTBs in the care of neuro-oncology patients.

Adham M. Khalafallah, MD
Adrian E. Jimenez, BS
Debraj Mukherjee, MD, MPH

Johns Hopkins University School of Medicine, Baltimore, MD

References

1. Lee B, Kim K, Choi JY, et al. Efficacy of the multidisciplinary tumor board conference in gynecologic oncology: a prospective study. *Medicine (Baltimore)*. 2017;96(48):e8089.
2. Rittman T, Corns R, Kumar A, et al. Is referral to the neuro-oncology MDT safe? *Br J Neurosurg*. 2012;26(3):321–324.
3. deSouza RM, Shaweis H, Han C, et al. Has the survival of patients with glioblastoma changed over the years? *Br J Cancer*. 2016;114(2):146–150.
4. Wheless SA, McKinney KA, Zanation AM. A prospective study of the clinical impact of a multidisciplinary head and neck tumor board. *Otolaryngol Head Neck Surg*. 2010;143(5):650–654.
5. Cella D, Yount S, Rothrock N, et al. The Patient-Reported Outcomes Measurement Information System (PROMIS): progress of an NIH Roadmap cooperative group during its first two years. *Med Care*. 2007;45(5 Suppl 1):S3–S11.
6. Lai J-S, Jensen SE, Beaumont JL, et al. Development of a symptom index for patients with primary brain tumors. *Value Health*. 2014;17(1):62–69.
7. Jones LK Jr. The roadmap to value-based care. *JAMA*. 2016; 73(10):1173–1174.
8. Emanuel EJ, Persad G, Upshur R, et al. Fair allocation of scarce medical resources in the time of Covid-19. *N Engl J Med*. 2020;382(21):2049–2055.
9. Khalafallah AM, Jimenez AE, Lee RP, et al. Impact of COVID-19 on an academic neurosurgery department: The Johns Hopkins experience. *World Neurosurg*. 2020;139:e877–e884.

INCLUDE WHEN CITING

Published online February 26, 2021; DOI: 10.3171/2021.1.JNS204273.

©AANS 2021, except where prohibited by US copyright law

Neurosurgical residency abroad: chasing a dream in unknown lands

TO THE EDITOR: We read with great interest the article by Morcos¹ (Morcos JJ. Brief encounters that last a lifetime: an immigrant neurosurgeon's reflections on American exceptionalism, George Floyd, sunlight, and race. *J Neurosurg*. 2020;133[5]:1612–1615). The author, through lived experiences, tries to make us aware of the importance that small moments play in the life of an immigrant professional, focusing on his devotion to the career of medicine as he crossed all the barriers that appear on that complex road. First of all, we want to thank the author for

sharing such a moving anecdote, which will surely have a positive impact on all readers of this prestigious journal.

As stated in the article, every doctor has the fortune or misfortune (depending on the perspective) of experiencing such narrated events. These experiences do not necessarily have to be so traumatic and heartbreaking to have such a revealing effect. Some of the mentioned moments can overflow with happiness, gratitude, and well-being for those who live it, without the need for a drop of suffering or pain. And if we thought that injustices were left behind in the past, we are wrong, because, in 2020, George Floyd suffered the agony of begging for his life to a man who lacked mercy or compassion.¹ There are then thousands of situations from which a positive message could be drawn, that have applicability in our medical exercise, with just having a little more empathy.

There are groups of people called “dreamers,” who have far greater aspirations than their present situation. A clear example of these people are the so-called international medical graduates,² who go in search of unfamiliar lands that welcome them to carry out their residencies and promote their careers. It is these kinds of experiences that generate awareness about all the work that needs to be done to build a just and promising future for future generations that guarantees the best possible quality of life. All this to surpass themselves and to demonstrate that, despite any heartbreaking circumstance, the impetus is there to provide security and hope to those in need, as did that American marine, despite the discrimination and hatred to which he was subjected for not being a Lebanese citizen.¹ That's right; that marine of the unknown name was a dreamer who, without thinking it, inspired the story of another great dreamer, a neurosurgeon, who illuminates students, doctors, and specialists who in essence seek the same thing with his words: to pursue their dreams and discover their happy place and ideal profession.

In Latin America, a region with high rates of corruption and inequality in the world,³ this path becomes more difficult because there is no solid support from the state for those who do not belong to a good socioeconomic class and cannot access a university that has active agreements for international exchanges and internships. Likewise, there are still sociocultural barriers such as scientific racism, where opportunities are overshadowed by the fact of not belonging to societies of reputation or being from distant places where one aspires to participate, wasting the scientific potential of those professionals.

There are obstacles that will have to be overcome, such as the economic gap, the admission of dreamers, the government policies of the elected country and the institutional policies of a university, and the time limit, among many others,⁴ but, for a dreamer, these difficulties will not be greater than their aspirations. On the contrary, they will be an opportunity to better prepare for the competition.

Those of us who are looking for a neurosurgical residency abroad, looking for a better quality of life and preparation under reconnaissance tutors, hope that many of these barriers will disappear to facilitate this process and so that we can have the opportunity to demonstrate that from anywhere in the world, medical-scientific potential can be found.

Juan Maiguel-Lapeira, MS
Ivan Lozada-Martinez, MS
Daniela Torres-Llinás, MS

Medical-Surgical Research Center, University of Cartagena,
 Cartagena, Colombia

Luis Moscote-Salazar, MD

Biomedical Research Center, University of Cartagena,
 Cartagena, Colombia

References

1. Morcos JJ. Brief encounters that last a lifetime: an immigrant neurosurgeon's reflections on American exceptionalism, George Floyd, sunlight, and race. *J Neurosurg*. 2020;133(5):1612–1615.
2. Kuczewski MG, Brubaker L. Medical education for “Dreamers”: barriers and opportunities for undocumented immigrants. *Acad Med*. 2014;89(12):1593–1598.
3. Pring C, Vrushni J. Global corruption barometer: Latin America & the Caribbean 2019—citizens' views and opinions of corruption. Transparency International. Accessed January 26, 2021. <https://www.transparency.org/en/publications/global-corruption-barometer-latin-america-and-the-caribbean-2019#>
4. Berns JS, Ghosn M, Altamirano RR. International medical graduates in nephrology: roles, rules, and future risks. *Am J Kidney Dis*. 2018;72(1):113–117.

Disclosures

The authors report no conflict of interest.

Correspondence

Ivan Lozada-Martinez: ivandavidloma@gmail.com.

INCLUDE WHEN CITING

Published online February 19, 2021; DOI: 10.3171/2020.11.JNS204112.

Response

I thank the authors very much for their interest in my nontraditional publication. It was borne from the unsettling socioeconomic times our nation is facing. I am delighted that it also inspired the authors to extend the theme of my essay and include their own reflections about what it means to be a dreamer on the global neurosurgical stage. Inequities with regard to access to a quality neurosurgical education abound everywhere in our world and permeate every level: race, gender, ethnicity, religion, and economical and geographical factors, among others. Our global community of 50,000 neurosurgeons has been quite adept at describing the problems but is not very well equipped at solving them. To solve them, a true and highly complex multidisciplinary approach is needed, one that relies on the buy-in and coordination of nations, governments, and institutions as well as the corpus of neurosurgery: not an easy task.

Neurosurgery remains the “queen of the surgical disci-

plines” as far as I am concerned. We seem to live in a time where some of my generation find it trendy to perhaps pre-judge those young millennial medical students that are said to be attracted to softer paths in life; whose souls some say lack the “right stuff,” the grit and focus that surely define the quintessential neurosurgeon of yesteryears. To those skeptics who would like to believe that the generational gap has doomed the future of neurosurgery, I say look no further than the makeup of candidates for neurosurgical residency positions today. It never ceases to amaze me how more driven, brilliant, and accomplished each match group is, year after year. Competition for residency training positions in the US is fiercer than ever, because the talent pool is stronger than ever. Yes, indeed, “foreign graduates”—like my former self at one time—still face today an enormous figurative wall. It may feel unfair, as it certainly did when I faced it, but it is entirely understandable. It is not a wall of discrimination but a wall of opportunity. To the high jumper, it will be just a pebble on the road, a stepping-stone to a future of great accomplishments.

I commend the authors for recognizing that indeed “these difficulties will not be greater than their aspirations.” But, unlike the authors, I do not wish that these barriers disappear. I only wish them to be equal-opportunity barriers, applicable, in different forms, and perhaps at different stages, to foreign and domestic applicants alike. Only then will we know that our mission as educators is being fulfilled, and that we have selected to train the best hearts and minds available today that will be equipped tomorrow to outshine their mentors and carry neurosurgery to a level unimaginable to their teachers' generation.

I look forward to one day seeing the authors, in their personal quest to follow their dreams, on top of that wall. They will love the landscape from there. When I do, I will remind them that dreaming is not a prerogative of the young. In fact, like a heartbeat, it never ceases in a fulfilled mind, and a mind that stops dreaming is a spirit that dies. They will in time realize that dreaming is not a temporary means to an end, but a way of life. It is a good thing, when done well. But they do not need to take it from me. A much more eloquent writer said it:

I prefer to be a dreamer among the humblest, with visions to be realized, than lord among those without dreams and desires.

— *Gibran Khalil Gibran*

Jacques J. Morcos, MD, FRCS(Eng), FRCS(Ed)
 University of Miami, FL

INCLUDE WHEN CITING

Published online February 19, 2021; DOI: 10.3171/2021.1.JNS204133.

©AANS 2021, except where prohibited by US copyright law