Neurosurgical Forum LETTERS TO THE EDITOR

Risk assessment of rupture in patients with small unruptured cerebral aneurysms

TO THE EDITOR: Ikawa et al.¹ evaluated the rupture risk in patients with small unruptured cerebral aneurysms (Ikawa F, Morita A, Tominari S, et al. Rupture risk of small unruptured cerebral aneurysms. *J Neurosurg*. 2020;132[1]:69–78). Treatment and rupture numbers of small unruptured cerebral aneurysms (UCAs) were 1132 and 23, respectively. The adjusted hazard ratios (95% confidence intervals [CIs]) of a screening brain checkup, a history of subarachnoid hemorrhage (SAH), uncontrolled hypertension, and location on the anterior communicating artery (ACoA) for rupture were 4.1 (1.2–14.4), 10.8 (2.3–51.1), 5.2 (1.8–15.3), and 5.0 (1.6–15.5), respectively. I have two comments on their study.

First, the authors concluded that the possible selection criteria for treating small UCAs included a history of SAH, uncontrolled hypertension, location on the ACoA, and young patients, which was inconsistent with multivariate analysis. Although the log-rank test showed no significant association of rupture with screening brain checkup, there is no statistical evidence of selecting young patients as a risk factor for rupture.

Second, they handled a small number of events for the multivariate analysis. Peduzzi et al. evaluated the effect of events per independent variable (EPV) in proportional hazards regression analysis.^{2,3} One conclusion of this evaluation was that EPV values < 10 have some problems in maintaining validity of the statistical model. Although there is an opinion that EPV values < 10 are also acceptable for investigating the association,⁴ a prediction model requires a stable number of events. Regarding this point, results presented by Ikawa et al. need further statistical considerations. The number of events was 23, and 2 independent variables can be used for the analysis. They presented relatively wide 95% CI ranges, which might be derived from unstable estimates. In any case, I recommend that EPV should be increased by using a longer follow-up duration or involving much larger sample sizes for risk estimation.

Tomoyuki Kawada, MD
Nippon Medical School, Tokyo, Japan

References

1. Ikawa F, Morita A, Tominari S, et al. Rupture risk of small unruptured cerebral aneurysms. *J Neurosurg*. 2020;132(1):69–78.
2. Concato J, Peduzzi P, Holford TR, Feinstein AR. Importance of events per independent variable in proportional hazards analysis. I. Background, goals, and general strategy. *J Clin Epidemiol*. 1995;48(12):1495–1501.
3. Peduzzi P, Concato J, Feinstein AR, Holford TR. Importance of events per independent variable in proportional hazards regression analysis. II. Accuracy and precision of regression estimates. *J Clin Epidemiol*. 1995;48(12):1503–1510.
4. Vittinghoff E, McCulloch CE. Relaxing the rule of ten events per variable in logistic and Cox regression. *Am J Epidemiol*. 2007;165(6):710–718.

Disclosures

The author reports no conflict of interest.

Correspondence

Tomoyuki Kawada: kawada@nms.ac.jp.

INCLUDE WHEN CITING

Published online November 5, 2021; DOI: 10.3171/2021.8.JNS211955.

Response

We thank Prof. Tomoyuki Kawada for his important comments on our article. We would like to respond to these comments and questions.

As Professor Kawada commented, there is no statistical evidence for selecting young patients as a risk factor of rupture, and we could agree with your reasonable comment. Although patient age was not a significant variable in our multivariate Cox proportional hazards analysis, the patients with treated UCAs were significantly younger than the patients with untreated UCAs (see Tables 4 and 5). As discussed in our article, younger age was a significant risk factor for rupture in small UCAs in previous reports.^{1,2} Additionally, young age was the common existing variable of surgical indication for UCA.^{3,4} Therefore, we added the young patients to the possible indication of interventional treatment for UCA in our conclusions.

Second, we also appreciate Professor Kawada pointing out the effect of EPV in our proportional hazards regression analysis. We should have discussed that point in our paper, as you mentioned. The small EPV in our study could lead to unstable estimates and overfitted models.

As we stated in the section of our paper on study limitations, this study was limited by the short follow-up dura-

tion based on post hoc analysis, not the planned prospective study targeting only the natural course of small UCAs (Unruptured Cerebral Aneurysm Study [UCAS] Japan); therefore, we conclude that further large prospective and longitudinal registered trials are needed to elucidate these issues. However, our study contained one of the largest cohorts of patients with UCAs in the world, and we believe the study can provide valuable information for neurosurgeons worldwide.

Fusao Ikawa, MD, PhD¹

Akio Morita, MD, PhD²

Shinjiro Tominari, MD, PhD³

The Japan Neurosurgical Society for UCAS Japan Investigators

¹Graduate School of Biomedical and Health Sciences, Hiroshima University, Hiroshima, Japan

²Nippon Medical School, Tokyo, Japan

³Kyoto University School of Public Health, Kyoto, Japan

References

1. Sonobe M, Yamazaki T, Yonekura M, Kikuchi H. Small unruptured intracranial aneurysm verification study: SUAVE study, Japan. *Stroke*. 2010;41(9):1969-1977.
2. Suzuki T, Takao H, Rapaka S, et al. Rupture risk of small unruptured intracranial aneurysms in Japanese adults. *Stroke*. 2020;51(2):641-643.
3. Etminan N, Brown RD Jr, Beseoglu K, et al. The unruptured intracranial aneurysm treatment score: a multidisciplinary consensus. *Neurology*. 2015;85(10):881-889.
4. Juvela S. Treatment scoring of unruptured intracranial aneurysms. *Stroke*. 2019;50(9):2344-2350.

INCLUDE WHEN CITING

Published online November 5, 2021; DOI: 10.3171/2021.9.JNS211981.

©AANS 2022, except where prohibited by US copyright law

Clinical Rating Scale for Tremor: a needed clarification

TO THE EDITOR: We read with great interest the article of Sinai and colleagues¹ and we would like to clarify some issues related to the Clinical Rating Scale for Tremor (CRST), one of the most commonly used tremor assessment scales (Sinai A, Nassar M, Eran A, et al. Magnetic resonance-guided focused ultrasound thalamotomy for essential tremor: a 5-year single-center experience. *J Neurosurg*. 2020;133[2]:417-424).

First published in 1988, the CRST was developed by Fahn, Tolosa, and Marin, and contains three parts.² Part A quantifies tremor at rest (R) and/or on action (postural [P] or kinetic [K]) of the following body sites: 1) face (R); 2) tongue (R/P); 3) voice (K); 4) head (R/P); 5) trunk (R/P); 6 and 7) right and left upper limbs (R/P/K); and 8 and 9) right and left lower limbs (R/P/K). Part B assesses kinetic tremor of the upper limbs during writing (only dominant hand), three drawing tasks, and water pouring (both hands). Part C assesses the functional disability and includes speaking, eating, drinking, hygiene, dressing, writing, and working.

Each item is rated on a scale from 0 to 4. The maximum

possible scores of the first version of CRST are 80, 36, and 28 for Parts A, B, and C, respectively; thus the maximum possible total score is 144.² In 1993, a revised version of the CRST was published adding two items to part A (face postural and orthostatic tremor) and social activities to Part C, thus resulting in a maximum possible score of 156.³

There are several noteworthy inaccuracies in the original CRST papers.^{2,3} The first version² did not contain the “face (P)” item either in the description or in the calculations part; however, a scoring box was provided for it in the scorecard. In the second version,³ the “face (P)” score was written in the text but the assessment method while posture holding was not specified. Another inaccuracy of this version is the presence of a box for the “voice (P)” item in the scorecard, although it was not referred to in the text and calculations. In fact, face tremor cannot be assessed while holding a posture. In our clinical practice we have been using a modified version of the CRST omitting the “face (P)” item with a maximum score of 152 instead of 156. However, to keep in line with the existing literature, we have been forced to use the original 156-point scale in our prospective trials.

A version of the CRST with a 160-point maximum score as stated by Sinai and colleagues¹ and Xiong and colleagues in their response letter⁴ has never existed. The confusion may result from two scoring boxes placed near the handwriting section (scored only for the dominant hand) in the original article.³ A similar error exists in a *Neurosurgical Focus* article,⁵ in which the maximum CRST and Part A score was noted as 160 and 32, instead of 156 and 88, respectively.

Although available digitally,⁶ the original CRST forms may be challenging to access. We note that unreviewed web sources exist that share problematic forms. To avoid such misuse, herein we share the CRST form designed for the Bilateral focused ultrasound thalamotomy for essential tremor (BEST-FUS) trial in a sequence optimized for video-recorded physical examinations.^{7,8}

Can Sarica, MD¹

Anton Fomenko, MD¹

Christian Iorio-Morin, MD, PhD²

Ajmal Zemmar, MD, PhD^{1,3,4}

Kazuaki Yamamoto, MD¹

Artur Vetkas, MD, PhD^{1,5}

Andres M. Lozano, MD, PhD^{1,6}

Alfonso Fasano, MD, PhD⁶⁻⁸

¹Toronto Western Hospital, University of Toronto, ON, Canada

²Université de Sherbrooke, QC, Canada

³Henan University School of Medicine, Zhengzhou, China

⁴University of Louisville, School of Medicine, Louisville, KY

⁵School of Medicine, University of Tartu, Estonia

⁶Krembil Brain Institute, Toronto, ON, Canada

⁷Morton and Gloria Shulman Movement Disorders Clinic, Toronto Western Hospital, UHN, Toronto, ON, Canada

⁸University of Toronto, ON, Canada

References

1. Sinai A, Nassar M, Eran A, et al. Magnetic resonance-guided focused ultrasound thalamotomy for essential tremor: a

- 5-year single-center experience. *J Neurosurg.* 2020;133(2):417-424.
2. Fahn S, Tolosa E, Marin C. Clinical rating scale for tremor. In: Jankovic J, Tolosa E, eds. *Parkinson's Disease and Movement Disorders.* Urban & Schwarzenberg; 1988:225-234.
 3. Fahn S, Tolosa E, Marin C. Clinical rating scale for tremor. In: Jankovic J, Tolosa E, ed. *Parkinson's Disease and Movement Disorders.* 2nd ed. Williams & Wilkins; 1993:271-280.
 4. Xiong Y, He J, Lou X. Errors about the use of the Clinical Rating Scale for Tremor score. Letter. *J Neurosurg.* 2020; 133(2):605.
 5. Mohammed N, Patra D, Nanda A. A meta-analysis of outcomes and complications of magnetic resonance-guided focused ultrasound in the treatment of essential tremor. *Neurosurg Focus.* 2018;44(2):E4.
 6. University Consortium. Hathitrust Digital Library. Accessed September 17, 2021. <https://www.hathitrust.org>
 7. Iorio-Morin C, Yamamoto K, Sarica C, et al. Bilateral focused ultrasound thalamotomy for essential tremor (BEST-FUS phase II trial). *Mov Disord.* Published online July 20, 2021. doi:10.1002/mds.28716
 8. Iorio-Morin C, Yamamoto K, Sarica C, et al. Bilateral focused ultrasound thalamotomy for essential tremor (BEST-FUS phase II trial). Scorecard. July 2021. Accessed September 17, 2021. https://www.researchgate.net/publication/353437520_CRST_scorecardpdf

Disclosures

Dr. Lozano is a consultant for Medtronic, Abbott, Boston Scientific, and Insightec. He is scientific director of Functional Neuromodulation.

Correspondence

Can Sarica: can.sarica@mail.utoronto.ca.

Response

No response was received from the authors of the original article.

INCLUDE WHEN CITING

Published online November 5, 2021; DOI: 10.3171/2021.7.JNS211783.

Response

No response was received from the authors of the original article.

©AANS 2022, except where prohibited by US copyright law

The place of brain biopsy in patients with presumed CNS relapse of systemic lymphoma

TO THE EDITOR: We read with great interest the article by Brown et al.¹ (Brown DA, Goyal A, Richter KR, et al. Clinical utility of brain biopsy for presumed CNS relapse of systemic lymphoma. *J Neurosurg.* Published online July 2, 2021. doi:10.3171/2020.12.JNS202517). We commend the authors for sharing their valuable experience on the management of patients with systemic lymphoma. In this retrospective study, the authors concluded that brain biopsy offered little diagnostic help but carried high risks of morbidity and mortality in patients with a history of systemic lymphoma and presumed cerebral relapse. In

this clinical situation, they advocate for empirical treatment instead of obtaining histological confirmation with a brain biopsy.

Based on our experience and findings, we would like to offer our view and contribution regarding the place and role of brain biopsy in the management of systemic lymphoma patients. First, because of their immunocompromised status induced by the disease itself and/or antineoplastic therapies, patients with systemic lymphoma may have a large variety of diseases affecting the brain (opportunistic infections, inflammatory diseases, and other types of neoplasia).² The frequency and diversity of neurological diseases in this patient subgroup render brain biopsies highly informative.³ Moreover, cerebral dual pathologies concern 5% of immunocompromised patients.⁴

Second, cerebral relapse must be histologically proven for inclusion in most of the currently ongoing clinical trials for patients with relapsed systemic lymphoma (e.g., NCT04608487, NCT01148173). In this context, performing a brain biopsy is critical to enabling patients to benefit from the latest medical developments.

For the reasons given above, it is obvious that histological confirmation of cerebral relapse is needed in most cases. Nevertheless, we acknowledge that lymphomas, regardless of their subtype, have a higher tendency for postbiopsy bleeding than other brain lesions,^{5,6} which may lead to potentially severe complications.⁷ However, in our series of 177 consecutive patients who underwent a stereotactic brain biopsy that provided the diagnosis of cerebral lymphoma (among 1646 patients who underwent brain biopsy between April 2009 and February 2020), postbiopsy complications rates remained low: we noted a 4.5% rate of symptomatic complications and no biopsy-related deaths.⁵ Moreover, the use of 5-aminolevulinic acid (5-ALA),⁸ intraoperative histological examination using frozen section or smear,⁹ and flow cytometry analysis of brain biopsy rinse fluid¹⁰ may prevent taking too many tissue samples during stereotactic biopsy and thereby may decrease the postbiopsy morbidity rate. We strongly believe that it is the neurosurgeon's duty is to make brain biopsies as safe as possible so that the indications for biopsy are not limited by the potential risks associated with the procedure. Regardless, the risk of complications of a brain biopsy should always be weighed against the risk of misdiagnosis and therefore inappropriate treatment. In most cases, the latter is often more life-threatening than the former.

In our view, the balance between the benefits and risks of brain biopsy in patients with presumed cerebral relapse of systemic lymphoma should be systematically assessed and weighed on a case-by-case basis in a multidisciplinary discussion involving a neurosurgeon, with consideration of the patient's medical history, clinical status, treatment plan, and location of the targeted lesion.

Bertrand Mathon, MD, MSc^{1,2}

Aymeric Amelot, MD, PhD^{1,3}

On behalf of the PSL BRAIN-BIOPSY Study Group

¹La Pitié-Salpêtrière Hospital, APHP, Sorbonne University, Paris, France

²Paris Brain Institute, ICM, INSERM U 1127, CNRS UMR 7225, Sorbonne University, Paris, France

³Bretonneau Hospital, Tours, France

References

1. Brown DA, Goyal A, Richter KR, et al. Clinical utility of brain biopsy for presumed CNS relapse of systemic lymphoma. *J Neurosurg*. Published online July 2, 2021. doi: 10.3171/2020.12.JNS202517
2. Recht LD. Neurologic complications of systemic lymphoma. *Neurol Clin*. 1991;9(4):1001-1015.
3. Mathon B, Le Joncour A, Bielle F, et al. Neurological diseases of unknown etiology: brain-biopsy diagnostic yields and safety. *Eur J Intern Med*. 2020;80:78-85.
4. Vitelli M, Malaizé H, Bielle F, et al. A diagnosis can hide another: the value of brain biopsy in neurological lesion of HIV patients. *J Acquir Immune Defic Syndr*. 2021;86(1):e6-e9.
5. Mathon B, Marijon P, Riche M, Amelot A; PSL Brain-Biopsy Study Group. Letter to the editor regarding: "Hemorrhagic attitude in frameless and frame-based stereotactic biopsy for deep-seated primary central nervous system." *World Neurosurg*. 2021;152:242-243.
6. Riche M, Marijon P, Amelot A, et al. Severity, timeline and management of complications after stereotactic brain biopsy. *J Neurosurg*. Published online September 10, 2021. doi: 10.3171/2021.3.JNS21134
7. Riche M, Amelot A, Peyre M, Capelle L, Carpentier A, Mathon B. Complications after frame-based stereotactic brain biopsy: a systematic review. *Neurosurg Rev*. 2021;44(1):301-307.
8. Kiesel B, Millesi M, Woehrer A, et al. 5-ALA-induced fluorescence as a marker for diagnostic tissue in stereotactic biopsies of intracranial lymphomas: experience in 41 patients. *Neurosurg Focus*. 2018;44(6):E7.
9. Mathon B, Amelot A, Mokhtari K, Bielle F. Increasing the diagnostic yield of stereotactic brain biopsy using intraoperative histological smear. *Clin Neurol Neurosurg*. 2019;186:105544.
10. Deblieux A, Voirin J, Harzallah I, et al. Cytomorphology and flow cytometry of brain biopsy rinse fluid enables faster and multidisciplinary diagnosis of large B-cell lymphoma of the central nervous system. *Cytometry B Clin Cytom*. 2018;94(1):182-188.

Disclosures

The authors report no conflict of interest.

Correspondence

Bertrand Mathon: bertrand.mathon@aphp.fr.

INCLUDE WHEN CITING

Published online November 5, 2021; DOI: 10.3171/2021.7.JNS211699.

Response

No response was received from the authors of the original article.

©AANS 2022, except where prohibited by US copyright law

Outcome prediction and assessment after surgery in CM-I: there is more to it than meets the eye

TO THE EDITOR: We read with great interest the article by Feghali et al.¹ published in the *Journal of Neurosurgery* (Feghali J, Xie Y, Chen Y, et al. External validation of current prediction systems of improvement after decom-

pression surgery in Chiari malformation type I patients: can we do better? *J Neurosurg*. 2021;134[5]:1466-1471). We congratulate the authors for their paper but would like to add some insights based on our experience.

We believe that the Chiari Severity Index (CSI)² and Thakar et al.'s point score³ are not flawless. The CSI talks about headaches and myelopathy clinically, whereas in real life, many patients with Chiari malformation type I (CM-I) will have paresthesia, numbness, and small hand muscle atrophy only. This led us to propose new clinical disease severity criteria in our paper.⁴ The "minimally symptomatic" category comprised patients with headache or neck pain with or without mild paresthesia that was not bothersome. On the contrary, the presence of severe paresthesia, motor symptoms (myelopathy), and atrophy of the hands with or without dissociative anesthesia were considered as "clinically severe" disease.⁴ Radiologically, the CSI talks about the presence or absence of syringomyelia, and when present it comprises syrinx thickness as a measure of disease severity.² However, we measured the vertical extent of the syrinx as an alternative method.⁴ The latter is more easily visible and measurable in most clinical situations. Moreover, we combined the extent of tonsillar descent in deriving the radiological disease severity estimation, which is the essence of CM-I diagnosis, easily measurable, and not included in the CSI. Thus, we agree with the authors when they say that there is a need for developing better preoperative clinoradiological predicting systems for CM-I.¹ Similarly, the parameters proposed by Thakar et al.³ may not be reproducible in the everyday busy clinical setting.

We also wish to comment on the Chicago Chiari Outcome Scale (CCOS).⁵ We have used this system in our study,⁴ believing firmly that our efforts to assess outcomes should be as objective as possible. However, we noted some outliers in this system, also pointed out by Aliaga et al.⁵ in their original article. It underestimated the improvement related to surgery in the short term by nearly 25%.⁴ We inferred that in the CCOS system (a 4-component score), even if the overall score is 11 or more (indicating an improvement), a low score in any specific domain thought to be important by the patient may lead him/her to report an unimproved or a worse outcome. Unlike other studies, we also looked at the long-term CCOS score and found that the CCOS cutoff (> 11) overestimated the favorable outcome, despite late deteriorations in our 4 patients.⁴ Therefore, the CCOS score clearly has some limitations. We explored the difference in the CCOS score between the time of discharge and at follow-up. We statistically demonstrated that a minimum score difference of 2 between the scores correlated well with patient-perceived outcome in the long run.⁴ This new matrix could capture surgical outcomes closer to the real-life patient perceptions.

Therefore, there is definitely more to this than what we know as of now. We once again congratulate the authors and hope the additional insights from us will strengthen their points of view for the readers.

Kuntal Kanti Das, MCh
Sanjay Gandhi Postgraduate Institute of
Medical Sciences, Lucknow, India

Jaskaran Singh Gosal, MCh, DNB

All India Institute of Medical Sciences (AIIMS), Jodhpur, India

Deepak Khatri, MCh

Lenox-Hill Hospital, New York, NY

Arun K. Srivastava, MCh

Sanjay Gandhi Postgraduate Institute of
Medical Sciences, Lucknow, India

References

1. Feghali J, Xie Y, Chen Y, et al. External validation of current prediction systems of improvement after decompression surgery in Chiari malformation type I patients: can we do better? *J Neurosurg.* 2021;134(5):1466-1471.
2. Greenberg JK, Yarbrough CK, Radmanesh A, et al. The Chiari Severity Index: a preoperative grading system for Chiari malformation type 1. *Neurosurgery.* 2015;76(3):279-285.
3. Thakar S, Sivaraju L, Jacob KS, et al. A points-based algorithm for prognosticating clinical outcome of Chiari malformation Type I with syringomyelia: results from a predictive model analysis of 82 surgically managed adult patients. *J Neurosurg Spine.* 2018;28(1):23-32.
4. Rangari K, Das KK, Singh S, et al. Type I Chiari malformation without concomitant bony instability: assessment of different surgical procedures and outcomes in 73 patients. *Neurospine.* 2021;18(1):126-138.
5. Aliaga L, Hekman KE, Yassari R, et al. A novel scoring system for assessing Chiari malformation type I treatment outcomes. *Neurosurgery.* 2012;70:656-665.

Disclosures

The authors report no conflict of interest.

Correspondence:

Kuntal Kanti Das: drkuntalkantidas@gmail.com.

INCLUDE WHEN CITING

Published online October 15, 2021; DOI: 10.3171/2021.5.JNS211329.

Response

We thank Das et al. for their insightful comments. As our external validation study demonstrates, predictive accuracy of both tools was limited in predicting surgical outcomes of CM-I as measured by the CCOS. Das et al. point out one possible reason, which is that these scores may neglect other variables that could enhance predictive accuracy. In our single-institution experience, we identified additional variables that demonstrate potential for predicting outcome, such as odontoid retroflexion, race, and visual symptoms, but external validation efforts are still needed.¹ The publication of online web-based calculator platforms has the potential to facilitate external validation efforts by independent investigators.

The outcome scale of interest is also an important point that Das et al. stress. The CSI, for instance, was developed to predict outcome on a brief quality-of-life questionnaire, unlike the CCOS, which is designed to capture outcome based on a retrospective chart review.² This represents another potential contributing factor to the poor performance of the CSI on external validation. Although our study evaluated the CSI in adult patients (different than the pediatric population of the original study), Ahluwalia et al. similarly demonstrated poor performance of the CSI in predicting

surgical outcome in 65 pediatric patients as measured by an abbreviated version of the CCOS.³

Based on the published literature regarding surgical outcome prediction of CM-I, much remains to be answered. Recently, an increasing emphasis has been placed on cranial morphometric measurements and on machine learning approaches to data interpretation and modeling. A catalog summarizing the important morphometric variables that have been studied represents a useful tool in planning and structuring future studies. The use of multiinstitutional data and the publication of testable models (e.g., calculators, scoring systems, decision trees, etc.) that encourage external validation efforts will be crucial steps moving forward. Predictive modeling should also attempt to discern which operative techniques (e.g., arachnoid dissection, tonsillectomy, duraplasty, cranioplasty) have the greatest chance of producing satisfactory outcomes depending on the patient's baseline presentation characteristics.

James Feghali, MD

Johns Hopkins University School of Medicine, Baltimore, MD

Sean Li

Pratt School of Engineering, Duke University, Durham, NC

Judy Huang, MD

Johns Hopkins University School of Medicine, Baltimore, MD

References

1. Feghali J, Xie Y, Chen Y, et al. The SHORE score: a novel predictive tool for improvement after decompression surgery in adult Chiari malformation type I. *World Neurosurg.* 2020; 142:e195-e202.
2. Greenberg JK, Limbrick DD. The Chiari Severity Index and the role of external validation. Letter. *J Neurosurg Pediatr.* 2021;27(2):241-242.
3. Ahluwalia R, Foster J, Brooks E, et al. Chiari type I malformation: role of the Chiari Severity Index and Chicago Chiari Outcome Scale. *J Neurosurg Pediatr.* 2020;26(3):262-268.

INCLUDE WHEN CITING

Published online October 15, 2021; DOI: 10.3171/2021.6.JNS211343.

©AANS 2022, except where prohibited by US copyright law

Overcoming obstacles and breaking barriers for women in neurosurgery

TO THE EDITOR: More women attend medical school than men.¹ Yet, female representation in neurosurgery remains low.² Donaldson and colleagues³ highlight the imbalance between the numbers of male and female residents and faculty members in neurosurgical training programs, offering further evidence of an ongoing discrepancy between the number of women who become physicians and the number who successfully train to become neurosurgeons; we aim to supplement these efforts (Donaldson K, Callahan KE, Gelinne A, et al. Gender diversity in United States neurosurgery training programs. *J Neurosurg.* 2021;135[3]:943-948).

The relatively small number of women in neurosurgery

reflects the “pipeline effect,” which suggests that gender differences within the neurosurgical field can be attributed to fewer females entering the discipline and a consequential delayed elevation in rank.³ However, the evidence does not support this explanation. Although female representation in neurosurgery has continuously increased since the 1990s, it has done so more slowly and gradually than other heavily male-dominated medical fields.^{2,4} Indeed, neurosurgery remains one of the specialties with the lowest female representation.³ On the contrary, fields such as general surgery have seen substantial gains in gender equity⁵ in recent years. To promote quality of patient care and advance the field, we need to attract, support, and encourage women to become neurosurgeons.

Understanding how formerly male-dominated fields such as business have addressed underrepresentation of women can be beneficial. In corporate leadership, gender bias and feelings of loneliness are recognized as factors that limit women’s advancement,⁶ and many women have argued that their experiences as some of the few females working in male-dominated spheres have negatively impacted their successes.³ This is to say that as more and more women enter the field of neurosurgery, their doing so may also increase the odds of succeeding generations of women entering the specialty. Particularly within a field that prides itself on the “see one, do one, teach one” approach, women interested in neurosurgery must be able to model their pursuits after women who have entered the field and succeeded before them.

With a greater number of role models, women will be better equipped to face the challenges of pursuing a career in neurosurgery.⁷ As the number of women in neurosurgery continues to increase, we expect to see changes in the next decade, including an emphasis on hiring diverse faculty, expanded options for parental leave and on-site day care, a focus on inviting female neurosurgeons as keynote speakers, equal compensation as men, and zero-tolerance harassment policies.² Additionally, as other fields have found that the solution to curtailing gender bias and reducing gender discrimination lies in providing more equitable educational and professional environments,^{4,7–10} we anticipate that the formation of more organizations such as Women in Neurosurgery (WINS) may have an impact on recruitment of women to neurosurgery.

Chidinma M. Wilson, BA, BS

Evalyn S. Mackenzie, BA

Mikhail A. Yudien, BA

Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA

Antoinette J. Charles, BS

Duke University School of Medicine, Durham, NC

Marianne I. J. Tissot, BA

Sydney J. Churchill, BS

Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA

Nolan J. Brown, BS

University of California, Irvine, Orange, CA

Jared M. Shulkin, AB

Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA

Donald K. E. Detchou, BA

Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA
Hospital of the University of Pennsylvania, Philadelphia, PA

Vamsi P. Reddy, MD

UT Southwestern Medical Center, Dallas, TX

Lola B. Chambless, MD

Vanderbilt University Medical Center, Nashville, TN

References

1. Boyle P. More women than men are enrolled in medical school. AAMC; 2019. Accessed August 6, 2021. <https://www.aamc.org/news-insights/more-women-men-are-enrolled-medical-school>
2. Abosch A, Rutka JT. Women in neurosurgery: inequality redux. *J Neurosurg*. 2018;129(2):277-281.
3. Donaldson K, Callahan KE, Gelinne A, et al. Gender diversity in United States neurosurgery training programs. *J Neurosurg*. 2021;135(3):943-948.
4. Corley J, Kim E, Philips CA, et al. One hundred years of neurosurgery: contributions of American women. *J Neurosurg*. 2021;134(2):337-342.
5. Linscheid LJ, Holliday EB, Ahmed A, et al. Women in academic surgery over the last four decades. *PLoS One* 2020; 15(12):e0243308.
6. Berry P, Franks TJ. Women in the world of corporate business: looking at the glass ceiling. *Contemp Issues Educ Res (Littleton)*. 2010;3(2):1-10.
7. Renfrow JJ, Rodriguez A, Wilson TA, Germano IM, Abosch A, Wolfe SQ. Tracking career paths of women in neurosurgery. *Neurosurgery*. 2018;82(4):576-582.
8. Patel EA, Aydin A, Cearns M, Dasgupta P, Ahmed K. A systematic review of simulation-based training in neurosurgery, part 1: cranial neurosurgery. *World Neurosurg*. 2020;133:e850-e873.
9. Howard SD, Lucas TH. The Supreme Court’s recent decision is a call for increased diversity in neurosurgery. *Letter. Neurosurgery*. 2020;87(6):E715-E716.
10. Lubelski D, Xiao R, Mukherjee D, et al. Improving medical student recruitment to neurosurgery. *J Neurosurg*. 2020; 133(3):848-854.

Disclosures

The authors report no conflict of interest.

Correspondence

Donald K. E. Detchou: donald.detchou@pennmedicine.upenn.edu.

Response

No response was received from the authors of the original article.

INCLUDE WHEN CITING

Published online October 15, 2021; DOI: 10.3171/2021.5.JNS211269.

©AANS 2022, except where prohibited by US copyright law