Thalamomesencephalic cavernoma: anterior transcallosal transchoroidal approach

Francesco Certo, MD, Giada Toccaceli, MD, Roberto Altieri, MD, and Giuseppe M. V. Barbagallo, MD

Department of Neurological Surgery, Policlinico "G. Rodolico" University Hospital, Catania, Italy

We present a case of a 62-year-old man with acute onset of diplopia, headache, and vomiting for a bleeding thalamomesencephalic cavernoma. The lesion was removed via the anterior transcallosal transchoroidal approach. His head was slightly flexed and a right paramedian craniotomy for an interhemispheric approach was performed. The interhemispheric fissure was split and, after callosotomy, the choroidal fissure was opened along the tenia fornicis to enter the velum interpositum and enlarge the foramen of Monro. The cavernoma was then identified and resected. There were no long-term postoperative neurological deficits. This approach is a valid alternative for thalamomesencephalic lesions.

The video can be found here: <https://youtu.be/DJdorbzDnH0>.

KEYWORDS cavernoma; third ventricle; transcallosal; approach; video

Transcript

We present a case of thalamomesencephalic cavernoma removed by anterior transcallosal transchoroidal approach.

The patient is a 62-year-old man who presented, after mild brain injury, acute onset of diplopia, headache, and vomiting. Because of persistence of symptoms, he performed a brain CT that showed a right thalamomesencephalic bleeding, with a maximum diameter of 12 mm.

For these reasons, patient was admitted in Neurosurgery Department and performed an MRI scan that showed a right periaqueductal gray mesencephalic and thalamic cavernoma and the presence of a minimal intraventricular bleeding.

1:29 Mesencephalic cavernous malformations (Lee and Lui, 1996; Garrett and Spetzler, 2009; Beechar et al., 2017; Ren et al., 2017) are usually approached via:

Anterior approaches (ventral cavernomas)

- Orbitozygomatic or frontotemporal-orbitozygomatic approaches
- Subtemporal approach
- Pterional transsylvian approach

Posterior approaches (dorsal cavernomas)

- Supracerebellar infratentorial approach
- Occipital transtentorial/interhemispheric

Superior approaches (ventral cavernomas surfacing the floor of third ventricle)

- Transcallosal transchoroidal
- Transcallosal transforaminal

For the reported case, an anterior route should be advised (OZ, FTOZ, pterional), but these approaches are associated with increased risks of:

- Pyramidal tract injury
- Mechanical or vascular injuries

Superior surface of cavernoma was directly visible in the context of the floor of third ventricle.

Due to the symptomatic onset with bleeding, we decide to operate the patient and we choose the anterior transcallosal transchoroidal approach.

The patient was in anti-Trendelenburg supine position with the head slightly flexed. We performed a C-shaped skin incision and the craniotomy is tailored with respect to the angle of approach.

SUBMITTED March 1, 2019. **ACCEPTED** May 13, 2019.

INCLUDE WHEN CITING Published online July 1, 2019; DOI: <http://thejns.org/doi/abs/10.3171/2019.7.FocusVid.191213>

© 2019, Francesco Certo, Giada Toccaceli, Roberto Altieri, and Giuseppe M. V. Barbagallo, CC BY 4.0 (<http://creativecommons.org/licenses/by/4.0/>)

We used, as surgical tools, intraoperative CT, neuro-navigation, and neurophysiological monitoring.

In this image we can see both the AxiEM neuronavigation and the radiolucent Mayfield head holder, useful to perform the intraoperative CT scan.

After coagulating and dissecting some bridging veins we proceeded to open the interhemispheric fissure, detect and separate pericallosal arteries, and find the corpus callosum.

3:18 In this phase, neuronavigation is useful to be guided in the surgical trajectory.

3:21 After callosotomy we could enter in the right lateral ventricle, opening a wide fissure.

3:38 The dissection of the choroidal fissure to release the choroid plexus from the fornix must be performed extremely carefully.

So, having identified the choroid plexus, the thalamostriate veins, and, anteriorly, the Monro foramen, we proceeded to a careful dissection of choroidal fissure, to release the choroid plexus from the fornix, avoiding any kind of injury.

4:03 Identification of thalamostriate veins and foramen of Monro

The choroidal fissure was opened along the tenia fornicea to enter the velum interpositum and enlarge the foramen of Monro

We entered in the third ventricle and we detected the cavernoma. We checked the position with the neuronavigation and then we proceeded to remove it.

4:37 Excision of cavernoma

In this phase neuromonitoring is useful to avoid neurological damages.

5:35 Hemostasis

6:16 Watertight dural closure

Postoperative neuroimaging showed the complete resection of cavernoma with minimal cerebral manipulation.

In conclusion, thalamomesencephalic cavernomas continue to represent a considerable microsurgical challenge. Even if most ventral thalamomesencephalic cavernomas can be removed via transylvian, subtemporal, or frontotemporal-zygomatic approach, the anterior transcallosal transchoroidal approach can serve as valuable alternative surgical corridor.

References

1. Beechar VB, Srinivasan VM, Reznik OE, Sen A, Klisch TJ, Ropper AE: Intraventricular cavernomas of the third ventricle: report of 2 cases and a systematic review of the literature. **World Neurosurg** **105**:935–943, 943.e1–943.e3, 2017
2. Garrett M, Spetzler RF: Surgical treatment of brainstem cavernous malformations. **Surg Neurol** **72** (Suppl 2):S3–S10, 2009
3. Hosainey SA, Meling TR: A 34-year-old woman with brainstem cavernous malformation: the anterior transcallosal transchoroidal approach and literature review. **J Neurol Surg Rep** **75**:e236–e240, 2014
4. Lee CC, Lui TN: Mesencephalic cavernous hemangioma. **J Formos Med Assoc** **95**:493–495, 1996
5. Ren Y, Li J, Tao C, Zheng J, Zhang S, Xiao A: Surgical treatment of cavernous malformations involving the midbrain: a single-center case series of 34 patients. **World Neurosurg** **107**:753–763, 2017
6. Ulm AJ, Russo A, Albanese E, Tanriover N, Martins C, Mericle RM: Limitations of the transcallosal transchoroidal approach to the third ventricle. **J Neurosurg** **111**:600–609, 2009

Correspondence

Giada Toccaceli, Policlinico “G. Rodolico” University Hospital, Catania, Italy. toccaceli.giada@gmail.com.

Disclosures

The authors report no conflict of interest concerning the materials or methods used in this study or the findings specified in this article.