Defining postoperative values for successful resection of prolactinomas

Clinical pearl

William T. Couldwell, M.D., Ph.D., and Martin H. Weiss, M.D.

*Departments of Surgery and Neuroscience, University of North Dakota, Minot, North Dakota; and
Department of Neurosurgery, University of Southern California, Los Angeles, California*

Key Words * hyperprolactinemia * microadenoma

We, as other authors, have noted that late relapse of hyperprolactinemia following apparent successful adenomectomy in women harboring microprolactinomas is not rare.[1-3]

On retrospective review of a series of tumors treated at the University of Southern California teaching hospitals, we have found that the immediate postoperative prolactin level may be predictive of late recurrence of hyperprolactinemia.[4] It is our routine to ascertain prolactin levels at 3 days and 6 weeks postoperatively. Of 133 patients undergoing microadenomectomy who had 3-day postoperative prolactin levels of less than 10 $\mu\text{g/L}$ (the vast majority of these were less than 3 $\mu\text{g/L}$), 132 or 99% had prolactin levels of less than 20 $\mu\text{g/L}$ at their 6-week follow-up examination. At 5 years postoperatively, 130 (98%) of the original 133 patients still had normal prolactin levels. However, of 43 patients who had 3-day postoperative prolactin levels in the range of 10 to 20 $\mu\text{g/L}$, only 32 (74%) had prolactin levels in the normal range at 6 weeks postoperatively, and only five (16%) of this group of 32 patients had normal prolactin levels at 5-year follow-up review. Therefore, although 18% of patients who have a normal postoperative prolactin level at 6 weeks will develop recurrent hyperprolactinemia when followed for 5 years, only three (7%) of 41 represent patients with immediate prolactin levels of less than 10 $\mu\text{g/L}$, whereas the remainder were in the group with immediate postoperative prolactin levels in the 10 to 20 $\mu\text{g/L}$ range.

These data indicate that the traditional criteria used to determine successful microadenoma resection in the immediate postoperative period (prolactin level of less than 20 $\mu\text{g/L}$) may be too liberal; in our series the patient with an immediate postoperative prolactin of less than 10 $\mu\text{g/L}$ had an approximately 98% chance of long-term chemical cure, and we have yet to see a recurrence of hyperprolactinemia in a patient whose immediate postoperative prolactin level was less than 3 $\mu\text{g/L}$.

References

1. Rodman EF, Molitch ME, Post KD, et al: Long-term follow-up of transsphenoidal selective adenomectomy for prolactinoma. **JAMA** **252**:921-924, 1984
 2. Serri O, Hardy J, Massoud F: Relapse of hyperprolactinemia revisited. **N Engl J Med** **329**:1357, 1993 (Letter)
 3. Serri O, Rasio E, Beauregard H, et al.: Recurrence of hyperprolactinemia after selective transsphenoidal adenomectomy in women with prolactinoma. **N Engl J Med** **309**:280-283, 1983
 4. Weiss MH: Role of surgical intervention for prolactinomas, in Melmed S, Robbins RJ, (eds): **Molecular and Clinical Advances in Pituitary Disorders**. Blackwell Scientific Publications, 1991, pp 255-260
-

Manuscript received June 3, 1996.

Accepted in final form June 10, 1996.

Address reprint requests to: William T. Couldwell, M.D., Ph.D., Trinity Medical Center, One Burdick Expressway West, Minot, North Dakota 58701.