Letter to the editor

Bullock MR, Merchant RE, and Choi SC, et al: Outcome measures for clinical trials in neurotrauma. **Neurosurg Focus** 13 (1):Clinical Pearl 1, 2002

TO THE EDITOR:

Although the review reported by Bullock and colleagues of the current state of outcome measures in neurotrauma² is quite comprehensive, its final receipt date (January 2002) has resulted in inadvertent omission of two recent developments in this rapidly changing field. Both of these relate to the use of oxidative stress as a measure of severity of neurotrauma, based on the excitotoxicity and free radical hypothesis.³ In the first of these developments,⁴ published in March 2002, the authors described a technique in which magnetic resonance spectroscopy was used for noninvasive measurement of nitric oxide synthase activity, a key focus of interest in the design of neuroprotective pharmacological agents. In the second,¹ published in May 2002, the investigators described the measurement of cerebrospinal fluid free radicals and antioxidant reserve as markers of the severity of oxidative stress. As noted by Bullock and colleagues, "Future research should be focused on development of sensitive and specific surrogate outcome measures that can be applied to Phase II trials and used to detect a neuroprotective effect." Undoubtedly, the closer the relationship between these measures and the pathophysiology of neural injury—as in oxidative stress—the greater the likelihood of achieving a significant response to pharmacological antagonists to the mechanism of injury.

MORRY SILBERSTEIN, M.D.

Assistant Dean, Academic Programs
Monash Medical School
Victoria, Australia

References

1. Bayir H, Kagan VE, Tyurina YY, et al: Assessment of antioxidant reserves and oxidative stress in cerebrospinal fluid after severe traumatic brain injury in infants and children. **Pediatr Res** 51:571–578, 2002
2. Bullock MR, Merchant RE, Choi SC, et al. Outcome measures for clinical trials in neurotrauma. **Neurosurg Focus** 13 (1):Clinical Pearl 1, 2002
3. Gilgun-Sherki Y, Rosenbaum Z, Melamed E, et al: Antioxidant therapy in acute central nervous system injury: current state. **Pharmacol Rev** 54:271–284, 2002
4. Silberstein M, Lane D, Dodd S, et al: Identification of a by-product of nitric oxide synthase activity in human acute brain injury with in vivo proton magnetic resonance spectroscopy. **AJNR Am J Neuroradiol** 23:389–392, 2002

RESPONSE:

We thank Dr. Silberstein for his comments regarding our manuscript, "Outcome measures for clinical trials in neurotrauma." We agree completely with him that novel outcome measures based on the excitotoxicity and free radical hypotheses are important and interesting.

The definition of a surrogate outcome measure is that it should very closely correlate with the actual outcome, which is best portrayed by one of the outcome scales such as the Glasgow Outcome Scale, Functional Independence Measure, and Disability Rating Scale.

As yet, the measures, which are described in the two references noted by Dr. Silberstein, have not been validated for use in human traumatic brain injury.

We absolutely agree, however, that such validation studies should be performed to assess these and other surrogate outcome measures. Our own institution has been dedicated to the evaluation of outcome measures such as brain oxygen tension, dialysate glutamate, and lactate glucose ratio, but perhaps the most cohesive evidence in support of a chemical "surrogate outcome measure" rests with S100 at this time. This aspect is discussed in the manuscript.

Again, we thank Dr. Silberstein for his comments

M. ROSS BULLOCK, M.D., PH.D.

Reynolds Professor
Neurosurgery and Anesthesiology
Virginia Commonwealth University
Richmond, Virginia