Minimally invasive transforaminal lumbosacral interbody fusion

Peng-Yuan Chang, MD, and Michael Y. Wang, MD

Departments of Neurosurgery and Rehabilitation Medicine, University of Miami Miller School of Medicine, Miami, Florida

In minimally invasive spinal fusion surgery, transforaminal lumbar (sacral) interbody fusion (TLIF) is one of the most common procedures that provides both anterior and posterior column support without retraction or violation to the neural structure. Direct and indirect decompression can be done through this single approach. Preoperative plain radiographs and MR scan should be carefully evaluated. This video demonstrates a standard approach for how to perform a minimally invasive transforaminal lumbosacral interbody fusion.

The video can be found here: <https://youtu.be/bhEeafKJ370>.

KEYWORDS minimally invasive surgery; MIS; TLIF; expandable cage; percutaneous screw; video

SUBMITTED March 1, 2016. **ACCEPTED** April 12, 2016.

INCLUDE WHEN CITING Published online July 1, 2016; DOI: <http://thejns.org/doi/abs/10.3171/2016.2.FocusVid.1694>.

CORRESPONDENCE Michael Y. Wang, Department of Neurosurgery, University of Miami Miller School of Medicine, Lois Pope Life Center, 1095 Northwest 14th Terrace, Miami, FL 33136. email: mwang2@med.miami.edu.