Awake craniotomy for right inferior lobule glioma

Rafael T. M. Leal, MD, and Jose A. Landeiro, MD, PhD

Department of Neurosurgery, Universidade Federal Fluminense, Hospital Universitário Antonio Pedro, Niterói, Rio de Janeiro, Brazil

Awake craniotomy is a method usually used for brain tumors near or within language and/or sensorimotor areas. Preservation of these important neurological functions, however, may not be enough for a normal life. The authors present a case of a 45-year-old woman with an inferior parietal lobule low-grade glioma who underwent an awake craniotomy with sensorimotor and spatial cognition mapping.

The video can be found here: https://youtu.be/HMA_pZYg3Ms.

KEYWORDS inferior parietal lobule; awake craniotomy; glioma; video

SUBMITTED July 15, 2018. **ACCEPTED** August 9, 2018.

INCLUDE WHEN CITING Published online October 1, 2018; DOI: <http://thejns.org/doi/abs/10.3171/2018.10.FocusVid.18275>.

DISCLOSURES The authors report no conflict of interest concerning the materials or methods used in this study or the findings specified in this paper.

CORRESPONDENCE Rafael T. M. Leal: Hospital Universitário Antonio Pedro, Niterói, Rio de Janeiro, Brazil. rafael_tmleal@yahoo.com.br.